Nai-Liang Tian

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10105213/publications.pdf Version: 2024-02-01



NALLIANC TIAN

#	Article	IF	CITATIONS
1	Impact of the Complexity of Bifurcation Lesions Treated With Drug-Eluting Stents. JACC: Cardiovascular Interventions, 2014, 7, 1266-1276.	2.9	153
2	3-Year Outcomes of the ULTIMATE TrialÂComparing Intravascular Ultrasound Versus Angiography-Guided Drug-Eluting Stent Implantation. JACC: Cardiovascular Interventions, 2021, 14, 247-257.	2.9	149
3	Angiographic and clinical comparisons of intravascular ultrasound- versus angiography-guided drug-eluting stent implantation for patients with chronic total occlusion lesions: two-year results from a randomised AIR-CTO study. EuroIntervention, 2015, 10, 1409-1417.	3.2	139
4	Multicentre, randomized comparison of two-stent and provisional stenting techniques in patients with complex coronary bifurcation lesions: the DEFINITION II trial. European Heart Journal, 2020, 41, 2523-2536.	2.2	124
5	Cutoff Value and Long-Term Prediction ofÂClinical Events by FFR Measured Immediately After Implantation of a Drug-Eluting Stent in Patients With Coronary Artery Disease. JACC: Cardiovascular Interventions, 2017, 10, 986-995.	2.9	111
6	Incidence and Clinical Outcomes ofÂStentÂFractures on the Basis of 6,555ÂPatientsÂandÂ16,482 Drug-Eluting StentsÂFrom 4ÂCenters. JACC: Cardiovascular Interventions, 2016, 9, 1115-1123.	2.9	66
7	Stenting strategy for coronary artery bifurcation with drug-eluting stents: a meta-analysis of nine randomised trials and systematic review. EuroIntervention, 2014, 10, 561-569.	3.2	66
8	Comparison of one-year clinical outcomes between intravascular ultrasound-guided versus angiography-guided implantation of drug-eluting stents for left main lesions: a single-center analysis of a 1,016-patient cohort. Patient Preference and Adherence, 2014, 8, 1299.	1.8	43
9	MicroRNA-210 promotes angiogenesis in acute myocardial infarction. Molecular Medicine Reports, 2018, 17, 5658-5665.	2.4	39
10	Intravascular ultrasound-guided drug-eluting stent implantation is associated with improved clinical outcomes in patients with unstable angina and complex coronary artery true bifurcation lesions. International Journal of Cardiovascular Imaging, 2018, 34, 1685-1696.	1.5	34
11	<p>Comparisons between ticagrelor and clopidogrel following percutaneous coronary intervention in patients with acute coronary syndrome: a comprehensive meta-analysis</p> . Drug Design, Development and Therapy, 2019, Volume 13, 719-730.	4.3	21
12	The Acute Changes of Fractional Flow Reserve in DK (Double Kissing), Crush, and 1‣tent Technique for True Bifurcation Lesions. Journal of Interventional Cardiology, 2010, 23, 341-345.	1.2	19
13	Serial intravascular ultrasound analysis comparing double kissing and classical crush stenting for coronary bifurcation lesions. Catheterization and Cardiovascular Interventions, 2011, 78, 729-736.	1.7	18
14	The impact of everolimus versus other rapamycin derivative-eluting stents on clinical outcomes in patients with coronary artery disease: A meta-analysis of 16 randomized trials. Journal of Cardiology, 2014, 64, 185-193.	1.9	15
15	High platelet reactivity affects the clinical outcomes of patients undergoing percutaneous coronary intervention. BMC Cardiovascular Disorders, 2016, 16, 240.	1.7	15
16	The clinical outcomes of triple antiplatelet therapy versus dual antiplatelet therapy for high-risk patients after coronary stent implantation: a meta-analysis of 11 clinical trials and 9,553 patients. Drug Design, Development and Therapy, 2016, Volume 10, 3435-3448.	4.3	14
17	Obstructive sleep apnea affects the clinical outcomes of patients undergoing percutaneous coronary intervention. Patient Preference and Adherence, 2016, 10, 871.	1.8	13
18	Comparison of intravascular ultrasound-guided with angiography-guided double kissing crush stenting for patients with complex coronary bifurcation lesions: Rationale and design of a prospective, randomized, and multicenter DKCRUSH VIII trial. American Heart Journal, 2021, 234, 101-110.	2.7	12

NAI-LIANG TIAN

#	Article	IF	CITATIONS
19	Treatment effects of systematic two-stent and provisional stenting techniques in patients with complex coronary bifurcation lesions: rationale and design of a prospective, randomised and multicentre DEFINITION II trial. BMJ Open, 2018, 8, e020019.	1.9	7
20	Obstructive Sleep Apnea Affecting Platelet Reactivity in Patients Undergoing Percutaneous Coronary Intervention. Chinese Medical Journal, 2018, 131, 1023-1029.	2.3	7
21	Stent fracture is associated with a higher mortality in patients with type-2 diabetes treated by implantation of a second-generation drug-eluting stent. International Journal of Cardiovascular Imaging, 2017, 33, 1873-1881.	1.5	5
22	The outcomes of intra-aortic balloon pump usage in patients with acute myocardial infarction: a comprehensive meta-analysis of 33 clinical trials and 18,889 patients. Patient Preference and Adherence, 2016, 10, 297.	1.8	4
23	Post-dilatation improves stent apposition in patients with ST-segment elevation myocardial infarction receiving primary percutaneous intervention: A multicenter, randomized controlled trial using optical coherence tomography. World Journal of Emergency Medicine, 2020, 11, 87.	1.0	3
24	NOBORIâ,"¢ biodegradable-polymer biolimus-eluting stent versus durable-polymer drug-eluting stents: A meta-analysis. International Journal of Cardiology, 2014, 174, 151-153.	1.7	2
25	The Clinical Outcomes of Ventricular Septal Rupture Secondary to Acute Myocardial Infarction: A Retrospective, Observational Trial. Journal of Interventional Cardiology, 2021, 2021, 1-7.	1.2	2
26	Maintained P2Y ₁₂ inhibitor monotherapy after shorterâ€duration of dual antiplatelet therapy in patients undergoing coronary drugâ€eluting stents implantation: An updated metaâ€analysis of randomized trials. Journal of Clinical Pharmacy and Therapeutics, 2022, , .	1.5	0