Min-Jian Qin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1019637/publications.pdf

Version: 2024-02-01

304743 395702 1,313 61 22 33 citations h-index g-index papers 65 65 65 1850 all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	An integrated study of Violae Herba (Viola philippica) and five adulterants by morphology, chemical compositions and chloroplast genomes: insights into its certified plant origin. Chinese Medicine, 2022, 17, 32.	4.0	5
2	Influence of different pretreatments and drying methods on the chemical compositions and bioactivities of Smilacis Glabrae Rhizoma. Chinese Medicine, 2022, 17, 54.	4.0	5
3	Iris domestica (iso)flavone 7- and 3′-O-Glycosyltransferases Can Be Induced by CuCl2. Frontiers in Plant Science, 2021, 12, 632557.	3.6	3
4	The Effect of Hispidulin, a Flavonoid from Salvia plebeia, on Human Nasopharyngeal Carcinoma CNE-2Z Cell Proliferation, Migration, Invasion, and Apoptosis. Molecules, 2021, 26, 1604.	3.8	4
5	The complete chloroplast genome of a Chinese medicinal plant, <i>Peristrophe japonica</i> (Thunb.) Bremek. (Lamiales: Acanthaceae) from Nanjing, China. Mitochondrial DNA Part B: Resources, 2021, 6, 1888-1889.	0.4	2
6	The complete chloroplast genome of Clerodendrum lindleyi Decne. ex Planch. (Tubiflorae:) Tj ETQq0 0 0 rgBT /O	verlock 10	Tf 50 542 Td
7	Chemical constituents and antioxidative, anti-inflammatory and anti-proliferative activities of wild and cultivated Corydalis saxicola. Industrial Crops and Products, 2021, 169, 113647.	5.2	12
8	A New Arylbenzofuran Derivative from Mentha canadensis. Chemistry of Natural Compounds, 2021, 57, 44-46.	0.8	0
9	Copper stress-induced changes in biomass accumulation, antioxidant activity and flavonoid contents in Belamcanda chinensis calli. Plant Cell, Tissue and Organ Culture, 2020, 142, 299-311.	2.3	15
10	Separation of acteoside and linarin from Buddlejae Flos by highâ€speed countercurrent chromatography and their antiâ€inflammatory activities. Journal of Separation Science, 2020, 43, 1450-1457.	2.5	20
11	Dynamic analysis of secondary metabolites in various parts of Scrophularia ningpoensis by liquid chromatography tandem mass spectrometry. Journal of Pharmaceutical and Biomedical Analysis, 2020, 186, 113307.	2.8	15
12	Chemical profiles and quality evaluation of Buddleja officinalis flowers by HPLC-DAD and HPLC-Q-TOF-MS/MS. Journal of Pharmaceutical and Biomedical Analysis, 2019, 164, 283-295.	2.8	35
13	Molecular diversity analysis of Tetradium ruticarpum (WuZhuYu) in China based on inter-primer binding site (iPBS) markers and inter-simple sequence repeat (ISSR) markers. Chinese Journal of Natural Medicines, 2018, 16, 1-9.	1.3	20
14	Qualitative and Quantitative Analysis of C-glycosyl-flavones of Iris lactea Leaves by Liquid Chromatography/Tandem Mass Spectrometry. Molecules, 2018, 23, 3359.	3.8	10
15	Global Transcriptome Analyses Reveal Differentially Expressed Genes of Six Organs and Putative Genes Involved in (Iso)flavonoid Biosynthesis in Belamcanda chinensis. Frontiers in Plant Science, 2018, 9, 1160.	3.6	9
16	Organ-Specific Metabolic Shifts of Flavonoids in Scutellaria baicalensis at Different Growth and Development Stages. Molecules, 2018, 23, 428.	3.8	33
17	Baicalin regulates SirT1/STAT3 pathway and restrains excessive hepatic glucose production. Pharmacological Research, 2018, 136, 62-73.	7.1	29
18	Determination of ginsenosides in Asian and American ginsengs by liquid chromatography–quadrupole/time-of-flight MS: assessing variations based on morphological characteristics. Journal of Ginseng Research, 2017, 41, 10-22.	5.7	38

#	Article	IF	Citations
19	Alkaloids from the Rhizomes of Iris germanica. Chemistry of Natural Compounds, 2017, 53, 196-198.	0.8	6
20	Antihyperglycemic, antihyperlipidemic and antioxidant effects of standard ethanol extract of Bombax ceiba leaves in high-fat-diet- and streptozotocin-induced Type 2 diabetic rats. Chinese Journal of Natural Medicines, 2017, 15, 168-177.	1.3	27
21	Four new C -glycosylflavones from the leaves of Iris lactea Pall. var . chinensis (Fisch.) Koidz Phytochemistry Letters, 2017, 22, 33-38.	1.2	10
22	Optimization of the Extraction Conditions for Buddleja officinalis Maxim. Using Response Surface Methodology and Exploration of the Optimum Harvest Time. Molecules, 2017, 22, 1877.	3.8	10
23	Optimization of the Ultrasonic-Assisted Extraction of Bioactive Flavonoids from Ampelopsis grossedentata and Subsequent Separation and Purification of Two Flavonoid Aglycones by High-Speed Counter-Current Chromatography. Molecules, 2016, 21, 1096.	3.8	25
24	Simultaneous Determination of Catalpol, Aucubin, and Geniposidic Acid in Different Developmental Stages of <i>Rehmannia glutinosa </i> Leaves by High Performance Liquid Chromatography. Journal of Analytical Methods in Chemistry, 2016, 2016, 1-6.	1.6	64
25	Chemical Constituents of Pyrrosia calvata. Natural Product Communications, 2015, 10, 1934578X1501000.	0.5	1
26	Effects of drying methods on the phytochemicals contents and antioxidant properties of chrysanthemum flower heads harvested at two developmental stages. Journal of Functional Foods, 2015, 19, 786-795.	3.4	56
27	Tissue-specific metabolite profiling and quantitative analysis of ginsenosides in Panax quinquefolium using laser microdissection and liquid chromatography–quadrupole/time of flight-mass spectrometry. Chemistry Central Journal, 2015, 9, 66.	2.6	9
28	Cause and control of Radix Ophiopogonis browning during storage. Chinese Journal of Natural Medicines, 2015, 13, 73-80.	1.3	3
29	Localization of ginsenosides in the rhizome and root of Panax ginseng by laser microdissection and liquid chromatography–quadrupole/time of flight-mass spectrometry. Journal of Pharmaceutical and Biomedical Analysis, 2015, 105, 121-133.	2.8	44
30	New flavonoids with cytotoxicity from the roots of Flemingia latifolia. Fìtoterapìâ, 2015, 104, 97-101.	2.2	9
31	Multiple responses optimization of ultrasonic-assisted extraction by response surface methodology (RSM) for rapid analysis of bioactive compounds in the flower head of Chrysanthemum morifolium Ramat Industrial Crops and Products, 2015, 74, 192-199.	5.2	68
32	Chemical Differentiation and Quality Evaluation of Commercial Asian and American Ginsengs based on a UHPLC–QTOF/MS/MS Metabolomics Approach. Phytochemical Analysis, 2015, 26, 145-160.	2.4	49
33	Four new eudesmane-type sesquiterpenes from the basal leaves of Salvia plebeia R. Br Fìtoterapìâ, 2014, 94, 142-147.	2.2	20
34	Stilbenoids from the seeds of Oroxylum indicum. Biochemical Systematics and Ecology, 2014, 54, 36-39.	1.3	3
35	Flavonoids from Mentha haplocalyx. Chemistry of Natural Compounds, 2014, 50, 124-125.	0.8	4
36	Phenolic metabolite profiles and antioxidants assay of three Iridaceae medicinal plants for traditional Chinese medicine "She-gan―by on-line HPLC–DAD coupled with chemiluminescence (CL) and ESI-Q-TOF-MS/MS. Journal of Pharmaceutical and Biomedical Analysis, 2014, 98, 40-51.	2.8	49

#	Article	IF	Citations
37	Dynamic Changes of Flavonoids Contents in the Different Parts of Rhizome of Belamcanda chinensis During the Thermal Drying Process. Molecules, 2014, 19, 10440-10454.	3.8	41
38	The Spectrum-Effect integrated fingerprint of Polygonum cus-pidatum based on HPLC-diode array detection-flow injection-chemiluminescence. Chinese Journal of Natural Medicines, 2014, 11, 546-552.	1.3	12
39	Application of an efficient strategy for discovery and purification of bioactive compounds from Chinese herbal medicines, a case study on the Puerariae thomsonii Flos. Journal of Pharmaceutical and Biomedical Analysis, 2013, 75, 25-32.	2.8	17
40	Chemical constituents from Mentha canadensis. Biochemical Systematics and Ecology, 2013, 49, 144-147.	1.3	10
41	The Spectrum-Effect integrated fingerprint of Polygonum cuspidatum based on HPLC-diode array detection-flow injection-chemiluminescence. Chinese Journal of Natural Medicines, 2013, 11, 546-552.	1.3	7
42	Kakkalide ameliorates endothelial insulin resistance by suppressing reactive oxygen speciesâ€associated inflammation (eʻʻ花跷通过æŠʻ制活性氧ç¸å³çš"ç,Žç—‡æ"¹å−"å†çš®ç»†èƒžèƒ°å²¸ç´æŠµæŠ—). Journa	al ¹ 68Diabe	etes, 2013, 5
43	New isoflavones with cytotoxic activity from the rhizomes of <i>lris germanica </i> L Natural Product Research, 2013, 27, 2173-2177.	1.8	22
44	Tectorigenin Attenuates Palmitate-Induced Endothelial Insulin Resistance via Targeting ROS-Associated Inflammation and IRS-1 Pathway. PLoS ONE, 2013, 8, e66417.	2.5	31
45	C-glycosylflavones from the leaves of Iris tectorum Maxim Acta Pharmaceutica Sinica B, 2012, 2, 598-601.	12.0	15
46	Qualitative and Quantitative Evaluation of Phenolic Compounds in <i>Iris dichotoma</i> Pall. Phytochemical Analysis, 2012, 23, 197-207.	2.4	21
47	Phylogenetic study of Oryzoideae species and related taxa of the Poaceae based on atpB-rbcL and ndhF DNA sequences. Molecular Biology Reports, 2012, 39, 5737-5744.	2.3	13
48	Authentication of an endangered herb Changium smyrnioides from different producing areas based on rDNA ITS sequences and allele-specific PCR. Archives of Pharmacal Research, 2012, 35, 701-708.	6.3	4
49	A new coumestan with immunosuppressive activities from Flemingia philippinensis. Fìtoterapìâ, 2011, 82, 615-619.	2.2	30
50	A new belamcandaquinone from the seeds of Iris bungei Maxim Fìtoterapìâ, 2011, 82, 1137-1139.	2,2	14
51	Comparative study of isoflavones in wild and cultivated soybeans as well as bean products by high-performance liquid chromatography coupled with mass spectrometry and chemometric techniques. European Food Research and Technology, 2011, 233, 869-880.	3.3	20
52	Characterization and determination of the major constituents in Belamcandae Rhizoma by HPLC–DAD–ESI-MSn. Journal of Pharmaceutical and Biomedical Analysis, 2011, 56, 304-314.	2.8	66
53	Characterization of homoisoflavonoids in different cultivation regions of Ophiopogon japonicus and related antioxidant activity. Journal of Pharmaceutical and Biomedical Analysis, 2010, 52, 757-762.	2.8	51
54	New features on the fragmentation patterns of homoisoflavonoids in ⟨i⟩Ophiopogon japonicus⟨ i⟩ by highâ€performance liquid chromatography/diodeâ€array detection/electrospray ionization with multiâ€stage tandem mass spectrometry. Rapid Communications in Mass Spectrometry, 2010, 24, 2193-2206.	1.5	19

#	Article	IF	CITATION
55	Analysis of catalpol derivatives by characteristic neutral losses using liquid chromatography combined with electrospray ionization multistage and timeâ€ofâ€flight mass spectrometry. Rapid Communications in Mass Spectrometry, 2010, 24, 2680-26863.	1.5	22
56	Analysis of Flavonoids and Phenolic Acids in <l>lris tectorum</l> by HPLC-DAD-ESI-MS ⁿ . Chinese Journal of Natural Medicines, 2010, 8, 202-207.	1.3	34
57	Chemical Constituents of <i>Viola yedoensis</i> . Chinese Journal of Natural Medicines, 2009, 7, 290-292.	1.3	10
58	Improved quality evaluation of Radix Salvia miltiorrhiza through simultaneous quantification of seven major active components by high-performance liquid chromatography and principal component analysis. Biomedical Chromatography, 2007, 21, 931-939.	1.7	39
59	New Isoflavonoid Glycosides from the Rhizomes of Iris leptophylla Lingelsh Journal of Integrative Plant Biology, 2007, 49, 213-217.	8.5	4
60	A New Isoflavonoid from Belamcanda chinensis (L.) DC Journal of Integrative Plant Biology, 2005, 47, 1404-1408.	8.5	17
61	Diterpenoids from the Rhizomes of Alpinia calcarata. Journal of Natural Products, 2000, 63, 939-942.	3.0	58