
## Alexandra Mant

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1038103/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Bone marrow transplantation for MHC class I deficiency corrects T-cell immunity but dissociates natural killer cell repertoire formation from function. Journal of Allergy and Clinical Immunology, 2016, 138, 1733-1736.e2.	2.9	7
2	Acid sensitive background potassium channels K2P3.1 and K2P9.1 undergo rapid dynamin-dependent endocytosis. Channels, 2013, 7, 288-292.	2.8	4
3	N-Glycosylation-dependent Control of Functional Expression of Background Potassium Channels K2P3.1 and K2P9.1. Journal of Biological Chemistry, 2013, 288, 3251-3264.	3.4	20
4	The pathway of crossâ€presentation is influenced by the particle size of phagocytosed antigen. Immunology, 2012, 136, 163-175.	4.4	52
5	Protein Kinase A Is Central for Forward Transport of Two-pore Domain Potassium Channels K2P3.1 and K2P9.1. Journal of Biological Chemistry, 2011, 286, 14110-14119.	3.4	21
6	Polymer microarrays: Identification of substrates for phagocytosis assays. Biomaterials, 2006, 27, 5299-5306.	11.4	40
7	The Properties of the Positively Charged Loop Region in PSI-G Are Essential for Its "Spontaneous― Insertion into Thylakoids and Rapid Assembly into the Photosystem I Complex. Journal of Biological Chemistry, 2006, 281, 10548-10554.	3.4	17
8	Insertion of the plant photosystem I subunit G into the thylakoid membrane. FEBS Journal, 2005, 272, 4002-4010.	4.7	13
9	Â-Glucan, water dikinase (GWD): A plastidic enzyme with redox-regulated and coordinated catalytic activity and binding affinity. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 1785-1790.	7.1	131
10	Arabidopsis VARIEGATED 3 encodes a chloroplast-targeted, zinc-finger protein required for chloroplast and palisade cell development. Journal of Cell Science, 2004, 117, 4807-4818.	2.0	65
11	Identification of an Arabidopsis inorganic pyrophosphatase capable of being imported into chloroplasts. FEBS Letters, 2004, 565, 101-105.	2.8	30
12	Three Isoforms of Isoamylase Contribute Different Catalytic Properties for the Debranching of Potato Glucans[W]. Plant Cell, 2003, 15, 133-149.	6.6	161
13	PSI-O, a new 10-kDa subunit of eukaryotic photosystem I. FEBS Letters, 2002, 510, 145-148.	2.8	52
14	Functional Characterization of Recombinant Chloroplast Signal Recognition Particle. Journal of Biological Chemistry, 2001, 276, 27778-27786.	3.4	70
15	Insertion of PsaK into the Thylakoid Membrane in a "Horseshoe―Conformation Occurs in the Absence of Signal Recognition Particle, Nucleoside Triphosphates, or Functional Albino3. Journal of Biological Chemistry, 2001, 276, 36200-36206.	3.4	50
16	Distinct Albino3-dependent and -independent Pathways for Thylakoid Membrane Protein Insertion. Journal of Biological Chemistry, 2001, 276, 40841-40846.	3.4	80
17	Conformation of a Purified "Spontaneously―Inserting Thylakoid Membrane Protein Precursor in Aqueous Solvent and Detergent Micelles. Journal of Biological Chemistry, 2001, 276, 14607-14613.	3.4	9
18	BUNDLE SHEATH DEFECTIVE2, a Novel Protein Required for Post-Translational Regulation of the rbcL Gene of Maize. Plant Cell, 1999, 11, 849-864.	6.6	149

ALEXANDRA MANT

#	Article	IF	CITATIONS
19	Distinct "Assisted―and "Spontaneous―Mechanisms for the Insertion of Polytopic Chlorophyll-binding Proteins into the Thylakoid Membrane. Journal of Biological Chemistry, 1999, 274, 4715-4721.	3.4	39
20	Dual Signal Peptides Mediate the Signal Recognition Particle/Sec-independent Insertion of a Thylakoid Membrane Polyprotein, PsbY. Journal of Biological Chemistry, 1999, 274, 4059-4066.	3.4	29
21	Characterisation of an Arabidopsis thaliana cDNA encoding a novel thylakoid lumen protein imported by the ΔpH-dependent pathway. Planta, 1999, 207, 624-627.	3.2	7
22	Multiple pathways for the targeting of thylakoid proteins in chloroplasts. Plant Molecular Biology, 1998, 38, 209-221.	3.9	58
23	AnArabidopsiscDNA encodes an apparent polyprotein of two non-identical thylakoid membrane proteins that are associated with photosystem II and homologous to algalycf32open reading frames. FEBS Letters, 1998, 423, 183-188.	2.8	23
24	Sec/SRP-independent insertion of two thylakoid membrane proteins bearing cleavable signal peptides. FEBS Letters, 1998, 424, 105-108.	2.8	38
25	Characterisation of anArabidopsiscDNA encoding a thylakoid lumen protein related to a novel `pentapeptide repeat' family of proteins. FEBS Letters, 1998, 428, 241-244.	2.8	26
26	Multiple pathways for the targeting of thylakoid proteins in chloroplasts. , 1998, , 209-221.		3
27	Targeting of proteins into and across the thylakoid membrane. Trends in Plant Science, 1997, 2, 431-437.	8.8	49
28	A Monomeric, Tightly Folded Stromal Intermediate on the â—³pH-dependent Thylakoidal Protein Transport Pathway. Journal of Biological Chemistry, 1995, 270, 1663-1669.	3.4	67
29	Determination of the unfrozen water content of maximally freeze-concentrated carbohydrate solutions. International Journal of Biological Macromolecules, 1993, 15, 227-232.	7.5	50