

Wei Xuan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1038392/publications.pdf>

Version: 2024-02-01

10
papers

454
citations

1307594

7
h-index

1372567

10
g-index

10
all docs

10
docs citations

10
times ranked

844
citing authors

#	ARTICLE	IF	CITATIONS
1	Inhaler reminders improve adherence with controller treatment in primary care patients with asthma. <i>Journal of Allergy and Clinical Immunology</i> , 2014, 134, 1260-1268.e3.	2.9	198
2	Effect of Inpatient Rehabilitation vs a Monitored Home-Based Program on Mobility in Patients With Total Knee Arthroplasty. <i>JAMA - Journal of the American Medical Association</i> , 2017, 317, 1037.	7.4	110
3	Independent Echocardiographic Markers of Cardiovascular Involvement in Chronic Kidney Disease: The Value of Left Atrial Function and Volume. <i>Journal of the American Society of Echocardiography</i> , 2016, 29, 359-367.	2.8	56
4	Patient satisfaction and perception of success after total knee arthroplasty are more strongly associated with patient factors and complications than surgical or anaesthetic factors. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 3156-3163.	4.2	32
5	Exercise testing can unmask right ventricular dysfunction in systemic sclerosis patients with normal resting pulmonary artery pressure. <i>International Journal of Cardiology</i> , 2016, 204, 179-186.	1.7	19
6	The effectiveness of inpatient rehabilitation after uncomplicated total hip arthroplasty: a propensity score matched cohort. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 236.	1.9	13
7	Changes in Right Ventricular Function with Exercise in Healthy Subjects: Optimal Parameters and Effects of Gender and Age. <i>Journal of the American Society of Echocardiography</i> , 2015, 28, 1441-1451.e1.	2.8	12
8	Timing and duration of antibiotic prophylaxis is associated with the risk of infection after hip and knee arthroplasty. <i>Bone & Joint Open</i> , 2022, 3, 252-260.	2.6	6
9	Patient factors associated with weight gain and weight loss after knee or hip arthroplasty. <i>Obesity Research and Clinical Practice</i> , 2019, 13, 371-377.	1.8	5
10	Evidence-Based Determination of Cut-Off Points for Increased Cardiac-Surgery Mortality Risk With EuroSCORE II and STS: The Best-Performing Risk Scoring Models in a Single-Centre Australian Population. <i>Heart Lung and Circulation</i> , 2022, 31, 590-601.	0.4	3