

Urban Åsby

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10386659/publications.pdf>

Version: 2024-02-01

55
papers

5,977
citations

186265

28
h-index

168389

53
g-index

60
all docs

60
docs citations

60
times ranked

8982
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Genome-wide association study identifies 30 loci associated with bipolar disorder. <i>Nature Genetics</i> , 2019, 51, 793-803. | 21.4 | 1,191 |
| 2 | Excess Mortality in Bipolar and Unipolar Disorder in Sweden. <i>Archives of General Psychiatry</i> , 2001, 58, 844. | 12.3 | 951 |
| 3 | Mortality and causes of death in schizophrenia in Stockholm County, Sweden. <i>Schizophrenia Research</i> , 2000, 45, 21-28. | 2.0 | 551 |
| 4 | Excess Mortality, Causes of Death and Life Expectancy in 270,770 Patients with Recent Onset of Mental Disorders in Denmark, Finland and Sweden. <i>PLoS ONE</i> , 2013, 8, e55176. | 2.5 | 486 |
| 5 | Genetic variants associated with response to lithium treatment in bipolar disorder: a genome-wide association study. <i>Lancet</i> , The, 2016, 387, 1085-1093. | 13.7 | 306 |
| 6 | Life Expectancy and Death by Diseases of the Circulatory System in Patients with Bipolar Disorder or Schizophrenia in the Nordic Countries. <i>PLoS ONE</i> , 2013, 8, e67133. | 2.5 | 250 |
| 7 | Evaluation of diagnostic procedures in Swedish patients with schizophrenia and related psychoses. <i>Nordic Journal of Psychiatry</i> , 2005, 59, 457-464. | 1.3 | 195 |
| 8 | Genome-wide association study of 40,000 individuals identifies two novel loci associated with bipolar disorder. <i>Human Molecular Genetics</i> , 2016, 25, 3383-3394. | 2.9 | 182 |
| 9 | Time trends in schizophrenia mortality in Stockholm County, Sweden: cohort study. <i>BMJ: British Medical Journal</i> , 2000, 321, 483-484. | 2.3 | 170 |
| 10 | Assessment of Response to Lithium Maintenance Treatment in Bipolar Disorder: A Consortium on Lithium Genetics (ConLiGen) Report. <i>PLoS ONE</i> , 2013, 8, e65636. | 2.5 | 156 |
| 11 | Cardiovascular mortality in bipolar disorder: a population-based cohort study in Sweden. <i>BMJ Open</i> , 2013, 3, e002373. | 1.9 | 154 |
| 12 | The Genetics of the Mood Disorder Spectrum: Genome-wide Association Analyses of More Than 185,000 Cases and 439,000 Controls. <i>Biological Psychiatry</i> , 2020, 88, 169-184. | 1.3 | 137 |
| 13 | Coeliac disease and risk of mood disorders – A general population-based cohort study. <i>Journal of Affective Disorders</i> , 2007, 99, 117-126. | 4.1 | 106 |
| 14 | Association of Polygenic Score for Schizophrenia and HLA Antigen and Inflammation Genes With Response to Lithium in Bipolar Affective Disorder. <i>JAMA Psychiatry</i> , 2018, 75, 65-74. | 11.0 | 102 |
| 15 | Mortality trends in cardiovascular causes in schizophrenia, bipolar and unipolar mood disorder in Sweden 1987–2010. <i>European Journal of Public Health</i> , 2016, 26, 867-871. | 0.3 | 77 |
| 16 | CRY2 Is Associated with Rapid Cycling in Bipolar Disorder Patients. <i>PLoS ONE</i> , 2010, 5, e12632. | 2.5 | 71 |
| 17 | Risk factors for suicide in schizophrenia: Findings from a Swedish population-based case-control study. <i>Schizophrenia Research</i> , 2009, 108, 231-237. | 2.0 | 70 |
| 18 | Carriers of the UGT1A4 142T>G gene variant are predisposed to reduced olanzapine exposure – an impact similar to male gender or smoking in schizophrenic patients. <i>European Journal of Clinical Pharmacology</i> , 2010, 66, 465-474. | 1.9 | 58 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Coeliac disease and risk of schizophrenia and other psychosis: A general population cohort study. <i>Scandinavian Journal of Gastroenterology</i> , 2007, 42, 179-185. | 1.5 | 49 |
| 20 | Suicide and hospitalization for mental disorders in Sweden: A population-based case-control study. <i>Journal of Psychiatric Research</i> , 2010, 44, 741-747. | 3.1 | 48 |
| 21 | Lithium treatment and cancer incidence in bipolar disorder. <i>Bipolar Disorders</i> , 2016, 18, 33-40. | 1.9 | 48 |
| 22 | Association of polygenic score for major depression with response to lithium in patients with bipolar disorder. <i>Molecular Psychiatry</i> , 2021, 26, 2457-2470. | 7.9 | 44 |
| 23 | Cognitive manic symptoms associated with the <i>P2RX7</i> gene in bipolar disorder. <i>Bipolar Disorders</i> , 2011, 13, 500-508. | 1.9 | 43 |
| 24 | Swedish clinical guidelines"Prevention and management of metabolic risk in patients with severe psychiatric disorders. <i>Nordic Journal of Psychiatry</i> , 2010, 64, 294-302. | 1.3 | 39 |
| 25 | Medication and suicide risk in schizophrenia: A nested case"control study. <i>Schizophrenia Research</i> , 2013, 150, 416-420. | 2.0 | 37 |
| 26 | Serotonergic medication enhances the association between suicide and sunshine. <i>Journal of Affective Disorders</i> , 2016, 189, 276-281. | 4.1 | 35 |
| 27 | P2RX7: Expression Responds to Sleep Deprivation and Associates with Rapid Cycling in Bipolar Disorder Type 1. <i>PLoS ONE</i> , 2012, 7, e43057. | 2.5 | 35 |
| 28 | Mitochondrial DNA copy number is associated with psychosis severity and anti-psychotic treatment. <i>Scientific Reports</i> , 2018, 8, 12743. | 3.3 | 34 |
| 29 | Time trends in first admissions for schizophrenia and paranoid psychosis in Stockholm County, Sweden. <i>Schizophrenia Research</i> , 2001, 47, 247-254. | 2.0 | 29 |
| 30 | Analysis of the Influence of microRNAs in Lithium Response in Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2018, 9, 207. | 2.6 | 28 |
| 31 | Combining schizophrenia and depression polygenic risk scores improves the genetic prediction of lithium response in bipolar disorder patients. <i>Translational Psychiatry</i> , 2021, 11, 606. | 4.8 | 25 |
| 32 | A comparison of cardiovascular risk factors for ten antipsychotic drugs in clinical practice. <i>Neuropsychiatric Disease and Treatment</i> , 2013, 9, 371. | 2.2 | 20 |
| 33 | Investigating polygenic burden in age at disease onset in bipolar disorder: Findings from an international multicentric study. <i>Bipolar Disorders</i> , 2019, 21, 68-75. | 1.9 | 20 |
| 34 | Mortality trends in external causes of death in people with mental health disorders in Sweden, 1987"2010. <i>Scandinavian Journal of Public Health</i> , 2019, 47, 121-126. | 2.3 | 20 |
| 35 | Suicide risk and antipsychotic side effects in schizophrenia: nested case-control study. <i>Human Psychopharmacology</i> , 2016, 31, 341-345. | 1.5 | 14 |
| 36 | Diagnostic profile and suicide risk in schizophrenia spectrum disorder. <i>Schizophrenia Research</i> , 2010, 123, 251-256. | 2.0 | 13 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Genetic and Clinical Factors Affecting Plasma Clozapine Concentration. primary care companion for CNS disorders, The, 2015, 17, . | 0.6 | 13 |
| 38 | Health-related quality of life and metabolic risk in patients with psychosis. Schizophrenia Research, 2014, 152, 295-299. | 2.0 | 12 |
| 39 | Association between history of psychosis and cardiovascular disease in bipolar disorder. Bipolar Disorders, 2015, 17, 518-527. | 1.9 | 12 |
| 40 | Depression-associated <i>ARNTL</i> and <i>PER2</i> genetic variants in psychotic disorders. Chronobiology International, 2015, 32, 579-584. | 2.0 | 12 |
| 41 | Suicide Ideation and Behavior as Risk Factors for Subsequent Suicide in Schizophrenia: A Nested Caseâ€“Control Study. Suicide and Life-Threatening Behavior, 2019, 49, 996-1005. | 1.9 | 12 |
| 42 | Gender influence on the bipolar disorder inpatient length of stay in Sweden, 2005â€“2014: A register-based study. Journal of Affective Disorders, 2019, 256, 183-191. | 4.1 | 12 |
| 43 | Psychotic disorder is an independent risk factor for increased fasting glucose and waist circumference. Nordic Journal of Psychiatry, 2014, 68, 251-258. | 1.3 | 11 |
| 44 | Risperidone metabolic ratio as a biomarker of individual CYP2D6 genotype in schizophrenic patients. European Journal of Clinical Pharmacology, 2014, 70, 695-699. | 1.9 | 10 |
| 45 | HLA-DRB1 and HLA-DQB1 genetic diversity modulates response to lithium in bipolar affective disorders. Scientific Reports, 2021, 11, 17823. | 3.3 | 10 |
| 46 | Cognitive Manic Symptoms in Bipolar Disorder Associated with Polymorphisms in the DAOA and COMT Genes. PLoS ONE, 2013, 8, e67450. | 2.5 | 7 |
| 47 | Diabetes and glucose disturbances in patients with psychosis in Sweden. BMJ Open Diabetes Research and Care, 2015, 3, e000120. | 2.8 | 6 |
| 48 | Outcome of a psychosocial health promotion intervention aimed at improving physical health and reducing alcohol use in patients with schizophrenia and psychotic disorders (MINT). Schizophrenia Research, 2019, 208, 138-144. | 2.0 | 6 |
| 49 | Plasma GDF15 level is elevated in psychosis and inversely correlated with severity. Scientific Reports, 2017, 7, 7906. | 3.3 | 5 |
| 50 | Genetic variants of increased waist circumference in psychosis. Psychiatric Genetics, 2017, 27, 210-218. | 1.1 | 4 |
| 51 | Symptoms and treatment of bipolar patients in Sweden. International Journal of Psychiatry in Clinical Practice, 2012, 16, 170-177. | 2.4 | 3 |
| 52 | Melatonin receptor 1B gene associated with hyperglycemia in bipolar disorder. Psychiatric Genetics, 2016, 26, 136-139. | 1.1 | 3 |
| 53 | Troponin T levels associated with genetic variants in NOTCH2 and MTNR1B in women with psychosis. Psychiatry Research, 2017, 250, 217-220. | 3.3 | 2 |
| 54 | The general health of people with schizophrenia. , 2012, , 91-106. | | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|--|----|-----------|
| 55 | The general health of people with schizophrenia. , 2012, , 91-106. | | 0 |