

Jeremy A Goettel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10412922/publications.pdf>

Version: 2024-02-01

19
papers

1,635
citations

567281

15
h-index

839539

18
g-index

19
all docs

19
docs citations

19
times ranked

3668
citing authors

#	ARTICLE	IF	CITATIONS
1	HLA-Restriction of Human Treg Cells Is Not Required for Therapeutic Efficacy of Low-Dose IL-2 in Humanized Mice. <i>Frontiers in Immunology</i> , 2021, 12, 630204.	4.8	12
2	Intestinal Regulatory T Cells as Specialized Tissue-Restricted Immune Cells in Intestinal Immune Homeostasis and Disease. <i>Frontiers in Immunology</i> , 2021, 12, 716499.	4.8	34
3	Utilizing a reductionist model to study host-microbe interactions in intestinal inflammation. <i>Microbiome</i> , 2021, 9, 215.	11.1	8
4	Humanized mouse models of genetic immune disorders and hematological malignancies. <i>Biochemical Pharmacology</i> , 2020, 174, 113671.	4.4	5
5	Low-Dose Interleukin-2 Ameliorates Colitis in a Preclinical Humanized Mouse Model. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2019, 8, 193-195.	4.5	25
6	WASP-mediated regulation of anti-inflammatory macrophages is IL-10 dependent and is critical for intestinal homeostasis. <i>Nature Communications</i> , 2018, 9, 1779.	12.8	40
7	Haematopoietic stem and progenitor cells from human pluripotent stem cells. <i>Nature</i> , 2017, 545, 432-438.	27.8	395
8	Enhanced TH17 Responses in Patients with IL10 Receptor Deficiency and Infantile-onset IBD. <i>Inflammatory Bowel Diseases</i> , 2017, 23, 1950-1961.	1.9	28
9	Macrophage dysfunction initiates colitis during weaning of infant mice lacking the interleukin-10 receptor. <i>ELife</i> , 2017, 6, .	6.0	26
10	AHR Activation Is Protective against Colitis Driven by T Cells in Humanized Mice. <i>Cell Reports</i> , 2016, 17, 1318-1329.	6.4	147
11	FOXP3+ Tregs require WASP to restrain Th2-mediated food allergy. <i>Journal of Clinical Investigation</i> , 2016, 126, 4030-4044.	8.2	53
12	Fatal autoimmunity in mice reconstituted with human hematopoietic stem cells encoding defective FOXP3. <i>Blood</i> , 2015, 125, 3886-3895.	1.4	33
13	Interleukin-10 Receptor Signaling in Innate Immune Cells Regulates Mucosal Immune Tolerance and Anti-Inflammatory Macrophage Function. <i>Immunity</i> , 2014, 40, 706-719.	14.3	455
14	Interleukin 10 Receptor Signaling. <i>Advances in Immunology</i> , 2014, 122, 177-210.	2.2	239
15	Colitis and Colon Cancer in WASP-Deficient Mice Require Helicobacter Species. <i>Inflammatory Bowel Diseases</i> , 2013, 19, 2041-2050.	1.9	31
16	Wiskott-Aldrich Syndrome Protein Deficiency in Innate Immune Cells Leads to Mucosal Immune Dysregulation and Colitis in Mice. <i>Gastroenterology</i> , 2012, 143, 719-729.e2.	1.3	32
17	TNF transactivation of EGFR stimulates cytoprotective COX-2 expression in gastrointestinal epithelial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 301, G220-G229.	3.4	39
18	TNFR1 Promotes Tumor Necrosis Factor-mediated Mouse Colon Epithelial Cell Survival through RAF Activation of NF- κ B. <i>Journal of Biological Chemistry</i> , 2008, 283, 29485-29494.	3.4	32

#	ARTICLE	IF	CITATIONS
19	Hematopoietic stem/progenitor cell conversion from human pluripotent stem cells. Protocol Exchange, 0, , .	0.3	1