

# Yang Yu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10584657/publications.pdf>

Version: 2024-02-01

45  
papers

1,552  
citations

257450

24  
h-index

315739

38  
g-index

47  
all docs

47  
docs citations

47  
times ranked

3403  
citing authors

#	ARTICLE	IF	CITATIONS
1	BX-795 inhibits neuroblastoma growth and enhances sensitivity towards chemotherapy. <i>Translational Oncology</i> , 2022, 15, 101272.	3.7	9
2	A genome-scale CRISPR Cas9 dropout screen identifies synthetically lethal targets in SRC-3 inhibited cancer cells. <i>Communications Biology</i> , 2021, 4, 399.	4.4	8
3	Mechanisms restraining break-induced replication at two-ended DNA double-strand breaks. <i>EMBO Journal</i> , 2021, 40, e104847.	7.8	45
4	Development of improved SRC-3 inhibitors as breast cancer therapeutic agents. <i>Endocrine-Related Cancer</i> , 2021, 28, 657-670.	3.1	7
5	Drug-induced PD-L1 expression and cell stress response in breast cancer cells can be balanced by drug combination. <i>Scientific Reports</i> , 2019, 9, 15099.	3.3	40
6	Rad52 Restrains Resection at DNA Double-Strand Break Ends in Yeast. <i>Molecular Cell</i> , 2019, 76, 699-711.e6.	9.7	37
7	Small molecule inhibitor agerafenib effectively suppresses neuroblastoma tumor growth in mouse models via inhibiting ERK MAPK signaling. <i>Cancer Letters</i> , 2019, 457, 129-141.	7.2	16
8	Delanzomib, a novel proteasome inhibitor, sensitizes breast cancer cells to doxorubicin-induced apoptosis. <i>Thoracic Cancer</i> , 2019, 10, 918-929.	1.9	6
9	Inhibition of Ubiquitin-Specific Protease 14 Suppresses Cell Proliferation and Synergizes with Chemotherapeutic Agents in Neuroblastoma. <i>Molecular Cancer Therapeutics</i> , 2019, 18, 1045-1056.	4.1	11
10	SRC-3 inhibition blocks tumor growth of pancreatic ductal adenocarcinoma. <i>Cancer Letters</i> , 2019, 442, 310-319.	7.2	17
11	Dna2 nuclease deficiency results in large and complex DNA insertions at chromosomal breaks. <i>Nature</i> , 2018, 564, 287-290.	27.8	33
12	Targeting LRH-1 in hepatoblastoma cell lines causes decreased proliferation. <i>Oncology Reports</i> , 2018, 41, 143-153.	2.6	14
13	Proteomic profiling identifies key coactivators utilized by mutant ER $\alpha$ proteins as potential new therapeutic targets. <i>Oncogene</i> , 2018, 37, 4581-4598.	5.9	51
14	MELK is a novel therapeutic target in high-risk neuroblastoma. <i>Oncotarget</i> , 2018, 9, 2591-2602.	1.8	22
15	Machine Learning Applications in Head and Neck Radiation Oncology: Lessons From Open-Source Radiomics Challenges. <i>Frontiers in Oncology</i> , 2018, 8, 294.	2.8	37
16	The second-generation ALK inhibitor alectinib effectively induces apoptosis in human neuroblastoma cells and inhibits tumor growth in a TH-MYCN transgenic neuroblastoma mouse model. <i>Cancer Letters</i> , 2017, 400, 61-68.	7.2	37
17	Targeting SRC Coactivators Blocks the Tumor-Initiating Capacity of Cancer Stem-like Cells. <i>Cancer Research</i> , 2017, 77, 4293-4304.	0.9	36
18	Yeast Sub1 and human PC4 are G-quadruplex binding proteins that suppress genome instability at co-transcriptionally formed G4 DNA. <i>Nucleic Acids Research</i> , 2017, 45, 5850-5862.	14.5	41

#	ARTICLE	IF	CITATIONS
19	EWS-FLI1 and RNA helicase A interaction inhibitor YK-4-279 inhibits growth of neuroblastoma. <i>Oncotarget</i> , 2017, 8, 94780-94792.	1.8	5
20	Novel multiple tyrosine kinase inhibitor ponatinib inhibits bFGF-activated signaling in neuroblastoma cells and suppresses neuroblastoma growth in vivo. <i>Oncotarget</i> , 2017, 8, 5874-5884.	1.8	17
21	Novel Src/Abl tyrosine kinase inhibitor bosutinib suppresses neuroblastoma growth via inhibiting Src/Abl signaling. <i>Oncotarget</i> , 2017, 8, 1469-1480.	1.8	25
22	Novel multi-targeted ErbB family inhibitor afatinib blocks EGF-induced signaling and induces apoptosis in neuroblastoma. <i>Oncotarget</i> , 2017, 8, 1555-1568.	1.8	22
23	TAK1 inhibitor 5Z-7-oxozeaenol sensitizes cervical cancer to doxorubicin-induced apoptosis. <i>Oncotarget</i> , 2017, 8, 33666-33675.	1.8	18
24	Small molecule inhibitor regorafenib inhibits RET signaling in neuroblastoma cells and effectively suppresses tumor growth <i>in vivo</i> . <i>Oncotarget</i> , 2017, 8, 104090-104103.	1.8	17
25	Novel proteasome inhibitor delanzomib sensitizes cervical cancer cells to doxorubicin-induced apoptosis via stabilizing tumor suppressor proteins in the p53 pathway. <i>Oncotarget</i> , 2017, 8, 114123-114135.	1.8	17
26	Microfluidic Cell Deformability Assay for Rapid and Efficient Kinase Screening with the CRISPR-Cas9 System. <i>Angewandte Chemie</i> , 2016, 128, 8703-8707.	2.0	6
27	Microfluidic Cell Deformability Assay for Rapid and Efficient Kinase Screening with the CRISPR-Cas9 System. <i>Angewandte Chemie - International Edition</i> , 2016, 55, 8561-8565.	13.8	26
28	Novel ALK inhibitor AZD3463 inhibits neuroblastoma growth by overcoming crizotinib resistance and inducing apoptosis. <i>Scientific Reports</i> , 2016, 6, 19423.	3.3	42
29	Wip1 inhibitor GSK2830371 inhibits neuroblastoma growth by inducing Chk2/p53-mediated apoptosis. <i>Scientific Reports</i> , 2016, 6, 38011.	3.3	29
30	Development of potent small-molecule inhibitors to drug the undruggable steroid receptor coactivator-3. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 4970-4975.	7.1	74
31	Multiple CDK inhibitor dinaciclib suppresses neuroblastoma growth via inhibiting CDK2 and CDK9 activity. <i>Scientific Reports</i> , 2016, 6, 29090.	3.3	60
32	Novel proteasome inhibitor ixazomib sensitizes neuroblastoma cells to doxorubicin treatment. <i>Scientific Reports</i> , 2016, 6, 34397.	3.3	20
33	Enrichment of Cdk1-cyclins at DNA double-strand breaks stimulates Fun30 phosphorylation and DNA end resection. <i>Nucleic Acids Research</i> , 2016, 44, 2742-2753.	14.5	39
34	Second-generation proteasome inhibitor carfilzomib sensitizes neuroblastoma cells to doxorubicin-induced apoptosis. <i>Oncotarget</i> , 2016, 7, 75914-75925.	1.8	17
35	Novel MDM2 inhibitor SAR405838 (MI-773) induces p53-mediated apoptosis in neuroblastoma. <i>Oncotarget</i> , 2016, 7, 82757-82769.	1.8	29
36	Stromal <i>CYR</i> 61 Confers Resistance to Mitoxantrone via Spleen Tyrosine Kinase Activation in Human Acute Myeloid Leukaemia. <i>British Journal of Haematology</i> , 2015, 170, 704-718.	2.5	27

#	ARTICLE	IF	CITATIONS
37	Tamoxifen Inhibits ER-negative Breast Cancer Cell Invasion and Metastasis by Accelerating Twist1 Degradation. <i>International Journal of Biological Sciences</i> , 2015, 11, 618-628.	6.4	29
38	Characterization of a Steroid Receptor Coactivator Small Molecule Stimulator that Overstimulates Cancer Cells and Leads to Cell Stress and Death. <i>Cancer Cell</i> , 2015, 28, 240-252.	16.8	69
39	Mus81 and converging forks limit the mutagenicity of replication fork breakage. <i>Science</i> , 2015, 349, 742-747.	12.6	162
40	Fission Yeast Pxd1 Promotes Proper DNA Repair by Activating Rad16XPF and Inhibiting Dna2. <i>PLoS Biology</i> , 2014, 12, e1001946.	5.6	19
41	Bufalin Is a Potent Small-Molecule Inhibitor of the Steroid Receptor Coactivators SRC-3 and SRC-1. <i>Cancer Research</i> , 2014, 74, 1506-1517.	0.9	145
42	Genome-wide Screens for Sensitivity to Ionizing Radiation Identify the Fission Yeast Nonhomologous End Joining Factor Xrc4. <i>G3: Genes, Genomes, Genetics</i> , 2014, 4, 1297-1306.	1.8	11
43	Identification of Verrucarin A as a Potent and Selective Steroid Receptor Coactivator-3 Small Molecule Inhibitor. <i>PLoS ONE</i> , 2014, 9, e95243.	2.5	33
44	A proteome-wide visual screen identifies fission yeast proteins localizing to DNA double-strand breaks. <i>DNA Repair</i> , 2013, 12, 433-443.	2.8	31
45	Phosphorylation of Thr-178 and Thr-184 in the TAK1 T-loop Is Required for Interleukin (IL)-1-mediated Optimal NF $\kappa$ B and AP-1 Activation as Well as IL-6 Gene Expression. <i>Journal of Biological Chemistry</i> , 2008, 283, 24497-24505.	3.4	94