
Christophe Hirtz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10645474/publications.pdf Version: 2024-02-01



CHDISTODUE HIDTZ

#	Article	IF	CITATIONS
1	Quantifying RNA modifications by mass spectrometry: a novel source of biomarkers in oncology. Critical Reviews in Clinical Laboratory Sciences, 2022, 59, 1-18.	6.1	14
2	Blood amyloid and tau biomarkers as predictors of cerebrospinal fluid profiles. Journal of Neural Transmission, 2022, 129, 231-237.	2.8	7
3	Changes Occurring on the Activity of Salivary Alpha-Amylase Proteoforms in Two Naturalistic Situations Using a Spectrophotometric Assay. Biology, 2021, 10, 227.	2.8	5
4	Tau protein in cerebrospinal fluid: a novel biomarker of the time of death?. International Journal of Legal Medicine, 2021, 135, 2081-2089.	2.2	7
5	Cytokine response following perturbation of the cervicovaginal milieu during HPV genital infection. Immunologic Research, 2021, 69, 255-263.	2.9	5
6	Use of plasma biomarkers for AT(N) classification of neurodegenerative dementias. Journal of Neurology, Neurosurgery and Psychiatry, 2021, 92, 1206-1214.	1.9	30
7	Serum glial fibrillary acidic protein is a predictor of brain metastases in patients with metastatic breast cancer. International Journal of Cancer, 2021, 149, 1605-1618.	5.1	8
8	Analytical comparison of ELISA and mass spectrometry for quantification of serum hepcidin in critically ill patients. Bioanalysis, 2021, 13, 1029-1035.	1.5	6
9	Concussion history in rugby union players is associated with depressed cerebrovascular reactivity and cognition. Scandinavian Journal of Medicine and Science in Sports, 2021, 31, 2291-2299.	2.9	7
10	Variation of human salivary alpha-amylase proteoforms in three stimulation models. Clinical Oral Investigations, 2020, 24, 475-486.	3.0	7
11	Detection of amyloid beta peptides in body fluids for the diagnosis of alzheimer's disease: Where do we stand?. Critical Reviews in Clinical Laboratory Sciences, 2020, 57, 99-113.	6.1	24
12	Cerebrospinal fluid A beta 1–40 peptides increase in Alzheimer's disease and are highly correlated with phospho-tau in control individuals. Alzheimer's Research and Therapy, 2020, 12, 123.	6.2	33
13	Hepcidin and ferritin levels in restless legs syndrome: a case–control study. Scientific Reports, 2020, 10, 11914.	3.3	21
14	Cerebrospinal Fluid and Plasma Biomarkers do not Differ in the Presenile and Late-Onset Behavioral Variants of Frontotemporal Dementia. Journal of Alzheimer's Disease, 2020, 74, 903-911.	2.6	9
15	Gravitational Transitions Increase Posterior Cerebral Perfusion and Systemic Oxidative-nitrosative Stress: Implications for Neurovascular Unit Integrity. Neuroscience, 2020, 441, 142-160.	2.3	9
16	Cerebrospinal fluid phospho-tau T217 outperforms T181 as a biomarker for the differential diagnosis of Alzheimer's disease and PET amyloid-positive patient identification. Alzheimer's Research and Therapy, 2020, 12, 26.	6.2	138
17	Soluble Intercellular Adhesion Molecule- (sICAM-) 1, Thrombospondin-1, and Vinculin for the Identification of Septic Shock Patients Suffering from an Invasive Fungal Infection. Mediators of Inflammation, 2020, 2020, 1-13.	3.0	2
18	Efficient extraction of intact HSA-Aβ peptide complexes from sera: Toward albuminome biomarker identification. Talanta, 2020, 216, 121002.	5.5	4

CHRISTOPHE HIRTZ

#	Article	IF	CITATIONS
19	Peripheral Blood and Salivary Biomarkers of Blood–Brain Barrier Permeability and Neuronal Damage: Clinical and Applied Concepts. Frontiers in Neurology, 2020, 11, 577312.	2.4	36
20	In Vivo Large-Scale Mapping of Protein Turnover in Human Cerebrospinal Fluid. Analytical Chemistry, 2019, 91, 15500-15508.	6.5	6
21	SILK studies — capturing the turnover of proteins linked to neurodegenerative diseases. Nature Reviews Neurology, 2019, 15, 419-427.	10.1	37
22	Stable Isotope Labeling Kinetics in CNS Translational Medicine: Introduction to SILK Technology. Handbook of Behavioral Neuroscience, 2019, 29, 173-190.	0.7	0
23	Biochemical markers of time since death in cerebrospinal fluid: A first step towards"Forensomics― Critical Reviews in Clinical Laboratory Sciences, 2019, 56, 274-286.	6.1	8
24	Natural history, dynamics, and ecology of human papillomaviruses in genital infections of young women: protocol of the PAPCLEAR cohort study. BMJ Open, 2019, 9, e025129.	1.9	17
25	Intact Protein Analysis by LC-MS for Characterizing Biomarkers in Cerebrospinal Fluid. Methods in Molecular Biology, 2019, 1959, 163-172.	0.9	0
26	The prognostic value of theÂTau protein serum level in metastatic breast cancer patients and its correlation with brain metastases. BMC Cancer, 2019, 19, 110.	2.6	20
27	Nano-flow vs standard-flow: Which is the more suitable LC/MS method for quantifying hepcidin-25 in human serum in routine clinical settings?. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2018, 1086, 110-117.	2.3	6
28	Association between serum hepcidin level and restless legs syndrome. Movement Disorders, 2018, 33, 618-627.	3.9	25
29	Assessing a multiplex-targeted proteomics approach for the clinical diagnosis of periodontitis using saliva samples. Bioanalysis, 2018, 10, 35-45.	1.5	12
30	Identification of multiple proteoforms biomarkers on clinical samples by routine Top-Down approaches. Data in Brief, 2018, 18, 1013-1021.	1.0	12
31	Towards a routine application of Top-Down approaches for label-free discovery workflows. Journal of Proteomics, 2018, 175, 12-26.	2.4	17
32	Sample Pooling and Inflammation Linked to the False Selection of Biomarkers for Neurodegenerative Diseases in Top–Down Proteomics: A Pilot Study. Frontiers in Molecular Neuroscience, 2018, 11, 477.	2.9	20
33	Impact of biological matrix on inflammatory protein biomarker quantification based on targeted mass spectrometry. Bioanalysis, 2018, 10, 1383-1399.	1.5	5
34	Plasma and CSF biomarkers for the diagnosis of Alzheimer's disease in adults with Down syndrome: a cross-sectional study. Lancet Neurology, The, 2018, 17, 860-869.	10.2	140
35	What sample preparation should be chosen for targeted MS monoclonal antibody quantification in human serum?. Bioanalysis, 2018, 10, 723-735.	1.5	12
36	Regulatory context and validation of assays for clinical mass spectrometry proteomics (cMSP) methods. Critical Reviews in Clinical Laboratory Sciences, 2018, 55, 346-358.	6.1	9

CHRISTOPHE HIRTZ

#	Article	IF	CITATIONS
37	Alzheimer's Disease: Advances in Drug Development. Journal of Alzheimer's Disease, 2018, 65, 3-13.	2.6	45
38	Clinical mass spectrometry proteomics (cMSP) for medical laboratory: What does the future hold?. Clinica Chimica Acta, 2017, 467, 51-58.	1.1	29
39	Characterizing Deep White Matter Hyperintensities in Patients with Symptomatic Isolated Cortical Superficial Siderosis. Journal of Stroke and Cerebrovascular Diseases, 2017, 26, 465-469.	1.6	1
40	Quantification of hepcidin-25 in human cerebrospinal fluid using LC–MS/MS. Bioanalysis, 2017, 9, 337-347.	1.5	12
41	Cerebrospinal fluid levels of orexin-A and histamine, and sleep profile within the Alzheimer process. Neurobiology of Aging, 2017, 53, 59-66.	3.1	76
42	Impurity determination for hepcidin by liquid chromatography-high resolution and ion mobility mass spectrometry for the value assignment of candidate primary calibrators. Analytical and Bioanalytical Chemistry, 2017, 409, 2559-2567.	3.7	16
43	Clinical perspectives of dried blood spot protein quantification using mass spectrometry methods. Critical Reviews in Clinical Laboratory Sciences, 2017, 54, 173-184.	6.1	19
44	Comparison of HbA1c detection in whole blood and dried blood spots using an automated ion-exchange HPLC system. Bioanalysis, 2017, 9, 427-434.	1.5	14
45	Impact of iron deficiency diagnosis using hepcidin mass spectrometry dosage methods on hospital stay and costs after a prolonged ICU stay: Study protocol for a multicentre, randomised, single-blinded medico-economic trial. Anaesthesia, Critical Care & Pain Medicine, 2017, 36, 391-396.	1.4	9
46	Cerebrospinal Fluid Alzheimer's Disease Biomarkers in Isolated Supratentorial Cortical Superficial Siderosis. Journal of Alzheimer's Disease, 2016, 54, 1291-1295.	2.6	16
47	From radioimmunoassay to mass spectrometry: a new method to quantify orexin-A (hypocretin-1) in cerebrospinal fluid. Scientific Reports, 2016, 6, 25162.	3.3	36
48	Differential Mass Spectrometry Profiles of Tau Protein in the Cerebrospinal Fluid of Patients with Alzheimer's Disease, Progressive Supranuclear Palsy, and Dementia with Lewy Bodies. Journal of Alzheimer's Disease, 2016, 51, 1033-1043.	2.6	104
49	Cerebrospinal Fluid Alzheimer's Disease Biomarkers in Cerebral Amyloid Angiopathy-Related Inflammation. Journal of Alzheimer's Disease, 2016, 50, 759-764.	2.6	23
50	Development of new quantitative mass spectrometry and semi-automatic isofocusing methods for the determination of Apolipoprotein E typing. Clinica Chimica Acta, 2016, 454, 33-38.	1.1	19
51	Absolute quantification of 35 plasma biomarkers in human saliva using targeted MS. Bioanalysis, 2016, 8, 43-53.	1.5	22
52	Tau Protein Quantification in Human Cerebrospinal Fluid by Targeted Mass Spectrometry at High Sequence Coverage Provides Insights into Its Primary Structure Heterogeneity. Journal of Proteome Research, 2016, 15, 667-676.	3.7	91
53	Stable Isotope Labeling by Amino acid <i>in Vivo</i> (SILAV): a new method to explore protein metabolism. Rapid Communications in Mass Spectrometry, 2015, 29, 1917-1925.	1.5	10
54	Antibody-free quantification of seven tau peptides in human CSF using targeted mass spectrometry. Frontiers in Neuroscience, 2015, 9, 302.	2.8	34

CHRISTOPHE HIRTZ

#	ARTICLE	IF	CITATIONS
55	What is the potential of dried matrix spot sampling for cerebrospinal fluid analysis?. Bioanalysis, 2015, 7, 2849-2851.	1.5	1
56	Quantitative detection of amyloid-l ² peptides by mass spectrometry: state of the art and clinical applications. Clinical Chemistry and Laboratory Medicine, 2015, 53, 1483-93.	2.3	30
57	Impact of harmonization of collection tubes on Alzheimer's disease diagnosis. , 2014, 10, S390-S394.e2.		58
58	Development and validation of dried matrix spot sampling for the quantitative determination of amyloid β peptides in cerebrospinal fluid. Clinical Chemistry and Laboratory Medicine, 2014, 52, 649-55.	2.3	8
59	Clinical measurement of Hepcidin-25 in human serum: Is quantitative mass spectrometry up to the job?. EuPA Open Proteomics, 2014, 3, 60-67.	2.5	19
60	Current and future use of "dried blood spot―analyses in clinical chemistry. Clinical Chemistry and Laboratory Medicine, 2013, 51, 1897-1909.	2.3	102
61	Quantitative Clinical Chemistry Proteomics (qCCP) using mass spectrometry: general characteristics and application. Clinical Chemistry and Laboratory Medicine, 2013, 51, 919-35.	2.3	47
62	From "Clinical Proteomics―to "Clinical Chemistry Proteomics― considerations using quantitative mass-spectrometry as a model approach. Clinical Chemistry and Laboratory Medicine, 2012, 50, 235-42.	2.3	7
63	Proteins and proteolysis in pre-term and term human milk and possible implications for infant formulae. International Dairy Journal, 2010, 20, 715-723.	3.0	56
64	Use of Reducing/Nonreducing Two-Dimensional Electrophoresis for the Study of Disulfide-Mediated Interactions between Proteins in Raw and Heated Bovine Milk. Journal of Agricultural and Food Chemistry, 2009, 57, 5948-5955.	5.2	54
65	Proteomic Studies of Saliva: A Proposal for a Standardized Handling of Clinical Samples. Clinical Proteomics, 2007, 3, 13-21.	2.1	23
66	Salivary protein profiling in type I diabetes using two-dimensional electrophoresis and mass spectrometry. Clinical Proteomics, 2006, 2, 117-127.	2.1	18
67	MS characterization of multiple forms of alpha-amylase in human saliva. Proteomics, 2005, 5, 4597-4607.	2.2	70