

# Elizabeth A Thomas

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10667781/publications.pdf>

Version: 2024-02-01

70  
papers

4,054  
citations

109321

35  
h-index

118850

62  
g-index

70  
all docs

70  
docs citations

70  
times ranked

5842  
citing authors

#	ARTICLE	IF	CITATIONS
1	Salivary S100 calcium-binding protein beta (S100B) and neurofilament light (NfL) after acute exposure to repeated head impacts in collegiate water polo players. <i>Scientific Reports</i> , 2022, 12, 3439.	3.3	5
2	Associations between prognostic index scores and plasma neurofilament light in Huntington's disease. <i>Parkinsonism and Related Disorders</i> , 2022, 97, 25-28.	2.2	6
3	Plasma neurofilament light in Huntington's disease: A marker for disease onset, but not symptom progression. <i>Parkinsonism and Related Disorders</i> , 2021, 87, 32-38.	2.2	19
4	Saliva testing as a means to monitor therapeutic lithium levels in patients with psychiatric disorders: Identification of clinical and environmental covariates, and their incorporation into a prediction model. <i>Bipolar Disorders</i> , 2021, 23, 679-688.	1.9	14
5	Levels of Interleukin-6 in Saliva, but Not Plasma, Correlate with Clinical Metrics in Huntington's Disease Patients and Healthy Control Subjects. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6363.	4.1	27
6	Reducing Mcl-1 gene dosage induces dopaminergic neuronal loss and motor impairments in Park2 knockout mice. <i>Communications Biology</i> , 2019, 2, 125.	4.4	11
7	Epigenetic mechanisms in Huntington's disease. , 2019, , 73-95.		3
8	Evaluation of Biochemical and Epigenetic Measures of Peripheral Brain-Derived Neurotrophic Factor (BDNF) as a Biomarker in Huntington's Disease Patients. <i>Frontiers in Molecular Neuroscience</i> , 2019, 12, 335.	2.9	41
9	Complex neuroprotective and neurotoxic effects of histone deacetylases. <i>Journal of Neurochemistry</i> , 2018, 145, 96-110.	3.9	55
10	Salivary levels of total huntingtin are elevated in Huntington's disease patients. <i>Scientific Reports</i> , 2018, 8, 7371.	3.3	25
11	Histone Posttranslational Modifications in Schizophrenia. <i>Advances in Experimental Medicine and Biology</i> , 2017, 978, 237-254.	1.6	26
12	Beneficial effects of glatiramer acetate in Huntington's disease mouse models: Evidence for BDNF-elevating and immunomodulatory mechanisms. <i>Brain Research</i> , 2017, 1673, 102-110.	2.2	16
13	The Role of Histone Deacetylase Inhibition in the Accumulation and Stability of Disease-Related Proteins. , 2017, , 159-179.		3
14	The Effects of Pharmacological Inhibition of Histone Deacetylase 3 (HDAC3) in Huntington's Disease Mice. <i>PLoS ONE</i> , 2016, 11, e0152498.	2.5	73
15	Increased cortical expression of the zinc transporter SLC39A12 suggests a breakdown in zinc cellular homeostasis as part of the pathophysiology of schizophrenia. <i>NPJ Schizophrenia</i> , 2016, 2, 16002.	3.6	47
16	DNA methylation in Huntington's disease: Implications for transgenerational effects. <i>Neuroscience Letters</i> , 2016, 625, 34-39.	2.1	33
17	HDAC inhibition imparts beneficial transgenerational effects in Huntington's disease mice via altered DNA and histone methylation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, E56-64.	7.1	95
18	Disease Modifying Potential of Glatiramer Acetate in Huntington's Disease. <i>Journal of Huntington's Disease</i> , 2014, 3, 311-316.	1.9	12

#	ARTICLE	IF	CITATIONS
19	Involvement of HDAC1 and HDAC3 in the Pathology of Polyglutamine Disorders: Therapeutic Implications for Selective HDAC1/HDAC3 Inhibitors. <i>Pharmaceuticals</i> , 2014, 7, 634-661.	3.8	34
20	Behavioral and transcriptome alterations in male and female mice with postnatal deletion of TrkB in dorsal striatal medium spiny neurons. <i>Molecular Neurodegeneration</i> , 2013, 8, 47.	10.8	11
21	Epigenetic changes at gene promoters in response to immune activation in utero. <i>Brain, Behavior, and Immunity</i> , 2013, 30, 168-175.	4.1	78
22	Selective histone deacetylase (HDAC) inhibition imparts beneficial effects in Huntington's disease mice: implications for the ubiquitin-proteasomal and autophagy systems. <i>Human Molecular Genetics</i> , 2012, 21, 5280-5293.	2.9	128
23	Egr-1 Induces DARPP-32 Expression in Striatal Medium Spiny Neurons via a Conserved Intragenic Element. <i>Journal of Neuroscience</i> , 2012, 32, 6808-6818.	3.6	21
24	Forkhead box protein p1 is a transcriptional repressor of immune signaling in the CNS: implications for transcriptional dysregulation in Huntington disease. <i>Human Molecular Genetics</i> , 2012, 21, 3097-3111.	2.9	55
25	Differential age- and disease-related effects on the expression of genes related to the arachidonic acid signaling pathway in schizophrenia. <i>Psychiatry Research</i> , 2012, 196, 201-206.	3.3	34
26	Histone deacetylase (HDAC) inhibitors targeting HDAC3 and HDAC1 ameliorate polyglutamine-elicited phenotypes in model systems of Huntington's disease. <i>Neurobiology of Disease</i> , 2012, 46, 351-361.	4.4	157
27	Sphingolipid abnormalities in psychiatric disorders: a missing link in pathology?. <i>Frontiers in Bioscience - Landmark</i> , 2011, 16, 1797.	3.0	24
28	Gene expression profiling of R6/2 transgenic mice with different CAG repeat lengths reveals genes associated with disease onset and progression in Huntington's disease. <i>Neurobiology of Disease</i> , 2011, 42, 459-467.	4.4	63
29	Phospholipase C Beta 1 Expression in the Dorsolateral Prefrontal Cortex from Patients with Schizophrenia at Different Stages of Illness. <i>Australian and New Zealand Journal of Psychiatry</i> , 2011, 45, 140-147.	2.3	29
30	In vivo cell-autonomous transcriptional abnormalities revealed in mice expressing mutant huntingtin in striatal but not cortical neurons. <i>Human Molecular Genetics</i> , 2011, 20, 1049-1060.	2.9	56
31	Changes in Gene Expression in Subjects with Schizophrenia Associated with Disease Progression. , 2011, , 237-251.		1
32	Genome-Wide Identification of Bcl11b Gene Targets Reveals Role in Brain-Derived Neurotrophic Factor Signaling. <i>PLoS ONE</i> , 2011, 6, e23691.	2.5	53
33	Cerebellar lipid differences between R6/1 transgenic mice and humans with Huntington's disease. <i>Journal of Neurochemistry</i> , 2010, 115, 748-758.	3.9	36
34	Chronic monoacylglycerol lipase blockade causes functional antagonism of the endocannabinoid system. <i>Nature Neuroscience</i> , 2010, 13, 1113-1119.	14.8	534
35	Low density lipoprotein receptor-related protein and apolipoprotein E expression is altered in schizophrenia. <i>Frontiers in Psychiatry</i> , 2010, 1, 19.	2.6	12
36	Coexpression network analysis of neural tissue reveals perturbations in developmental processes in schizophrenia. <i>Genome Research</i> , 2010, 20, 403-412.	5.5	127

#	ARTICLE	IF	CITATIONS
37	A role of apolipoprotein D in triglyceride metabolism. <i>Journal of Lipid Research</i> , 2010, 51, 1298-1311.	4.2	59
38	Evidence for disruption of sphingolipid metabolism in schizophrenia. <i>Journal of Neuroscience Research</i> , 2009, 87, 278-288.	2.9	85
39	Focal Nature of Neurological Disorders Necessitates Isotype-Selective Histone Deacetylase (HDAC) Inhibitors. <i>Molecular Neurobiology</i> , 2009, 40, 33-45.	4.0	50
40	Normal human aging and early-stage schizophrenia share common molecular profiles. <i>Aging Cell</i> , 2009, 8, 339-342.	6.7	41
41	Apolipoprotein D mRNA expression is elevated in PDAPP transgenic mice. <i>Journal of Neurochemistry</i> , 2008, 79, 1059-1064.	3.9	25
42	Molecular profiles of schizophrenia in the CNS at different stages of illness. <i>Brain Research</i> , 2008, 1239, 235-248.	2.2	178
43	Functional roles for the striatal-enriched transcription factor, Bcl11b, in the control of striatal gene expression and transcriptional dysregulation in Huntington's disease. <i>Neurobiology of Disease</i> , 2008, 31, 298-308.	4.4	63
44	The HDAC inhibitor 4b ameliorates the disease phenotype and transcriptional abnormalities in Huntington's disease transgenic mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 15564-15569.	7.1	271
45	Regulator of G-Protein Signalling 4 Expression is Not Altered in the Prefrontal Cortex in Schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 2008, 42, 740-745.	2.3	12
46	Gene Expression Profiling in Brodmann's Area 46 from Subjects with Schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 2007, 41, 308-320.	2.3	74
47	Clozapine specifically alters the arachidonic acid pathway in mice lacking apolipoprotein D. <i>Schizophrenia Research</i> , 2007, 89, 147-153.	2.0	23
48	Chronic haloperidol treatment results in a decrease in the expression of myelin/oligodendrocyte-related genes in the mouse brain. <i>Journal of Neuroscience Research</i> , 2007, 85, 757-765.	2.9	52
49	Glycolipid and ganglioside metabolism imbalances in Huntington's disease. <i>Neurobiology of Disease</i> , 2007, 27, 265-277.	4.4	120
50	Selective deficits in the expression of striatal-enriched mRNAs in Huntington's disease. <i>Journal of Neurochemistry</i> , 2006, 96, 743-757.	3.9	125
51	Molecular Profiling of Antipsychotic Drug Function: Convergent Mechanisms in the Pathology and Treatment of Psychiatric Disorders. <i>Molecular Neurobiology</i> , 2006, 34, 109-128.	4.0	35
52	Striatal specificity of gene expression dysregulation in Huntington's disease. <i>Journal of Neuroscience Research</i> , 2006, 84, 1151-1164.	2.9	53
53	Association of plasma apolipoproteins D with RBC membrane arachidonic acid levels in schizophrenia. <i>Schizophrenia Research</i> , 2005, 72, 259-266.	2.0	19
54	Apolipoprotein D levels are elevated in prefrontal cortex of subjects with Alzheimer's disease. <i>Biological Psychiatry</i> , 2003, 54, 136-141.	1.3	51

#	ARTICLE	IF	CITATIONS
55	Apolipoprotein D modulates arachidonic acid signaling in cultured cells: implications for psychiatric disorders. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2003, 69, 421-427.	2.2	34
56	Increased levels of apolipoprotein E in the frontal cortex of subjects with schizophrenia. <i>Biological Psychiatry</i> , 2003, 54, 616-622.	1.3	41
57	Novel Isoform of Insulin Receptor Substrate p53/p58 Is Generated by Alternative Splicing in the CRIB/SH3-binding Region. <i>Journal of Biological Chemistry</i> , 2002, 277, 24728-24734.	3.4	21
58	Evolutionarily Distinct Classes of S27 Ribosomal Proteins with Differential mRNA Expression in Rat Hypothalamus. <i>Journal of Neurochemistry</i> , 2002, 74, 2259-2267.	3.9	26
59	The Endogenous Lipid Oleamide Activates Serotonin 5-HT <sub>7</sub> Neurons in Mouse Thalamus and Hypothalamus. <i>Journal of Neurochemistry</i> , 2002, 72, 2370-2378.	3.9	40
60	The Neurobiology of Apolipoproteins in Psychiatric Disorders. <i>Molecular Neurobiology</i> , 2002, 26, 369-388.	4.0	32
61	Insulin receptor substrate protein p53 localization in rats suggests mechanism for specific polyglutamine neurodegeneration. <i>Neuroscience Letters</i> , 2001, 309, 145-148.	2.1	17
62	Clozapine increases apolipoprotein D expression in rodent brain: towards a mechanism for neuroleptic pharmacotherapy. <i>Journal of Neurochemistry</i> , 2001, 76, 789-796.	3.9	60
63	Pertussis toxin treatment prevents 5-HT <sub>5a</sub> receptor-mediated inhibition of cyclic AMP accumulation in rat C6 glioma cells. <i>Journal of Neuroscience Research</i> , 2000, 61, 75-81.	2.9	24
64	Allosteric regulation by oleamide of the binding properties of 5-hydroxytryptamine <sub>7</sub> receptors. <i>Biochemical Pharmacology</i> , 1999, 58, 1807-1813.	4.4	54
65	RGS9: A regulator of G-protein signalling with specific expression in rat and mouse striatum. <i>Journal of Neuroscience Research</i> , 1998, 52, 118-124.	2.9	76
66	Fatty acid amide hydrolase, the degradative enzyme for anandamide and oleamide, has selective distribution in neurons within the rat central nervous system. <i>Journal of Neuroscience Research</i> , 1997, 50, 1047-1052.	2.9	162
67	Fatty acid amide hydrolase, the degradative enzyme for anandamide and oleamide, has selective distribution in neurons within the rat central nervous system. <i>Journal of Neuroscience Research</i> , 1997, 50, 1047-1052.	2.9	3
68	Involvement of the M2 muscarinic receptor in contractions of the guinea pig trachea, guinea pig esophagus, and rat fundus. <i>Biochemical Pharmacology</i> , 1996, 51, 779-788.	4.4	40
69	The 5-HT <sub>5A</sub> serotonin receptor is expressed predominantly by astrocytes in which it inhibits cAMP accumulation: A mechanism for neuronal suppression of reactive astrocytes. , 1996, 17, 317-326.		87
70	Functional role of M2 muscarinic receptors in the guinea pig ileum. <i>Life Sciences</i> , 1995, 56, 965-971.	4.3	37