
Richard A Lutz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10701848/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Seasonal Changes in Shell Microstructure of Some Common Bivalve Molluscs in the Mid-Atlantic Region. Journal of Shellfish Research, 2022, 41, .	0.9	2
2	Shell Morphology and Morphometry of Larval and Post-Larval Donax fossor Say (Bivalvia: Donacidae). Journal of Shellfish Research, 2021, 40, .	0.9	0
3	Hydrothermal Vent Biota. , 2019, , 308-319.		3
4	Scanning Electron Microscopic Aids for Identification of Larval and Post-Larval Bivalves. Journal of Shellfish Research, 2018, 37, 247-448.	0.9	4
5	Optical Imaging and Molecular Sequencing of a Preserved Collection of Bivalve Larvae. Journal of Shellfish Research, 2018, 37, 449-466.	0.9	3
6	Microbial biofilms associated with fluid chemistry and megafaunal colonization at post-eruptive deep-sea hydrothermal vents. Deep-Sea Research Part II: Topical Studies in Oceanography, 2015, 121, 31-40.	1.4	25
7	Deep-sea hydrothermal vent <i>Epsilonproteobacteria</i> encode a conserved and widespread nitrate reduction pathway (Nap). ISME Journal, 2014, 8, 1510-1521.	9.8	86
8	A Dive to Challenger Deep. Science, 2012, 336, 301-302.	12.6	25
9	Phorcysia thermohydrogeniphila gen. nov., sp. nov., a thermophilic, chemolithoautotrophic, nitrate-ammonifying bacterium from a deep-sea hydrothermal vent. International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 2388-2394.	1.7	20
10	Genetic diversity and demographic instability in Riftia pachyptilatubeworms from eastern Pacific hydrothermal vents. BMC Evolutionary Biology, 2011, 11, 96.	3.2	34
11	Salinisphaera hydrothermalis sp. nov., a mesophilic, halotolerant, facultatively autotrophic, thiosulfate-oxidizing gammaproteobacterium from deep-sea hydrothermal vents, and emended description of the genus Salinisphaera. International Journal of Systematic and Evolutionary Microbiology, 2009, 59, 1497-1503.	1.7	38
12	Pre- and post-eruption diffuse flow variability among tubeworm habitats at 9°50â€2 north on the East Pacific Rise. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 1607-1615.	1.4	19
13	Phylogenetic diversity of methanogenic, sulfate-reducing and methanotrophic prokaryotes from deep-sea hydrothermal vents and cold seeps. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 1665-1674.	1.4	27
14	Paleodictyon nodosum: A living fossil on the deep-sea floor. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 1700-1712.	1.4	56
15	Hydrothermal Vent Mussel Habitat Chemistry, Pre- and Post-Eruption at 9°50â€2North on the East Pacific Rise. Journal of Shellfish Research, 2008, 27, 169-175.	0.9	29
16	Interrelationships Between Vent Fluid Chemistry, Temperature, Seismic Activity, and Biological Community Structure at a Mussel-Dominated, Deep-Sea Hydrothermal Vent Along the East Pacific Rise. Journal of Shellfish Research, 2008, 27, 177-190.	0.9	31
17	Vertical distribution and diversity of bacteria and archaea in sulfide and methane-rich cold seep sediments located at the base of the Florida Escarpment. Extremophiles, 2006, 10, 199-211.	2.3	59
18	Mercury Adaptation among Bacteria from a Deep-Sea Hydrothermal Vent. Applied and Environmental Microbiology, 2005, 71, 220-226.	3.1	109

RICHARD A LUTZ

#	Article	IF	CITATIONS
19	Thermovibrio ammonificans sp. nov., a thermophilic, chemolithotrophic, nitrate-ammonifying bacterium from deep-sea hydrothermal vents. International Journal of Systematic and Evolutionary Microbiology, 2004, 54, 175-181.	1.7	97
20	Experimental ecology at deep-sea hydrothermal vents: a perspective. Journal of Experimental Marine Biology and Ecology, 2004, 300, 273-307.	1.5	64
21	Chemical speciation drives hydrothermal vent ecology. Nature, 2001, 410, 813-816.	27.8	337
22	Neutral and Nonneutral Mitochondrial Genetic Variation in Deep-Sea Clams from the Family Vesicomyidae. Journal of Molecular Evolution, 2000, 50, 141-153.	1.8	43
23	Miocene Radiation of Deep-Sea Hydrothermal Vent Shrimp (Caridea: Bresiliidae): Evidence from Mitochondrial Cytochrome Oxidase Subunit I. Molecular Phylogenetics and Evolution, 1999, 13, 244-254.	2.7	113
24	Calcium carbonate dissolution rates in deep-sea bivalve shells on the East Pacific Rise at 21°N: results of an 8-year in-situ experiment. Palaeogeography, Palaeoclimatology, Palaeoecology, 1999, 154, 293-299.	2.3	26
25	Population genetics and biogeography of vestimentiferan tube worms. Deep-Sea Research Part II: Topical Studies in Oceanography, 1998, 45, 365-382.	1.4	32
26	Temporal and spatial patterns of biological community development at nascent deep-sea hydrothermal vents (9°50′N, East Pacific Rise). Deep-Sea Research Part II: Topical Studies in Oceanography, 1998, 45, 465-515.	1.4	366
27	Patterns of dispersal and larval development of archaeogastropod limpets at hydrothermal vents in the eastern Pacific. Journal of Experimental Marine Biology and Ecology, 1997, 210, 37-51.	1.5	37
28	Calcium carbonate dissolution rates in hydrothermal vent fields of the Guaymas Basin. Journal of Marine Research, 1994, 52, 969-982.	0.3	16
29	Cryptic species of deep-sea clams (Mollusca: Bivalvia: Vesicomyidae) from hydrothermal vent and cold-water seep environments. Deep-Sea Research Part I: Oceanographic Research Papers, 1994, 41, 1171-1189.	1.4	67
30	Rapid growth at deep-sea vents. Nature, 1994, 371, 663-664.	27.8	203
31	Ecology of deep-sea hydrothermal vent communities: A review. Reviews of Geophysics, 1993, 31, 211.	23.0	163
32	Electrophoretic identification and genetic analysis of bivalve larvae. Marine Biology, 1992, 113, 227-230.	1.5	33
33	Biomineralization of barite in the shell of the freshwater Asiatic clam Corbicula fluminea (Molluscs:) Tj ETQq1 1	0.784314 ı 3.1	rgBŢ ₃ /Overloo
34	The relationship of larval shell morphology to mode of development in marine prosobranch gastropods. Journal of the Marine Biological Association of the United Kingdom, 1990, 70, 611-637.	0.8	42
35	Shell morphology of larval and post-larval mytilids from the north-western Atlantic. Journal of the Marine Biological Association of the United Kingdom, 1989, 69, 181-218.	0.8	37
36	Procedures for Accurate Documentation of Shapes and Dimensions of Larval Bivalve Shells with Scanning Electron Microscopy. Transactions of the American Microscopical Society, 1989, 108, 58.	0.3	12

#	Article	IF	CITATIONS
37	A comparison of bivalve (Calyptogena magnifica) growth at two deep-sea hydrothermal vents in the eastern Pacific. Deep-sea Research Part A, Oceanographic Research Papers, 1988, 35, 1793-1810.	1.5	32

Temporal change in megafauna at the Rose Garden hydrothermal vent (Galapagos Rift; eastern tropical) Tj ETQq0 0.0 rgBT /Overlock 10

39	Mussel Aquaculture in the United States. , 1985, , 311-363.		4
40	Seasonal and geographic variation in the shell microstructure of a salt-marsh bivalve (<l>Geukensia demissa</l> (Dillwyn)). Journal of Marine Research, 1984, 42, 943-956.	0.3	16
41	LARVAL ECOLOGY OF MARINE BENTHIC INVERTEBRATES: PALEOBIOLOGICAL IMPLICATIONS. Biological Reviews, 1983, 58, 21-89.	10.4	507
42	Molluscan Larval Shell Morphology. Topics in Geobiology, 1980, , 323-377.	0.5	112
43	Hinge morphogenesis in the shells of larval and early post-larval mussels (<i>Mytilus edulis</i> L. and) Tj ETQq1 1 1979, 59, 111-121.	0.784314 0.8	4 rgBT /Ove 39
44	Larval ecology of extinct molluscs: Comment on larval development of hyolithids. Lethaia, 1979, 12, 306-306.	1.4	3
45	LARVAL DEVELOPMENT OF THE NORTHERN HORSE MUSSEL, MODIOLUS MODIOLUS (L.), INCLUDING A COMPARISON WITH THE LARVAE OF MYTILUS EDULIS L. AS AN AID IN PLANKTONIC IDENTIFICATION. Biological Bulletin, 1976, 150, 348-360.	1.8	46