Vassilis Golfinopoulos

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10740115/publications.pdf

Version: 2024-02-01

567281 794594 4,137 19 15 19 citations h-index g-index papers 20 20 20 6349 docs citations times ranked citing authors all docs

| # | Article | IF | CITATIONS |
|----|--|------|---------------------|
| 1 | Trabectedin for recurrent WHO grade 2 or 3 meningioma: A randomized phase II study of the EORTC Brain Tumor Group (EORTC-1320-BTG). Neuro-Oncology, 2022, 24, 755-767. | 1.2 | 25 |
| 2 | Non-IDH1-R132H IDH1/2 mutations are associated with increased DNA methylation and improved survival in astrocytomas, compared to IDH1-R132H mutations. Acta Neuropathologica, 2021, 141, 945-957. | 7.7 | 32 |
| 3 | Prognostic significance of genome-wide DNA methylation profiles within the randomized, phase 3, EORTC CATNON trial on non-1p/19q deleted anaplastic glioma. Neuro-Oncology, 2021, 23, 1547-1559. | 1.2 | 34 |
| 4 | Impact of depatuxizumab mafodotin on health-related quality of life and neurological functioning in the phase II EORTC 1410/INTELLANCE 2 trial for EGFR-amplified recurrent glioblastoma. European Journal of Cancer, 2021, 147, 1-12. | 2.8 | 10 |
| 5 | Adjuvant and concurrent temozolomide for $1p/19q$ non-co-deleted anaplastic glioma (CATNON; EORTC) Tj ETQq1 Oncology, The, 2021, 22, 813-823. | | 14 rgBT /Ove 132 |
| 6 | INTELLANCE 2/EORTC 1410 randomized phase II study of Depatux-M alone and with temozolomide vs temozolomide or lomustine in recurrent EGFR amplified glioblastoma. Neuro-Oncology, 2020, 22, 684-693. | 1.2 | 126 |
| 7 | Temporal muscle thickness is an independent prognostic marker in patients with progressive glioblastoma: translational imaging analysis of the EORTC 26101 trial. Neuro-Oncology, 2019, 21, 1587-1594. | 1.2 | 56 |
| 8 | Imaging necrosis during treatment is associated with worse survival in EORTC 26101 study. Neurology, 2019, 92, e2754-e2763. | 1.1 | 9 |
| 9 | Bevacizumab and temozolomide in patients with first recurrence of WHO grade II and III glioma, without 1p/19q co-deletion (TAVAREC): a randomised controlled phase 2 EORTC trial. Lancet Oncology, The, 2018, 19, 1170-1179. | 10.7 | 80 |
| 10 | Is more better? The impact of extended adjuvant temozolomide in newly diagnosed glioblastoma: a secondary analysis of EORTC and NRG Oncology/RTOG. Neuro-Oncology, 2017, 19, 1119-1126. | 1.2 | 107 |
| 11 | Short-Course Radiation plus Temozolomide in Elderly Patients with Glioblastoma. New England Journal of Medicine, 2017, 376, 1027-1037. | 27.0 | 810 |
| 12 | Interim results from the CATNON trial (EORTC study 26053-22054) of treatment with concurrent and adjuvant temozolomide for $1p/19q$ non-co-deleted anaplastic glioma: a phase 3, randomised, open-label intergroup study. Lancet, The, 2017, 390, 1645-1653. | 13.7 | 307 |
| 13 | Lomustine and Bevacizumab in Progressive Glioblastoma. New England Journal of Medicine, 2017, 377, 1954-1963. | 27.0 | 670 |
| 14 | Precision Medicine: From "Omics―to Economics towards Data-Driven Healthcare - Time for European Transformation. Biomedicine Hub, 2017, 2, 1-10. | 1.2 | 7 |
| 15 | Final results of the EORTC Brain Tumor Group randomized phase II TAVAREC trial on temozolomide with or without bevacizumab in 1st recurrence grade II/III glioma without 1p/19q co-deletion Journal of Clinical Oncology, 2017, 35, 2009-2009. | 1.6 | 8 |
| 16 | 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. New England Journal of Medicine, 2016, 375, 717-729. | 27.0 | 1,427 |
| 17 | Evaluating novel agent effects in multipleâ€treatments metaâ€regression. Statistics in Medicine, 2010, 29, 2369-2383. | 1.6 | 84 |
| 18 | Survival and disease-progression benefits with treatment regimens for advanced colorectal cancer: a meta-analysis. Lancet Oncology, The, 2007, 8, 898-911. | 10.7 | 149 |

| # | Article | IF | CITATIONS |
|-------|---|-----|-----------|
| 19 | Treatment of colorectal cancer in the elderly: A review of the literature. Cancer Treatment Reviews, 2006, 32, 1-8. | 7.7 | 64 |