

Jian-Xiang Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1077829/publications.pdf>

Version: 2024-02-01

51
papers

4,560
citations

186265

28
h-index

182427

51
g-index

51
all docs

51
docs citations

51
times ranked

4334
citing authors

#	ARTICLE	IF	CITATIONS
1	Gene delivery strategies for therapeutic proteins production in plants: Emerging opportunities and challenges. <i>Biotechnology Advances</i> , 2022, 54, 107845.	11.7	6
2	bZIP17 regulates heat stress tolerance at reproductive stage in <i>Arabidopsis</i> . <i>ABIOTECH</i> , 2022, 3, 1-11.	3.9	22
3	Regulation of Chloroplast Development and Function at Adverse Temperatures in Plants. <i>Plant and Cell Physiology</i> , 2022, , .	3.1	6
4	UBA domain protein SUF1 interacts with NatA complex subunit NAA15 to regulate thermotolerance in <i>Arabidopsis</i> . <i>Journal of Integrative Plant Biology</i> , 2022, 64, 1297-1302.	8.5	2
5	REVEILLE 7 inhibits the expression of the circadian clock gene <i>EARLY FLOWERING 4</i> to fine-tune hypocotyl growth in response to warm temperatures. <i>Journal of Integrative Plant Biology</i> , 2022, 64, 1310-1324.	8.5	9
6	Histone H3K4 methyltransferases SDG25 and ATX1 maintain heat stress gene expression during recovery in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2021, 105, 1326-1338.	5.7	41
7	Ectopic overexpression of a membrane-ethered transcription factor gene <i>NAC60</i> from oilseed rape positively modulates programmed cell death and age-triggered leaf senescence. <i>Plant Journal</i> , 2021, 105, 600-618.	5.7	21
8	Protein Quality Control in Plant Organelles: Current Progress and Future Perspectives. <i>Molecular Plant</i> , 2021, 14, 95-114.	8.3	77
9	Two B-box domain proteins, BBX28 and BBX29, regulate flowering time at low ambient temperature in <i>Arabidopsis</i> . <i>Plant Molecular Biology</i> , 2021, 106, 21-32.	3.9	17
10	Chromatin remodeling factors regulate environmental stress responses in plants. <i>Journal of Integrative Plant Biology</i> , 2021, 63, 438-450.	8.5	42
11	XBAT31 regulates thermoresponsive hypocotyl growth through mediating degradation of the thermosensor ELF3 in <i>Arabidopsis</i> . <i>Science Advances</i> , 2021, 7, .	10.3	42
12	The E3 ligase XBAT35 mediates thermoresponsive hypocotyl growth by targeting ELF3 for degradation in <i>Arabidopsis</i> . <i>Journal of Integrative Plant Biology</i> , 2021, 63, 1097-1103.	8.5	24
13	Timing to grow: roles of clock in thermomorphogenesis. <i>Trends in Plant Science</i> , 2021, 26, 1248-1257.	8.8	16
14	Phosphoproteomic Analysis of Thermomorphogenic Responses in <i>Arabidopsis</i> . <i>Frontiers in Plant Science</i> , 2021, 12, 753148.	3.6	3
15	The FtsH-Inactive Protein FtsHi5 Is Required for Chloroplast Development and Protein Accumulation in Chloroplasts at Low Ambient Temperature in <i>Arabidopsis</i> . <i>Frontiers in Plant Science</i> , 2021, 12, 830390.	3.6	5
16	Roles of plant hormones in thermomorphogenesis. <i>Stress Biology</i> , 2021, 1, .	3.1	2
17	A membrane-associated NAC transcription factor OsNTL3 is involved in thermotolerance in rice. <i>Plant Biotechnology Journal</i> , 2020, 18, 1317-1329.	8.3	126
18	Quantitative Proteomic Analysis of ER Stress Response Reveals both Common and Specific Features in Two Contrasting Ecotypes of <i>Arabidopsis thaliana</i> . <i>International Journal of Molecular Sciences</i> , 2020, 21, 9741.	4.1	8

#	ARTICLE	IF	CITATIONS
19	NAC103, a NAC family transcription factor, regulates ABA response during seed germination and seedling growth in Arabidopsis. <i>Planta</i> , 2020, 252, 95.	3.2	14
20	Mutation of DELAYED GREENING1 impairs chloroplast RNA editing at elevated ambient temperature in Arabidopsis. <i>Journal of Genetics and Genomics</i> , 2020, 47, 201-212.	3.9	14
21	BLISTER-regulated vegetative growth is dependent on the protein kinase domain of ER stress modulator IRE1A in Arabidopsis thaliana. <i>PLoS Genetics</i> , 2019, 15, e1008563.	3.5	15
22	The 25 subunit is essential for intact 26S proteasome assembly to specifically promote plant autotrophic growth under salt stress. <i>New Phytologist</i> , 2019, 221, 1359-1368.	7.3	32
23	A Novel QTL qTGW3 Encodes the GSK3/SHAGGY-Like Kinase OsGSK5/OsSK41 that Interacts with OsARF4 to Negatively Regulate Grain Size and Weight in Rice. <i>Molecular Plant</i> , 2018, 11, 736-749.	8.3	201
24	Two B-Box Domain Proteins, BBX18 and BBX23, Interact with ELF3 and Regulate Thermomorphogenesis in Arabidopsis. <i>Cell Reports</i> , 2018, 25, 1718-1728.e4.	6.4	91
25	Chromatin remodeling factor CHR18 interacts with replication protein RPA1A to regulate the DNA replication stress response in Arabidopsis. <i>New Phytologist</i> , 2018, 220, 476-487.	7.3	8
26	Tissue-Specific Transcriptomics Reveals an Important Role of the Unfolded Protein Response in Maintaining Fertility upon Heat Stress in Arabidopsis. <i>Plant Cell</i> , 2017, 29, 1007-1023.	6.6	130
27	Cellulose synthesis genes <i>CESA6</i> and <i>CSI1</i> are important for salt stress tolerance in Arabidopsis. <i>Journal of Integrative Plant Biology</i> , 2016, 58, 623-626.	8.5	45
28	Managing the protein folding demands in the endoplasmic reticulum of plants. <i>New Phytologist</i> , 2016, 211, 418-428.	7.3	165
29	Membrane-associated transcription factor peptidase, site-2 protease, antagonizes ABA signaling in Arabidopsis. <i>New Phytologist</i> , 2015, 208, 188-197.	7.3	46
30	Transcription factor interaction with COMPASS-like complex regulates histone H3K4 trimethylation for specific gene expression in plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 2900-2905.	7.1	106
31	Site-1 protease cleavage site is important for the ER stress-induced activation of membrane-associated transcription factor bZIP28 in Arabidopsis. <i>Science China Life Sciences</i> , 2015, 58, 270-275.	4.9	35
32	The Membrane-Associated Transcription Factor NAC089 Controls ER-Stress-Induced Programmed Cell Death in Plants. <i>PLoS Genetics</i> , 2014, 10, e1004243.	3.5	178
33	A plasma membrane-ethered transcription factor, <i>NAC062/ANAC062/NTL6</i> , mediates the unfolded protein response in Arabidopsis. <i>Plant Journal</i> , 2014, 79, 1033-1043.	5.7	113
34	The plant-specific transcription factor gene <i>NAC103</i> is induced by bZIP60 through a new cis-regulatory element to modulate the unfolded protein response in Arabidopsis. <i>Plant Journal</i> , 2013, 76, 274-286.	5.7	110
35	Identification and characterization of <i>OsEBS</i> , a gene involved in enhanced plant biomass and spikelet number in rice. <i>Plant Biotechnology Journal</i> , 2013, 11, 1044-1057.	8.3	23
36	The Lumen-Facing Domain Is Important for the Biological Function and Organelle-to-Organelle Movement of bZIP28 during ER Stress in Arabidopsis. <i>Molecular Plant</i> , 2013, 6, 1605-1615.	8.3	59

#	ARTICLE	IF	CITATIONS
37	Transcriptomic Analysis of Cadmium Stress Response in the Heavy Metal Hyperaccumulator <i>Sedum alfredii</i> Hance. <i>PLoS ONE</i> , 2013, 8, e64643.	2.5	100
38	Conservation of IRE1-Regulated bZIP74 mRNA Unconventional Splicing in Rice (<i>Oryza sativa</i> L.) Involved in ER Stress Responses. <i>Molecular Plant</i> , 2012, 5, 504-514.	8.3	106
39	Reversible and Irreversible Drought-Induced Changes in the Anther Proteome of Rice (<i>Oryza sativa</i> L.) Genotypes IR64 and Moroberekan. <i>Molecular Plant</i> , 2011, 4, 59-69.	8.3	65
40	Heat induces the splicing by IRE1 of a mRNA encoding a transcription factor involved in the unfolded protein response in <i>Arabidopsis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 7247-7252.	7.1	405
41	bZIP28 and NF-Y Transcription Factors Are Activated by ER Stress and Assemble into a Transcriptional Complex to Regulate Stress Response Genes in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2010, 22, 782-796.	6.6	356
42	Endoplasmic Reticulum Protein Quality Control and Its Relationship to Environmental Stress Responses in Plants. <i>Plant Cell</i> , 2010, 22, 2930-2942.	6.6	413
43	Overexpression of an <i>Arabidopsis</i> gene encoding a subtilase (AtSBT5.4) produces a clavata-like phenotype. <i>Planta</i> , 2009, 230, 687-697.	3.2	20
44	Regulation and processing of a plant peptide hormone, AtRALF23, in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2009, 59, 930-939.	5.7	174
45	Proteolytic processing of a precursor protein for a growth-promoting peptide by a subtilisin serine protease in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2008, 56, 219-227.	5.7	134
46	Stress-induced expression of an activated form of AtbZIP17 provides protection from salt stress in <i>Arabidopsis</i> . <i>Plant, Cell and Environment</i> , 2008, 31, 1735-1743.	5.7	116
47	Salt stress signaling in <i>Arabidopsis thaliana</i> involves a membrane-bound transcription factor AtbZIP17 as a signal transducer. <i>Plant Signaling and Behavior</i> , 2008, 3, 56-57.	2.4	14
48	An Endoplasmic Reticulum Stress Response in <i>Arabidopsis</i> Is Mediated by Proteolytic Processing and Nuclear Relocation of a Membrane-Associated Transcription Factor, bZIP28. <i>Plant Cell</i> , 2008, 19, 4111-4119.	6.6	394
49	Salt stress responses in <i>Arabidopsis</i> utilize a signal transduction pathway related to endoplasmic reticulum stress signaling. <i>Plant Journal</i> , 2007, 51, 897-909.	5.7	401
50	NMDA receptor antagonists reduce restraint-induced release of prolactin in male rats. <i>Neuroendocrinology Letters</i> , 2003, 24, 435-9.	0.2	1
51	Hypoxia alters testis development in neonatal rats. <i>Neuroendocrinology Letters</i> , 2002, 23, 231-7.	0.2	7