

Ronald N Jones

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10778812/publications.pdf>

Version: 2024-02-01

794
papers

44,296
citations

2565

99
h-index

8034

154
g-index

799
all docs

799
docs citations

799
times ranked

22784
citing authors

#	ARTICLE	IF	CITATIONS
1	Nosocomial Bloodstream Infections in United States Hospitals: A Three-Year Analysis. <i>Clinical Infectious Diseases</i> , 1999, 29, 239-244.	2.9	1,274
2	10 x '20 Progress--Development of New Drugs Active Against Gram-Negative Bacilli: An Update From the Infectious Diseases Society of America. <i>Clinical Infectious Diseases</i> , 2013, 56, 1685-1694.	2.9	612
3	Determining the value of antimicrobial surveillance programs. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 41, 171-175.	0.8	577
4	Microbial Etiologies of Hospital-Acquired Bacterial Pneumonia and Ventilator-Associated Bacterial Pneumonia. <i>Clinical Infectious Diseases</i> , 2010, 51, S81-S87.	2.9	570
5	Twenty Years of the SENTRY Antifungal Surveillance Program: Results for <i>Candida</i> Species From 1997-2016. <i>Open Forum Infectious Diseases</i> , 2019, 6, S79-S94.	0.4	456
6	Bacterial Pathogens Isolated from Patients with Bloodstream Infection: Frequencies of Occurrence and Antimicrobial Susceptibility Patterns from the SENTRY Antimicrobial Surveillance Program (United States and Canada, 1997). <i>Antimicrobial Agents and Chemotherapy</i> , 1998, 42, 1762-1770.	1.4	422
7	Prevalence of Antimicrobial Resistance Among Respiratory Tract Isolates of <i>Streptococcus pneumoniae</i> in North America: 1997 Results from the SENTRY Antimicrobial Surveillance Program. <i>Clinical Infectious Diseases</i> , 1998, 27, 764-770.	2.9	383
8	Oxazolidinone antibiotics. <i>Lancet</i> , The, 2001, 358, 1975-1982.	6.3	356
9	National Surveillance of Nosocomial Blood Stream Infection Due to Species of <i>Candida</i> Other than <i>Candida albicans</i> : Frequency of Occurrence and Antifungal Susceptibility in the SCOPE Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 30, 121-129.	0.8	331
10	Occurrence and antimicrobial resistance pattern comparisons among bloodstream infection isolates from the SENTRY Antimicrobial Surveillance Program (1997-2002). <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 50, 59-69.	0.8	326
11	Contemporary causes of skin and soft tissue infections in North America, Latin America, and Europe: Report from the SENTRY Antimicrobial Surveillance Program (1998-2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 7-13.	0.8	324
12	The Microbiology of Bloodstream Infection: 20-Year Trends from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2019, 63, .	1.4	307
13	Early Dissemination of NDM-1- and OXA-181-Producing <i>Enterobacteriaceae</i> in Indian Hospitals: Report from the SENTRY Antimicrobial Surveillance Program, 2006-2007. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 1274-1278.	1.4	303
14	Contemporary activity of colistin and polymyxin B against a worldwide collection of Gram-negative pathogens: results from the SENTRY Antimicrobial Surveillance Program (2006-09). <i>Journal of Antimicrobial Chemotherapy</i> , 2011, 66, 2070-2074.	1.3	295
15	Resistance Patterns Among Nosocomial Pathogens. <i>Chest</i> , 2001, 119, 397S-404S.	0.4	285
16	Antimicrobial resistance and molecular epidemiology of vancomycin-resistant enterococci from North America and Europe: a report from the SENTRY antimicrobial surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 58, 163-170.	0.8	280
17	Molecular characterization of SPM-1, a novel metallo-beta-lactamase isolated in Latin America: report from the SENTRY antimicrobial surveillance programme. <i>Journal of Antimicrobial Chemotherapy</i> , 2002, 50, 673-679.	1.3	277
18	Molecular Characterization of a β -Lactamase Gene, bla GIM-1 , Encoding a New Subclass of Metallo- β -Lactamase. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 4654-4661.	1.4	236

#	ARTICLE	IF	CITATIONS
19	Antimicrobial Susceptibility and Epidemiology of a Worldwide Collection of <i>Chryseobacterium</i> spp.: Report from the SENTRY Antimicrobial Surveillance Program (1997-2001). <i>Journal of Clinical Microbiology</i> , 2004, 42, 445-448.	1.8	230
20	Antimicrobial resistance among Gram-negative bacilli isolated from Latin America: results from SENTRY Antimicrobial Surveillance Program (Latin America, 2008-2010). <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 73, 354-360.	0.8	222
21	Microbiological Features of Vancomycin in the 21st Century: Minimum Inhibitory Concentration Creep, Bactericidal/Static Activity, and Applied Breakpoints to Predict Clinical Outcomes or Detect Resistant Strains. <i>Clinical Infectious Diseases</i> , 2006, 42, S13-S24.	2.9	218
22	<i>Candida</i> bloodstream infections: comparison of species distribution and resistance to echinocandin and azole antifungal agents in Intensive Care Unit (ICU) and non-ICU settings in the SENTRY Antimicrobial Surveillance Program (2008-2009). <i>International Journal of Antimicrobial Agents</i> , 2011, 38, 65-69.	1.1	216
23	Nosocomial Bloodstream Infections Caused by <i>Acinetobacter</i> Species in United States Hospitals: Clinical Features, Molecular Epidemiology, and Antimicrobial Susceptibility. <i>Clinical Infectious Diseases</i> , 2000, 31, 690-697.	2.9	215
24	Global Emergence of Trimethoprim/Sulfamethoxazole Resistance in <i>Stenotrophomonas maltophilia</i> Mediated by Acquisition of <i>sul</i> Genes. <i>Emerging Infectious Diseases</i> , 2007, 13, 559-565.	2.0	210
25	Echinocandin and Triazole Antifungal Susceptibility Profiles for Clinical Opportunistic Yeast and Mold Isolates Collected from 2010 to 2011: Application of New CLSI Clinical Breakpoints and Epidemiological Cutoff Values for Characterization of Geographic and Temporal Trends of Antifungal Resistance. <i>Journal of Clinical Microbiology</i> , 2013, 51, 2571-2581.	1.8	209
26	<i>Candida</i> Bloodstream Infections: Comparison of Species Distributions and Antifungal Resistance Patterns in Community-Onset and Nosocomial Isolates in the SENTRY Antimicrobial Surveillance Program, 2008-2009. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 561-566.	1.4	204
27	First Report of <i>cfr</i> -Mediated Resistance to Linezolid in Human Staphylococcal Clinical Isolates Recovered in the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 2244-2246.	1.4	203
28	Oxazolidinones. <i>Drugs</i> , 2000, 59, 7-16.	4.9	198
29	Linezolid update: Stable in vitro activity following more than a decade of clinical use and summary of associated resistance mechanisms. <i>Drug Resistance Updates</i> , 2014, 17, 1-12.	6.5	195
30	Antimicrobial susceptibility of Gram-negative organisms isolated from patients hospitalised with pneumonia in US and European hospitals: Results from the SENTRY Antimicrobial Surveillance Program, 2009-2012. <i>International Journal of Antimicrobial Agents</i> , 2014, 43, 328-334.	1.1	194
31	Geographic Variations in Species Distribution and Echinocandin and Azole Antifungal Resistance Rates among <i>Candida</i> Bloodstream Infection Isolates: Report from the SENTRY Antimicrobial Surveillance Program (2008 to 2009). <i>Journal of Clinical Microbiology</i> , 2011, 49, 396-399.	1.8	192
32	Epidemiologic typing of multiply drug-resistant <i>Pseudomonas aeruginosa</i> isolated from an outbreak in an intensive care unit. <i>Diagnostic Microbiology and Infectious Disease</i> , 1993, 17, 13-18.	0.8	188
33	Antimicrobial susceptibility of Gram-negative organisms isolated from patients hospitalized in intensive care units in United States and European hospitals (2009-2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 78, 443-448.	0.8	184
34	Antimicrobial activity and spectrum of the new glycolcycline, GAR-936 tested against 1,203 recent clinical bacterial isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 36, 19-36.	0.8	177
35	Antimicrobial Activity of Ceftolozane-Tazobactam Tested against Enterobacteriaceae and <i>Pseudomonas aeruginosa</i> with Various Resistance Patterns Isolated in U.S. Hospitals (2011-2012). <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 6305-6310.	1.4	177
36	Epidemiology and carbapenem resistance mechanisms of carbapenem-non-susceptible <i>Pseudomonas aeruginosa</i> collected during 2009-11 in 14 European and Mediterranean countries. <i>Journal of Antimicrobial Chemotherapy</i> , 2014, 69, 1804-1814.	1.3	173

#	ARTICLE	IF	CITATIONS
37	A nationwide, multicenter, case-control study comparing risk factors, treatment, and outcome for vancomycin-resistant and -susceptible enterococcal bacteremia†. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 36, 145-158.	0.8	170
38	Contemporary Diversity of Î²-Lactamases among Enterobacteriaceae in the Nine U.S. Census Regions and Ceftazidime-Avibactam Activity Tested against Isolates Producing the Most Prevalent Î²-Lactamase Groups. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 833-838.	1.4	170
39	Rationale for Revised Penicillin Susceptibility Breakpoints versus <i>Streptococcus pneumoniae</i> : Coping with Antimicrobial Susceptibility in an Era of Resistance. <i>Clinical Infectious Diseases</i> , 2009, 48, 1596-1600.	2.9	163
40	Assessment of pathogen occurrences and resistance profiles among infected patients in the intensive care unit: report from the SENTRY Antimicrobial Surveillance Program (North America, 2001). <i>International Journal of Antimicrobial Agents</i> , 2004, 24, 111-118.	1.1	162
41	<i>Haemophilus influenzae</i> and <i>Moraxella catarrhalis</i> from Patients with Community-Acquired Respiratory Tract Infections: Antimicrobial Susceptibility Patterns from the SENTRY Antimicrobial Surveillance Program (United States and Canada, 1997). <i>Antimicrobial Agents and Chemotherapy</i> , 1999, 43, 385-389.	1.4	161
42	Molecular Analysis of Tn 1546 in <i>Enterococcus faecium</i> Isolated from Animals and Humans. <i>Journal of Clinical Microbiology</i> , 1998, 36, 437-442.	1.8	161
43	Multicenter Studies of Tigecycline Disk Diffusion Susceptibility Results for <i>Acinetobacter</i> spp. <i>Journal of Clinical Microbiology</i> , 2007, 45, 227-230.	1.8	157
44	Occurrence and antimicrobial susceptibility patterns of pathogens isolated from skin and soft tissue infections: report from the SENTRY Antimicrobial Surveillance Program (United States and Canada, 1999-2008). <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 1078-1084.	1.1	156
45	Summary trends for the Meropenem Yearly Susceptibility Test Information Collection Program: a 10-year experience in the United States (1999-2008). <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 65, 414-426.	0.8	156
46	Antimicrobial susceptibility of uncommonly isolated non-enteric Gram-negative bacilli. <i>International Journal of Antimicrobial Agents</i> , 2005, 25, 95-109.	1.1	155
47	Emerging multiply resistant enterococci among clinical isolates I. Prevalence data from 97 medical center surveillance study in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 85-93.	0.8	152
48	Antimicrobial Activity of Quinupristin-Dalfopristin (RP 59500, Synercid®) Tested against Over 28,000 Recent Clinical Isolates from 200 Medical Centers in the United States and Canada. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 437-451.	0.8	152
49	Linezolid Resistance since 2001: SENTRY Antimicrobial Surveillance Program. <i>Annals of Pharmacotherapy</i> , 2003, 37, 769-774.	0.9	151
50	<i>In vitro</i> antimicrobial activity of S-649266, a catechol-substituted siderophore cephalosporin, when tested against non-fermenting Gram-negative bacteria. <i>Journal of Antimicrobial Chemotherapy</i> , 2016, 71, 670-677.	1.3	150
51	Nosocomial enterococcal blood stream infections in the SCOPE program: Antimicrobial resistance, species occurrence, molecular testing results, and laboratory testing accuracy. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 95-102.	0.8	148
52	International Surveillance of <i>Candida</i> spp. and <i>Aspergillus</i> spp.: Report from the SENTRY Antimicrobial Surveillance Program (2003). <i>Journal of Clinical Microbiology</i> , 2006, 44, 1782-1787.	1.8	146
53	Prevalence of important pathogens and antimicrobial activity of parenteral drugs at numerous medical centers in the United States I. Study on the threat of emerging resistances: Real or perceived?. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 19, 203-215.	0.8	144
54	Global Epidemiology of Antimicrobial Resistance among Community-Acquired and Nosocomial Pathogens: A Five-Year Summary from the SENTRY Antimicrobial Surveillance Program (1997-2001). <i>Seminars in Respiratory and Critical Care Medicine</i> , 2003, 24, 121-134.	0.8	144

#	ARTICLE	IF	CITATIONS
55	Variation in <i>Candida</i> spp. distribution and antifungal resistance rates among bloodstream infection isolates by patient age: report from the SENTRY Antimicrobial Surveillance Program (2008–2009). <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 68, 278-283.	0.8	141
56	Emergence and widespread dissemination of OXA-23, -24/40 and -58 carbapenemases among <i>Acinetobacter</i> spp. in Asia-Pacific nations: report from the SENTRY Surveillance Program. <i>Journal of Antimicrobial Chemotherapy</i> , 2008, 63, 55-59.	1.3	139
57	Impact of changing pathogens and antimicrobial susceptibility patterns in the treatment of serious infections in hospitalized patients. <i>American Journal of Medicine</i> , 1996, 100, 3S-12S.	0.6	137
58	Antimicrobial Activity and Spectrum of PPI-0903M (T-91825), a Novel Cephalosporin, Tested against a Worldwide Collection of Clinical Strains. <i>Antimicrobial Agents and Chemotherapy</i> , 2005, 49, 3501-3512.	1.4	137
59	Worldwide assessment of dalbavancin activity and spectrum against over 6,000 clinical isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 48, 137-143.	0.8	136
60	Regional variation in the prevalence of extended-spectrum β -lactamase-producing clinical isolates in the Asia-Pacific region (SENTRY 1998–2002). <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 323-329.	0.8	136
61	Antimicrobial Susceptibility of <i>Acinetobacter calcoaceticus</i> – <i>Acinetobacter baumannii</i> Complex and <i>Stenotrophomonas maltophilia</i> Clinical Isolates: Results From the SENTRY Antimicrobial Surveillance Program (1997–2016). <i>Open Forum Infectious Diseases</i> , 2019, 6, S34-S46.	0.4	136
62	Emerging Resistance to Antimicrobial Agents in Gram-Positive Bacteria. <i>Drugs</i> , 1996, 51, 6-12.	4.9	135
63	Activity and spectrum of 22 antimicrobial agents tested against urinary tract infection pathogens in hospitalized patients in Latin America: report from the second year of the SENTRY Antimicrobial Surveillance Program (1998). <i>Journal of Antimicrobial Chemotherapy</i> , 2000, 45, 295-303.	1.3	134
64	Bacterial pathogens isolated from patients with skin and soft tissue infections: frequency of occurrence and antimicrobial susceptibility patterns from the SENTRY Antimicrobial Surveillance Program (United States and Canada, 1997). <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 65-72.	0.8	133
65	Occurrence and Characterization of Carbapenemase-Producing Enterobacteriaceae: Report from the SENTRY Antimicrobial Surveillance Program (2000–2004). <i>Microbial Drug Resistance</i> , 2006, 12, 223-230.	0.9	133
66	Characterization of Vancomycin-Heteroresistant <i>Staphylococcus aureus</i> from the Metropolitan Area of Detroit, Michigan, over a 22-Year Period (1986 to 2007). <i>Journal of Clinical Microbiology</i> , 2008, 46, 2950-2954.	1.8	132
67	Background and Rationale for Revised Clinical and Laboratory Standards Institute Interpretive Criteria (Breakpoints) for Enterobacteriaceae and <i>Pseudomonas aeruginosa</i> . <i>Cephalosporins and Aztreonam</i> . <i>Clinical Infectious Diseases</i> , 2013, 56, 1301-1309.	2.9	132
68	Twenty-Year Trends in Antimicrobial Susceptibilities Among <i>Staphylococcus aureus</i> From the SENTRY Antimicrobial Surveillance Program. <i>Open Forum Infectious Diseases</i> , 2019, 6, S47-S53.	0.4	132
69	Antimicrobial Activities of Tigecycline and Other Broad-Spectrum Antimicrobials Tested against Serine Carbapenemase- and Metallo- β -Lactamase-Producing Enterobacteriaceae: Report from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 570-573.	1.4	131
70	Daptomycin activity and spectrum: a worldwide sample of 6737 clinical Gram-positive organisms. <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 53, 669-674.	1.3	130
71	Antimicrobial activity of ceftolozane/tazobactam tested against <i>Pseudomonas aeruginosa</i> and Enterobacteriaceae with various resistance patterns isolated in European hospitals (2011–12). <i>Journal of Antimicrobial Chemotherapy</i> , 2014, 69, 2713-2722.	1.3	130
72	bla VIM-7, an Evolutionarily Distinct Metallo- β -Lactamase Gene in a <i>Pseudomonas aeruginosa</i> Isolate from the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 329-332.	1.4	129

#	ARTICLE	IF	CITATIONS
73	Antimicrobial Activity of Ceftazidime-Avibactam against Gram-Negative Organisms Collected from U.S. Medical Centers in 2012. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 1684-1692.	1.4	129
74	OXA-163, an OXA-48-Related Class D β -Lactamase with Extended Activity Toward Expanded-Spectrum Cephalosporins. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 2546-2551.	1.4	128
75	Assessment of linezolid resistance mechanisms among <i>Staphylococcus epidermidis</i> causing bacteraemia in Rome, Italy. <i>Journal of Antimicrobial Chemotherapy</i> , 2010, 65, 2329-2335.	1.3	126
76	Emergence of serine carbapenemases (KPC and SME) among clinical strains of Enterobacteriaceae isolated in the United States Medical Centers: Report from the MYSTIC Program (1999-2005). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 367-372.	0.8	124
77	Detection of a New <i>bla</i> -Like Gene, <i>bla</i> (B), in <i>Enterococcus faecium</i> Isolates Recovered from Human Specimens in the United States as Part of the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 6256-6261.	1.4	124
78	Variations in the Occurrence of Resistance Phenotypes and Carbapenemase Genes Among Enterobacteriaceae Isolates in 20 Years of the SENTRY Antimicrobial Surveillance Program. <i>Open Forum Infectious Diseases</i> , 2019, 6, S23-S33.	0.4	124
79	Bacterial Resistance: A Worldwide Problem. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 379-388.	0.8	123
80	Ceftolozane/tazobactam activity tested against Gram-negative bacterial isolates from hospitalised patients with pneumonia in US and European medical centres (2012). <i>International Journal of Antimicrobial Agents</i> , 2014, 43, 533-539.	1.1	123
81	Pathogen of occurrence and susceptibility patterns associated with pneumonia in hospitalized patients in North America: results of the SENTRY Antimicrobial Surveillance Study (2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 45, 279-285.	0.8	122
82	Effect of the β -Lactamase Inhibitor Vaborbactam Combined with Meropenem against Serine Carbapenemase-Producing Enterobacteriaceae. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 5454-5458.	1.4	121
83	Characteristics of pathogens causing urinary tract infections in hospitals in North America: results from the SENTRY Antimicrobial Surveillance Program, 1997. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 55-63.	0.8	120
84	Clarithromycin, a unique macrolide. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 39-53.	0.8	118
85	Mutation-Driven β -Lactam Resistance Mechanisms among Contemporary Ceftazidime-Nonsusceptible <i>Pseudomonas aeruginosa</i> Isolates from U.S. Hospitals. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 6844-6850.	1.4	118
86	In vitro evaluation of BAL9141, a novel parenteral cephalosporin active against oxacillin-resistant staphylococci. <i>Journal of Antimicrobial Chemotherapy</i> , 2002, 50, 915-932.	1.3	117
87	Comparative activity of doripenem and three other carbapenems tested against Gram-negative bacilli with various β -lactamase resistance mechanisms. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 71-74.	0.8	117
88	<i>Staphylococcus aureus</i> and Coagulase-Negative Staphylococci from Blood Stream Infections: Frequency of Occurrence, Antimicrobial Susceptibility, and Molecular (<i>mecA</i>) Characterization of Oxacillin Resistance in the SCOPE Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 30, 205-214.	0.8	116
89	Omiganan Pentahydrochloride (MBI 226), a Topical 12-Amino-Acid Cationic Peptide: Spectrum of Antimicrobial Activity and Measurements of Bactericidal Activity. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 3112-3118.	1.4	115
90	Evaluation of Vancomycin and Daptomycin Potency Trends (MIC Creep) against Methicillin-Resistant <i>Staphylococcus aureus</i> Isolates Collected in Nine U.S. Medical Centers from 2002 to 2006. <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 4127-4132.	1.4	113

#	ARTICLE	IF	CITATIONS
91	Activity of Retapamulin (SB-275833), a Novel Pleuromutilin, against Selected Resistant Gram-Positive Cocci. <i>Antimicrobial Agents and Chemotherapy</i> , 2006, 50, 2583-2586.	1.4	112
92	Low Prevalence of <i>fls1</i> Hot Spot 1 Mutations in a Worldwide Collection of <i>Candida</i> Strains. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 2655-2659.	1.4	112
93	Antimicrobial Activity of CXA-101, a Novel Cephalosporin Tested in Combination with Tazobactam against Enterobacteriaceae, <i>Pseudomonas aeruginosa</i> , and <i>Bacteroides fragilis</i> Strains Having Various Resistance Phenotypes. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 2390-2394.	1.4	112
94	Inducible amp C β -lactamase producing gram-negative bacilli from blood stream infections: Frequency, antimicrobial susceptibility, and molecular epidemiology in a national surveillance program (SCOPE). <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 28, 211-219.	0.8	111
95	Prevalence of extended spectrum β -lactamase (ESBL)-producing clinical isolates in the Asia-Pacific region and South Africa: regional results from SENTRY Antimicrobial Surveillance Program (1998-99). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 42, 193-198.	0.8	111
96	Activities of Doripenem (S-4661) against Drug-Resistant Clinical Pathogens. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 3136-3140.	1.4	110
97	Doripenem (S-4661), a novel carbapenem: comparative activity against contemporary pathogens including bactericidal action and preliminary in vitro methods evaluations. <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 54, 144-154.	1.3	110
98	Tigecycline activity tested against 26,474 bloodstream infection isolates: a collection from 6 continents. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 181-186.	0.8	106
99	Ceftazidime-Avibactam Activity Tested against Enterobacteriaceae Isolates from U.S. Hospitals (2011 to 2015). <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 3509-3517.	1.4	104
100	Antimicrobial susceptibility patterns for pathogens isolated from patients in Latin American medical centers with a diagnosis of pneumonia: analysis of results from the SENTRY Antimicrobial Surveillance Program (1997). <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 32, 289-301.	0.8	103
101	Group B streptococci causing neonatal bloodstream infection: Antimicrobial susceptibility and serotyping results from SENTRY centers in the Western Hemisphere. <i>American Journal of Obstetrics and Gynecology</i> , 2000, 183, 859-862.	0.7	103
102	SENTRY antimicrobial surveillance program report: latin american and brazilian results for 1997 through 2001. <i>Brazilian Journal of Infectious Diseases</i> , 2004, 8, 25-79.	0.3	101
103	Susceptibility rates in Latin American nations: report from a regional resistance surveillance program (2011). <i>Brazilian Journal of Infectious Diseases</i> , 2013, 17, 672-681.	0.3	101
104	Contemporary in vitro spectrum of activity summary for antimicrobial agents tested against 18,569 strains non-fermentative Gram-negative bacilli isolated in the SENTRY Antimicrobial Surveillance Program (1997-2001). <i>International Journal of Antimicrobial Agents</i> , 2003, 22, 551-556.	1.1	100
105	LEADER surveillance program results for 2006: an activity and spectrum analysis of linezolid using clinical isolates from the United States (50 medical centers). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 59, 309-317.	0.8	100
106	Prevalence of β -Lactamase-Encoding Genes among Enterobacteriaceae Bacteremia Isolates Collected in 26 U.S. Hospitals: Report from the SENTRY Antimicrobial Surveillance Program (2010). <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 3012-3020.	1.4	100
107	Update of dalbavancin spectrum and potency in the USA: report from the SENTRY Antimicrobial Surveillance Program (2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 304-307.	0.8	100
108	Antimicrobial Activity and Spectrum Investigation of Eight Broad-Spectrum β -Lactam Drugs: A 1997 Surveillance Trial in 102 Medical Centers in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 30, 215-228.	0.8	99

#	ARTICLE	IF	CITATIONS
109	Important and Emerging β -Lactamase-mediated Resistances in Hospital-based Pathogens: The Amp C Enzymes. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 461-466.	0.8	97
110	In vitro antimicrobial activity of GAR-936 tested against antibiotic-resistant gram-positive blood stream infection isolates and strains producing extended-spectrum β -lactamases. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 40, 173-177.	0.8	97
111	Pathogen frequency and resistance patterns in Brazilian hospitals: summary of results from three years of the SENTRY antimicrobial surveillance program. <i>Brazilian Journal of Infectious Diseases</i> , 2001, 5, 200-14.	0.3	97
112	Characterization of methicillin-resistant <i>Staphylococcus aureus</i> displaying increased MICs of ceftaroline. <i>Journal of Antimicrobial Chemotherapy</i> , 2012, 67, 1321-1324.	1.3	97
113	Echinocandin and triazole antifungal susceptibility profiles for <i>Candida</i> spp., <i>Cryptococcus neoformans</i> , and <i>Aspergillus fumigatus</i> : application of new CLSI clinical breakpoints and epidemiologic cutoff values to characterize resistance in the SENTRY Antimicrobial Surveillance Program (2009). <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 69, 45-50.	0.8	96
114	Antimicrobial usage and resistance trend relationships from the MYSTIC Programme in North America (1999-2001). <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 53, 290-296.	1.3	95
115	Susceptibility patterns of orally administered antimicrobials among urinary tract infection pathogens from hospitalized patients in North America: comparison report to Europe and Latin America. Results from the SENTRY Antimicrobial Surveillance Program (2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 45, 295-301.	0.8	94
116	Antimicrobial activity of tigecycline tested against nosocomial bacterial pathogens from patients hospitalized in the intensive care unit. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 203-208.	0.8	94
117	Dissemination and diversity of metallo- β -lactamases in Latin America: report from the SENTRY Antimicrobial Surveillance Program. <i>International Journal of Antimicrobial Agents</i> , 2005, 25, 57-61.	1.1	93
118	Antimicrobial Susceptibilities of a Worldwide Collection of <i>Stenotrophomonas maltophilia</i> Isolates Tested against Tigecycline and Agents Commonly Used for <i>S. maltophilia</i> Infections. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 2735-2737.	1.4	93
119	Linezolid-resistant <i>Enterococcus faecium</i> isolated from a patient without prior exposure to an oxazolidinone: report from the SENTRY Antimicrobial Surveillance Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 42, 137-139.	0.8	92
120	Urinary tract infection trends in Latin American hospitals: report from the SENTRY antimicrobial surveillance program (1997-2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 289-299.	0.8	92
121	First Descriptions of <i>bla</i> KPC in <i>Raoultella</i> spp. (<i>R. planticola</i> and) Tj ETQq1 1 0.784314 rgBT /Overl Clinical Microbiology, 2009, 47, 4129-4130.	1.8	92
122	The emergent needs for basic research, education, and surveillance of antimicrobial resistance. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 25, 153-161.	0.8	91
123	Antimicrobial activity of ceftobiprole, a novel anti-methicillin-resistant <i>Staphylococcus aureus</i> cephalosporin, tested against contemporary pathogens: results from the SENTRY Antimicrobial Surveillance Program (2005-2006). <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 61, 86-95.	0.8	90
124	Antimicrobial susceptibility of Gram-positive bacteria isolated from US medical centers: results of the Daptomycin Surveillance Program (2007-2008). <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 65, 158-162.	0.8	90
125	Antimicrobial activity and spectrum of rifaximin, a new topical rifamycin derivative. <i>Diagnostic Microbiology and Infectious Disease</i> , 1993, 16, 111-118.	0.8	89
126	Increasing prevalence of antimicrobial resistance among <i>Pseudomonas aeruginosa</i> isolates in Latin American medical centres: 5 year report of the SENTRY Antimicrobial Surveillance Program (1997-2001). <i>Journal of Antimicrobial Chemotherapy</i> , 2003, 52, 140-141.	1.3	89

#	ARTICLE	IF	CITATIONS
127	Occurrence and molecular characterization of fusidic acid resistance mechanisms among <i>Staphylococcus</i> spp. from European countries (2008). <i>Journal of Antimicrobial Chemotherapy</i> , 2010, 65, 1353-1358.	1.3	89
128	Assessment of pathogen frequency and resistance patterns among pediatric patient isolates: Report from the 2004 SENTRY Antimicrobial Surveillance Program on 3 continents. <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 427-436.	0.8	88
129	<i>Candida guilliermondii</i> and Other Species of <i>Candida</i> Misidentified as <i>Candida famata</i> : Assessment by Vitek 2, DNA Sequencing Analysis, and Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry in Two Global Antifungal Surveillance Programs. <i>Journal of Clinical Microbiology</i> , 2013, 51, 117-124.	1.8	88
130	Potential synergy activity of the novel ceragenin, CSA-13, against clinical isolates of <i>Pseudomonas aeruginosa</i> , including multidrug-resistant <i>P. aeruginosa</i> . <i>Journal of Antimicrobial Chemotherapy</i> , 2007, 61, 365-370.	1.3	87
131	Nine-Hospital Study Comparing Broth Microdilution and Etest Method Results for Vancomycin and Daptomycin against Methicillin-Resistant <i>Staphylococcus aureus</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 3162-3165.	1.4	87
132	Activity of MK-3118, a new oral glucan synthase inhibitor, tested against <i>Candida</i> spp. by two international methods (CLSI and EUCAST). <i>Journal of Antimicrobial Chemotherapy</i> , 2013, 68, 858-863.	1.3	87
133	Integron Carrying a Novel Metallo- β -Lactamase Gene, blaIMP-16, and a Fused Form of Aminoglycoside-Resistant Gene aac(6)-30/aac(6)-Ib: Report from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 4693-4702.	1.4	86
134	Activity of a long-acting echinocandin, CD101, determined using CLSI and EUCAST reference methods, against <i>Candida</i> and <i>Aspergillus</i> spp., including echinocandin- and azole-resistant isolates. <i>Journal of Antimicrobial Chemotherapy</i> , 2016, 71, 2868-2873.	1.3	85
135	Ceftazidime, a pseudomonas-active cephalosporin: in-vitro antimicrobial activity evaluation including recommendations for disc diffusion susceptibility tests. <i>Journal of Antimicrobial Chemotherapy</i> , 1981, 8, 187-211.	1.3	84
136	Geographic and Temporal Patterns of Antimicrobial Resistance in <i>Pseudomonas aeruginosa</i> Over 20 Years From the SENTRY Antimicrobial Surveillance Program, 1997-2016. <i>Open Forum Infectious Diseases</i> , 2019, 6, S63-S68.	0.4	84
137	Italian metallo- β -lactamases: a national problem? Report from the SENTRY Antimicrobial Surveillance Programme. <i>Journal of Antimicrobial Chemotherapy</i> , 2005, 55, 61-70.	1.3	83
138	Four-year evaluation of frequency of occurrence and antimicrobial susceptibility patterns of bacteria from bloodstream infections in Latin American medical centers. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 273-280.	0.8	82
139	Evolving trends in <i>Streptococcus pneumoniae</i> resistance: implications for therapy of community-acquired bacterial pneumonia. <i>International Journal of Antimicrobial Agents</i> , 2010, 36, 197-204.	1.1	82
140	Activities of E1210 and Comparator Agents Tested by CLSI and EUCAST Broth Microdilution Methods against <i>Fusarium</i> and <i>Scedosporium</i> Species Identified Using Molecular Methods. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 352-357.	1.4	82
141	Regional data analysis of <i>Candida albicans</i> strains collected in United States medical sites over a 6-year period, 2006-2011. <i>Mycoses</i> , 2014, 57, 602-611.	1.8	82
142	In Vitro Activity of Newer Fluoroquinolones for Respiratory Tract Infections and Emerging Patterns of Antimicrobial Resistance: Data from the SENTRY Antimicrobial Surveillance Program. <i>Clinical Infectious Diseases</i> , 2000, 31, S16-S23.	2.9	81
143	Genetic characterization of a novel metallo- β -lactamase gene, blaIMP-13, harboured by a novel Tn5051-type transposon disseminating carbapenemase genes in Europe: report from the SENTRY worldwide antimicrobial surveillance programme. <i>Journal of Antimicrobial Chemotherapy</i> , 2003, 52, 583-590.	1.3	81
144	In Vitro Activities of the Novel Cephalosporin LB 11058 against Multidrug-Resistant <i>Staphylococci</i> and <i>Streptococci</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 53-62.	1.4	81

#	ARTICLE	IF	CITATIONS
145	Antimicrobial activity of the novel pleuromutilin antibiotic BC-3781 against organisms responsible for community-acquired respiratory tract infections (CARTIs). <i>Journal of Antimicrobial Chemotherapy</i> , 2012, 67, 1170-1175.	1.3	81
146	WCK 5222 (cefepime/zidebactam) antimicrobial activity tested against Gram-negative organisms producing clinically relevant β -lactamases. <i>Journal of Antimicrobial Chemotherapy</i> , 2017, 72, 1696-1703.	1.3	81
147	Antimicrobial activity of RU-66647, a new ketolide. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 27, 7-12.	0.8	80
148	United States resistance surveillance results for linezolid (LEADER Program for 2007). <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 62, 416-426.	0.8	80
149	Update on antimicrobial susceptibility trends among <i>Streptococcus pneumoniae</i> in the United States: report of ceftaroline activity from the SENTRY Antimicrobial Surveillance Program (1998-2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 107-109.	0.8	80
150	<i>In Vitro</i> Activity of a New Oral Glucan Synthase Inhibitor (MK-3118) Tested against <i>Aspergillus</i> spp. by CLSI and EUCAST Broth Microdilution Methods. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 1065-1068.	1.4	80
151	Linezolid Surveillance Results for the United States (LEADER Surveillance Program 2014). <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 2273-2280.	1.4	80
152	Rapid Emergence of β -CTX-M Among Enterobacteriaceae in U.S. Medical Centers: Molecular Evaluation from the MYSTIC Program (2007). <i>Microbial Drug Resistance</i> , 2008, 14, 211-216.	0.9	79
153	LEADER Program Results for 2009: an Activity and Spectrum Analysis of Linezolid Using 6,414 Clinical Isolates from 56 Medical Centers in the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 3684-3690.	1.4	79
154	Pharmacokinetics-Pharmacodynamics of Tazobactam in Combination with Ceftolozane in an <i>In Vitro</i> Infection Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 2809-2814.	1.4	79
155	Antimicrobial activity of tigecycline tested against organisms causing community-acquired respiratory tract infection and nosocomial pneumonia. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 187-193.	0.8	78
156	<i>In Vitro</i> Activity of Ceftaroline Against Multidrug-Resistant <i>Staphylococcus aureus</i> and <i>Streptococcus pneumoniae</i> : A Review of Published Studies and the AWARE Surveillance Program (2008-2010). <i>Clinical Infectious Diseases</i> , 2012, 55, S206-S214.	2.9	78
157	Resistance surveillance program report for selected European nations (2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 78, 429-436.	0.8	78
158	Daptomycin Bactericidal Activity and Correlation between Disk and Broth Microdilution Method Results in Testing of <i>Staphylococcus aureus</i> Strains with Decreased Susceptibility to Vancomycin. <i>Antimicrobial Agents and Chemotherapy</i> , 2006, 50, 2330-2336.	1.4	77
159	Fusidic Acid Resistance Rates and Prevalence of Resistance Mechanisms among <i>Staphylococcus</i> spp. Isolated in North America and Australia, 2007-2008. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 3614-3617.	1.4	77
160	Activities of Dalbavancin against a Worldwide Collection of 81,673 Gram-Positive Bacterial Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 1260-1263.	1.4	76
161	Linezolid surveillance program results for 2008 (LEADER Program for 2008). <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 65, 392-403.	0.8	76
162	Daptomycin activity tested against 164457 bacterial isolates from hospitalised patients: Summary of 8 years of a Worldwide Surveillance Programme (2005-2012). <i>International Journal of Antimicrobial Agents</i> , 2014, 43, 465-469.	1.1	76

#	ARTICLE	IF	CITATIONS
163	Contemporary Antimicrobial Susceptibility Patterns of Bacterial Pathogens Commonly Associated with Febrile Patients with Neutropenia. <i>Clinical Infectious Diseases</i> , 1999, 29, 495-502.	2.9	75
164	<i>In Vitro</i> Activities of Isavuconazole and Comparator Antifungal Agents Tested against a Global Collection of Opportunistic Yeasts and Molds. <i>Journal of Clinical Microbiology</i> , 2013, 51, 2608-2616.	1.8	75
165	Ceftolozane/tazobactam activity tested against aerobic Gram-negative organisms isolated from intra-abdominal and urinary tract infections in European and United States hospitals (2012). <i>Journal of Infection</i> , 2014, 69, 266-277.	1.7	75
166	Ceftazidime/avibactam tested against Gram-negative bacteria from intensive care unit (ICU) and non-ICU patients, including those with ventilator-associated pneumonia. <i>International Journal of Antimicrobial Agents</i> , 2015, 46, 53-59.	1.1	75
167	Geographic variations in activity of broad-spectrum β -lactams against <i>Pseudomonas aeruginosa</i> : summary of the worldwide SENTRY Antimicrobial Surveillance Program (1997-2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 239-243.	0.8	74
168	In vitro activity of tigecycline (GAR-936) tested against 11,859 recent clinical isolates associated with community-acquired respiratory tract and gram-positive cutaneous infections. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 201-209.	0.8	74
169	Antimicrobial Activities of Ceftaroline and ME1036 Tested against Clinical Strains of Community-Acquired Methicillin-Resistant <i>Staphylococcus aureus</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 1153-1155.	1.4	74
170	Regional Resistance Surveillance Program Results for 12 Asia-Pacific Nations (2011). <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 5721-5726.	1.4	74
171	Ceftazidime-Avibactam Activity against Multidrug-Resistant <i>Pseudomonas aeruginosa</i> Isolated in U.S. Medical Centers in 2012 and 2013. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 3656-3659.	1.4	74
172	Comparison of Agar Diffusion Methodologies for Antimicrobial Susceptibility Testing of <i>Pseudomonas aeruginosa</i> Isolates from Cystic Fibrosis Patients. <i>Journal of Clinical Microbiology</i> , 2000, 38, 1818-1822.	1.8	74
173	Emergence of multidrug-resistant <i>Streptococcus pneumoniae</i> : report from the SENTRY Antimicrobial Surveillance Program (1999-2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 69-74.	0.8	73
174	Antimicrobial susceptibility pattern comparisons among intensive care unit and general ward Gram-negative isolates from the Meropenem Yearly Susceptibility Test Information Collection Program (USA). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 57-62.	0.8	73
175	Serotype Replacement and Multiple Resistance in <i>Streptococcus pneumoniae</i> After the Introduction of the Conjugate Pneumococcal Vaccine. <i>Microbial Drug Resistance</i> , 2008, 14, 101-107.	0.9	73
176	Zyvox [®] Annual Appraisal of Potency and Spectrum program: linezolid surveillance program results for 2008. <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 65, 404-413.	0.8	73
177	Antimicrobial Activity of the Investigational Pleuromutilin Compound BC-3781 Tested against Gram-Positive Organisms Commonly Associated with Acute Bacterial Skin and Skin Structure Infections. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 1619-1623.	1.4	73
178	Antimicrobial Activity of the Pleuromutilin Antibiotic BC-3781 against Bacterial Pathogens Isolated in the SENTRY Antimicrobial Surveillance Program in 2010. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 4489-4495.	1.4	73
179	Evaluation of Vancomycin Susceptibility Testing for Methicillin-Resistant <i>Staphylococcus aureus</i> : Comparison of Etest and Three Automated Testing Methods. <i>Journal of Clinical Microbiology</i> , 2013, 51, 2077-2081.	1.8	73
180	Public Health Consequences of Macrolide Use in Food Animals: A Deterministic Risk Assessment. <i>Journal of Food Protection</i> , 2004, 67, 980-992.	0.8	72

#	ARTICLE	IF	CITATIONS
181	Emerging Metallo- β -Lactamase-Mediated Resistances: A Summary Report from the Worldwide SENTRY Antimicrobial Surveillance Program. <i>Clinical Infectious Diseases</i> , 2005, 41, S276-S278.	2.9	72
182	Significance of Molecular Identification and Antifungal Susceptibility of Clinically Significant Yeasts and Moulds in a Global Antifungal Surveillance Programme. <i>Mycopathologia</i> , 2012, 174, 259-271.	1.3	72
183	Frequency of pathogen occurrence and antimicrobial susceptibility among community-acquired respiratory tract infections in the respiratory surveillance program study: microbiology from the medical office practice environment. <i>American Journal of Medicine</i> , 2001, 111, 4-12.	0.6	71
184	Oxazolidinone susceptibility patterns in 2004: report from the Zyvox [®] Annual Appraisal of Potency and Spectrum (ZAAPS) Program assessing isolates from 16 nations. <i>Journal of Antimicrobial Chemotherapy</i> , 2006, 57, 279-287.	1.3	71
185	Summary of Ceftaroline Activity against Pathogens in the United States, 2010: Report from the Assessing Worldwide Antimicrobial Resistance Evaluation (AWARE) Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 2933-2940.	1.4	71
186	Multi-laboratory assessment of the linezolid spectrum of activity using the Kirby-Bauer disk diffusion method: Report of the Zyvox [®] Antimicrobial Potency Study (ZAPS) in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 40, 59-66.	0.8	70
187	A multicenter evaluation of linezolid antimicrobial activity in North America. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 75-83.	0.8	70
188	Antimicrobial susceptibility patterns of β -hemolytic and viridans group streptococci: report from the SENTRY Antimicrobial Surveillance Program (1997-2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 157-162.	0.8	70
189	Increased resistance to first-line agents among bacterial pathogens isolated from urinary tract infections in Latin America: time for local guidelines?. <i>Memorias Do Instituto Oswaldo Cruz</i> , 2006, 101, 741-748.	0.8	70
190	Activity of Ceftaroline-Avibactam Tested against Gram-Negative Organism Populations, including Strains Expressing One or More β -Lactamases and Methicillin-Resistant <i>Staphylococcus aureus</i> Carrying Various Staphylococcal Cassette Chromosome <i>mec</i> Types. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 4779-4785.	1.4	70
191	Temporal and Geographic Variation in Antimicrobial Susceptibility and Resistance Patterns of Enterococci: Results From the SENTRY Antimicrobial Surveillance Program, 1997-2016. <i>Open Forum Infectious Diseases</i> , 2019, 6, S54-S62.	0.4	70
192	Methods for Improved Detection of Oxacillin Resistance in Coagulase-Negative Staphylococci: Results of a Multicenter Study. <i>Journal of Clinical Microbiology</i> , 1999, 37, 4051-4058.	1.8	70
193	Dalbavancin activity against selected populations of antimicrobial-resistant Gram-positive pathogens. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 53, 307-310.	0.8	69
194	Carbapenem Resistance among <i>Pseudomonas aeruginosa</i> Strains from India: Evidence for Nationwide Endemicity of Multiple Metallo- β -Lactamase Clones (VIM-2, -5, -6, and -11 and the Newly) <i>Trends Microbiology</i> , 2007, 15, 107-111.	0.8	68
195	AWARE Ceftaroline Surveillance Program (2008-2010): Trends in Resistance Patterns Among <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> , and <i>Moraxella catarrhalis</i> in the United States. <i>Clinical Infectious Diseases</i> , 2012, 55, S187-S193.	2.9	68
196	Antibacterial activity of 41 antimicrobials tested against over 2773 bacterial isolates from hospitalized patients with pneumonia: results from the SENTRY Antimicrobial Surveillance Program (North) <i>Journal of Clinical Microbiology</i> , 2004, 42, 107-111.	0.8	67
197	Spectrum and potency evaluation of a new oxazolidinone, linezolid: report from the SENTRY Antimicrobial Surveillance Program, 1998-2000. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 65-73.	0.8	67
198	Activity of Ceftaroline against Recent Emerging Serotypes of <i>Streptococcus pneumoniae</i> in the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 2716-2719.	1.4	67

#	ARTICLE	IF	CITATIONS
199	Frequency of <i>FKS</i> Mutations among <i>Candida glabrata</i> Isolates from a 10-Year Global Collection of Bloodstream Infection Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 577-580.	1.4	67
200	Zyvox(R) Annual Appraisal of Potency and Spectrum (ZAAPS) Program: report of linezolid activity over 9 years (2004-12). <i>Journal of Antimicrobial Chemotherapy</i> , 2014, 69, 1582-1588.	1.3	67
201	A review of the in vitro activity of meropenem and comparative antimicrobial agents tested against 30,254 aerobic and anaerobic pathogens isolated world wide. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 28, 157-163.	0.8	66
202	Determination of the antimicrobial activity of 29 clinically important compounds tested against fastidious HACEK group organisms. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 73-76.	0.8	66
203	Comparison of <i>Streptococcus pneumoniae</i> and <i>Haemophilus influenzae</i> susceptibilities from community-acquired respiratory tract infections and hospitalized patients with pneumonia: five-year results for the SENTRY antimicrobial surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 46, 285-289.	0.8	66
204	Biochemical Characterization of the Acquired Metallo- β -Lactamase SPM-1 from <i>Pseudomonas aeruginosa</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 582-587.	1.4	66
205	Resistance trends of <i>Acinetobacter</i> spp. in Latin America and characterization of international dissemination of multi-drug resistant strains: five-year report of the SENTRY Antimicrobial Surveillance Program. <i>International Journal of Infectious Diseases</i> , 2004, 8, 284-291.	1.5	66
206	Activity of Contemporary Antifungal Agents, Including the Novel Echinocandin Anidulafungin, Tested against <i>Candida</i> spp., <i>Cryptococcus</i> spp., and <i>Aspergillus</i> spp.: Report from the SENTRY Antimicrobial Surveillance Program (2006 to 2007). <i>Journal of Clinical Microbiology</i> , 2009, 47, 1942-1946.	1.8	66
207	ZAAPS International Surveillance Program (2007) for linezolid resistance: results from 5591 Gram-positive clinical isolates in 23 countries. <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 64, 191-201.	0.8	66
208	In Vitro Activity of a Novel Broad-Spectrum Antifungal, E1210, Tested against <i>Aspergillus</i> spp. Determined by CLSI and EUCAST Broth Microdilution Methods. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 5155-5158.	1.4	66
209	Pharmacodynamics of β -Lactamase Inhibition by NXL104 in Combination with Ceftaroline: Examining Organisms with Multiple Types of β -Lactamases. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 258-270.	1.4	66
210	Antimicrobial activity of daptomycin against multidrug-resistant gram-positive strains collected worldwide. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 50, 201-204.	0.8	65
211	Trends in carbapenemase-producing <i>Escherichia coli</i> and <i>Klebsiella</i> spp. from Europe and the Americas: report from the SENTRY antimicrobial surveillance programme (2007-09). <i>Journal of Antimicrobial Chemotherapy</i> , 2011, 66, 1409-1411.	1.3	65
212	Use of a surfactant (polysorbate 80) to improve MIC susceptibility testing results for polymyxin B and colistin. <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 74, 412-414.	0.8	65
213	Cefuroxime, a New Parenteral Cephalosporin: Collaborative In Vitro Susceptibility Comparison with Cephalothin Against 5,887 Clinical Bacterial Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 1977, 12, 47-50.	1.4	64
214	Skin and soft tissue infections in Latin American medical centers: four-year assessment of the pathogen frequency and antimicrobial susceptibility patterns. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 281-288.	0.8	64
215	Tigecycline activity tested against antimicrobial resistant surveillance subsets of clinical bacteria collected worldwide (2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 217-221.	0.8	64
216	Activity of echinocandins and triazoles against a contemporary (2012) worldwide collection of yeast and moulds collected from invasive infections. <i>International Journal of Antimicrobial Agents</i> , 2014, 44, 320-326.	1.1	64

#	ARTICLE	IF	CITATIONS
217	Comparison of EUCAST and CLSI broth microdilution methods for the susceptibility testing of 10 Systemically active antifungal agents when tested against <i>Candida</i> spp.. <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 79, 198-204.	0.8	64
218	Activity of tigecycline tested against a global collection of Enterobacteriaceae, including tetracycline-resistant isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 209-213.	0.8	63
219	Antimicrobial susceptibility of Gram-positive bacterial isolates from the Asia-Pacific region and an in vitro evaluation of the bactericidal activity of daptomycin, vancomycin, and teicoplanin: a SENTRY Program Report (2003-2004). <i>International Journal of Antimicrobial Agents</i> , 2007, 30, 143-149.	1.1	63
220	Antimicrobial activities of doripenem and other carbapenems against <i>Pseudomonas aeruginosa</i> , other nonfermentative bacilli, and <i>Aeromonas</i> spp.. <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 63, 426-433.	0.8	63
221	Surveillance for linezolid resistance via the Zyvox Annual Appraisal of Potency and Spectrum (ZAAPS) programme (2014): evolving resistance mechanisms with stable susceptibility rates. <i>Journal of Antimicrobial Chemotherapy</i> , 2016, 71, 1860-1865.	1.3	63
222	WCK 5222 (Cefepime-Zidebactam) Antimicrobial Activity against Clinical Isolates of Gram-Negative Bacteria Collected Worldwide in 2015. <i>Antimicrobial Agents and Chemotherapy</i> , 2017, 61, .	1.4	63
223	Respiratory tract pathogens isolated from patients hospitalized with suspected pneumonia in Latin America: frequency of occurrence and antimicrobial susceptibility profile: results from the SENTRY Antimicrobial Surveillance Program (1997-2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 301-311.	0.8	62
224	Emergence of an IMP-like metallo-enzyme in an <i>Acinetobacter baumannii</i> clinical strain from a Brazilian teaching hospital. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 45, 77-79.	0.8	62
225	<i>Pseudomonas aeruginosa</i> strains harbouring an unusual blaVIM-4 gene cassette isolated from hospitalized children in Poland (1998-2001). <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 53, 451-456.	1.3	62
226	Pharmacological Basis of β -Lactamase Inhibitor Therapeutics: Tazobactam in Combination with Ceftolozane. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 5924-5930.	1.4	62
227	Ceftobiprole Activity against over 60,000 Clinical Bacterial Pathogens Isolated in Europe, Turkey, and Israel from 2005 to 2010. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 3882-3888.	1.4	62
228	Cefoperazone: A Review of Its Antimicrobial Spectrum, β -Lactamase Stability, Enzyme Inhibition, and Other in Vitro Characteristics. <i>Clinical Infectious Diseases</i> , 1983, 5, S108-S126.	2.9	61
229	A 1994-1995 survey of <i>Haemophilus influenzae</i> susceptibility to ten orally administered agents A 187 clinical laboratory center sample in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 27, 75-83.	0.8	61
230	Evaluation of the in vitro activity of daptomycin against 19615 clinical isolates of Gram-positive cocci collected in North American hospitals (2002-2005). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 459-465.	0.8	61
231	Ceftaroline activity against pathogens associated with complicated skin and skin structure infections: results from an international surveillance study. <i>Journal of Antimicrobial Chemotherapy</i> , 2010, 65, iv17-iv31.	1.3	61
232	Ceftazidime/avibactam activity tested against Gram-negative bacteria isolated from bloodstream, pneumonia, intra-abdominal and urinary tract infections in US medical centres (2012). <i>Journal of Antimicrobial Chemotherapy</i> , 2014, 69, 1589-1598.	1.3	61
233	Geographic variations and trends in antimicrobial resistance among <i>Enterococcus faecalis</i> and <i>Enterococcus faecium</i> in the SENTRY Antimicrobial Surveillance Program (1997-2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 46, 63-68.	0.8	60
234	Characterization of fluoroquinolone-resistant β -hemolytic <i>Streptococcus</i> spp. isolated in North America and Europe including the first report of fluoroquinolone-resistant <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i> : Report from the SENTRY Antimicrobial Surveillance Program (1997-2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 55, 119-127.	0.8	60

#	ARTICLE	IF	CITATIONS
235	Characterization of an Integron Carrying bla IMP-1 and a New Aminoglycoside Resistance Gene, aac(6)-31, and Its Dissemination among Genetically Unrelated Clinical Isolates in a Brazilian Hospital. <i>Antimicrobial Agents and Chemotherapy</i> , 2007, 51, 2611-2614.	1.4	60
236	Comparison of two commercial systems (Vitek and MicroScan-WalkAway) for antimicrobial susceptibility testing of <i>Pseudomonas aeruginosa</i> isolates from cystic fibrosis patients. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 39, 257-260.	0.8	59
237	Evaluation of the in vitro activity of six broad-spectrum β -lactam antimicrobial agents tested against recent clinical isolates from India: a survey of ten medical center laboratories. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 367-377.	0.8	59
238	Triazole and Echinocandin MIC Distributions with Epidemiological Cutoff Values for Differentiation of Wild-Type Strains from Non-Wild-Type Strains of Six Uncommon Species of <i>Candida</i> . <i>Journal of Clinical Microbiology</i> , 2011, 49, 3800-3804.	1.8	59
239	Use of Micafungin as a Surrogate Marker To Predict Susceptibility and Resistance to Caspofungin among 3,764 Clinical Isolates of <i>Candida</i> by Use of CLSI Methods and Interpretive Criteria. <i>Journal of Clinical Microbiology</i> , 2014, 52, 108-114.	1.8	59
240	<i>In Vitro</i> Spectrum of Pexiganan Activity When Tested against Pathogens from Diabetic Foot Infections and with Selected Resistance Mechanisms. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 1751-1754.	1.4	59
241	Sustained activity and spectrum of selected extended-spectrum β -lactams (carbapenems and cefepime) against <i>Enterobacter</i> spp. and ESBL-producing <i>Klebsiella</i> spp.: report from the SENTRY antimicrobial surveillance program (USA, 1997-2000). <i>International Journal of Antimicrobial Agents</i> , 2003, 21, 1-7.	1.1	58
242	Worldwide Antimicrobial Susceptibility Patterns and Pharmacodynamic Comparisons of Gatifloxacin and Levofloxacin against <i>Streptococcus pneumoniae</i> : Report from the Antimicrobial Resistance Rate Epidemiology Study Team. <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 292-296.	1.4	58
243	Evolution of an integron carrying blaVIM-2 in Eastern Europe: report from the SENTRY Antimicrobial Surveillance Program. <i>Journal of Antimicrobial Chemotherapy</i> , 2003, 52, 116-119.	1.3	58
244	Accuracy of Three Automated Systems (MicroScan WalkAway, VITEK, and VITEK 2) for Susceptibility Testing of <i>Pseudomonas aeruginosa</i> against Five Broad-Spectrum Beta-Lactam Agents. <i>Journal of Clinical Microbiology</i> , 2006, 44, 1101-1104.	1.8	58
245	Binaphthyl-Based Dicationic Peptoids with Therapeutic Potential. <i>Angewandte Chemie - International Edition</i> , 2010, 49, 537-540.	7.2	58
246	Tigecycline activity tested against carbapenem-resistant Enterobacteriaceae from 18 European nations: results from the SENTRY surveillance program (2010-2013). <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 83, 183-186.	0.8	58
247	An international activity and spectrum analysis of linezolid: ZAAPS Program results for 2011. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 206-213.	0.8	57
248	Macrolide and Fluoroquinolone (Levofloxacin) Resistances among <i>Streptococcus pneumoniae</i> Strains: Significant Trends from the SENTRY Antimicrobial Surveillance Program (North America, 1997-1999). <i>Journal of Clinical Microbiology</i> , 2000, 38, 4298-4299.	1.8	57
249	Susceptibility trends of <i>Haemophilus influenzae</i> and <i>Moraxella catarrhalis</i> against orally administered antimicrobial agents: five-year report from the SENTRY Antimicrobial Surveillance Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 47, 373-376.	0.8	56
250	Development of decreased susceptibility to daptomycin and vancomycin in a <i>Staphylococcus aureus</i> strain during prolonged therapy. <i>Journal of Antimicrobial Chemotherapy</i> , 2006, 58, 481-483.	1.3	56
251	Occurrence of vancomycin-tolerant and heterogeneous vancomycin-intermediate strains (hVISA) among <i>Staphylococcus aureus</i> causing bloodstream infections in nine USA hospitals. <i>Journal of Antimicrobial Chemotherapy</i> , 2009, 64, 1024-1028.	1.3	56
252	First Report of Staphylococcal Clinical Isolates in Mexico with Linezolid Resistance Caused by <i>In Vivo</i> Mobilization. <i>Journal of Clinical Microbiology</i> , 2010, 48, 3041-3043.	1.8	56

#	ARTICLE	IF	CITATIONS
253	Wild-type MIC distributions and epidemiologic cutoff values for fluconazole, posaconazole, and voriconazole when testing <i>Cryptococcus neoformans</i> as determined by the CLSI broth microdilution method. <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 71, 252-259.	0.8	56
254	Antimicrobial Susceptibility of <i>Streptococcus pneumoniae</i> from North America, Europe, Latin America, and the Asia-Pacific Region: Results From 20 Years of the SENTRY Antimicrobial Surveillance Program (1997â€“2016). <i>Open Forum Infectious Diseases</i> , 2019, 6, S14-S23.	0.4	56
255	E test, an antimicrobial susceptibility testing method with broad clinical and epidemiologic application. <i>Antimicrobial Newsletter</i> , 1992, 8, 1-7.	0.9	55
256	Genetic Relatedness of Multidrug-Resistant, Methicillin (Oxacillin)-Resistant <i>Staphylococcus aureus</i> Bloodstream Isolates from SENTRY Antimicrobial Resistance Surveillance Centers Worldwide, 1998. <i>Microbial Drug Resistance</i> , 2000, 6, 213-221.	0.9	55
257	Potency and spectrum trends for cefepime tested against 65746 clinical bacterial isolates collected in North American medical centers: Results from the SENTRY Antimicrobial Surveillance Program (1998â€“2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 265-273.	0.8	55
258	Potency and spectrum of tigecycline tested against an international collection of bacterial pathogens associated with skin and soft tissue infections (2000â€“2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 195-201.	0.8	55
259	Trends in linezolid susceptibility patterns: report from the 2002â€“2003 worldwide Zyvox Annual Appraisal of Potency and Spectrum (ZAAPS) Program. <i>International Journal of Antimicrobial Agents</i> , 2005, 26, 13-21.	1.1	55
260	Spectrum and potency of dalbavancin tested against 3322 Gram-positive cocci isolated in the United States Surveillance Program (2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 54, 149-153.	0.8	55
261	Outcomes evaluation of patients with ESBL- and non-ESBL-producing <i>Escherichia coli</i> and <i>Klebsiella</i> species as defined by CLSI reference methods: report from the SENTRY Antimicrobial Surveillance Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 54, 231-236.	0.8	55
262	Daptomycin antimicrobial activity tested against methicillin-resistant staphylococci and vancomycin-resistant enterococci isolated in European medical centers (2005). <i>BMC Infectious Diseases</i> , 2007, 7, 29.	1.3	55
263	Comparative activity of meropenem in US medical centers (2007): initiating the 2nd decade of MYSTIC program surveillance. <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 61, 203-213.	0.8	55
264	High Rates of Nonsusceptibility to Ceftazidime-avibactam and Identification of New Delhi Metallo- β -lactamase Production in <i>Enterobacteriaceae</i> Bloodstream Infections at a Major Cancer Center: Table 1.. <i>Clinical Infectious Diseases</i> , 2016, 63, 954-958.	2.9	55
265	The prevalence of staphylococcal resistance to penicillinase-resistant penicillins. <i>Diagnostic Microbiology and Infectious Disease</i> , 1989, 12, 385-394.	0.8	54
266	The current and future impact of antimicrobial resistance among nosocomial bacterial pathogens. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 1.	0.8	54
267	Occurrence of macrolide-lincosamide-streptogramin resistances among staphylococcal clinical isolates at a University Medical Center. <i>Diagnostic Microbiology and Infectious Disease</i> , 1993, 16, 205-213.	0.8	54
268	Antimicrobial resistance trends in medical centers using carbapenems: report of 1999 and 2000 results from the MYSTIC program (USA). <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 41, 177-182.	0.8	54
269	Antimicrobial spectrum of activity for meropenem and nine broad spectrum antimicrobials: report from the MYSTIC Program (2002) in North America. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 47, 365-372.	0.8	54
270	Declining antimicrobial susceptibility of <i>Streptococcus pneumoniae</i> in the United States: report from the SENTRY Antimicrobial Surveillance Program (1998â€“2009). <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 68, 334-336.	0.8	54

#	ARTICLE	IF	CITATIONS
271	Rapid Expansion of KPC-2-Producing <i>Klebsiella pneumoniae</i> Isolates in Two Texas Hospitals due to Clonal Spread of ST258 and ST307 Lineages. <i>Microbial Drug Resistance</i> , 2013, 19, 295-297.	0.9	54
272	Piperacillin/tazobactam (YTR 830) combination. <i>Diagnostic Microbiology and Infectious Disease</i> , 1989, 12, 489-494.	0.8	53
273	Respiratory tract pathogens isolated from patients hospitalized with suspected pneumonia: frequency of occurrence and antimicrobial susceptibility patterns from the SENTRY Antimicrobial Surveillance Program (United States and Canada, 1997). <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 37, 115-125.	0.8	53
274	Relationship between increased levofloxacin use and decreased susceptibility of <i>Streptococcus pneumoniae</i> in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 51, 31-37.	0.8	53
275	Carbapenem-Resistant Isolates of <i>Klebsiella pneumoniae</i> in China and Detection of a Conjugative Plasmid (<i>bla</i> _{KPC-2} plus <i>qnrB4</i>) and a <i>bla</i> _{IMP-4} Gene. <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 798-799.	1.4	53
276	In vitro activity of a novel broad-spectrum antifungal, E1210, tested against <i>Candida</i> spp. as determined by CLSI broth microdilution method. <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 71, 167-170.	0.8	53
277	Evaluation of Clonality and Carbapenem Resistance Mechanisms among <i>Acinetobacter baumannii</i> - <i>Acinetobacter calcoaceticus</i> Complex and Enterobacteriaceae Isolates Collected in European and Mediterranean Countries and Detection of Two Novel β -Lactamases, GES-22 and VIM-35. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 7358-7366.	1.4	53
278	Changes in the Frequencies of β -Lactamase Genes among Enterobacteriaceae Isolates in U.S. Hospitals, 2012 to 2014: Activity of Ceftazidime-Avibactam Tested against β -Lactamase-Producing Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 4770-4777.	1.4	53
279	Performance Accuracy of Antibacterial and Antifungal Susceptibility Test Methods: Report From the College of American Pathologists Microbiology Surveys Program (2001-2003). <i>Archives of Pathology and Laboratory Medicine</i> , 2006, 130, 767-778.	1.2	53
280	Integrating <i>Escherichia coli</i> Antimicrobial Susceptibility Data from Multiple Surveillance Programs. <i>Emerging Infectious Diseases</i> , 2005, 11, 873-882.	2.0	52
281	Comparative Antimicrobial Characterization of LBM415 (NVP PDF-713), a New Peptide Deformylase Inhibitor of Clinical Importance. <i>Antimicrobial Agents and Chemotherapy</i> , 2005, 49, 1468-1476.	1.4	52
282	Antimicrobial susceptibility of Gram-positive cocci isolated from skin and skin-structure infections in European medical centres. <i>International Journal of Antimicrobial Agents</i> , 2010, 36, 28-32.	1.1	52
283	Comparison of 30-min and 3-h infusion regimens for imipenem/cilastatin and for meropenem evaluated by Monte Carlo simulation. <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 68, 251-258.	0.8	52
284	Tigecycline antimicrobial activity tested against clinical bacteria from Latin American medical centres: results from SENTRY Antimicrobial Surveillance Program (2011-2014). <i>International Journal of Antimicrobial Agents</i> , 2016, 48, 144-150.	1.1	52
285	Nosocomial streptococcal blood stream infections in the SCOPE program: Species occurrence and antimicrobial resistance. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 259-263.	0.8	51
286	In Vitro Activity of Ceftaroline Alone and in Combination against Clinical Isolates of Resistant Gram-Negative Pathogens, Including β -Lactamase-Producing Enterobacteriaceae and <i>Pseudomonas aeruginosa</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 2360-2366.	1.4	51
287	Comparison of the Vitek Gram-Positive Susceptibility 106 Card and the MRSA-Screen Latex Agglutination Test for Determining Oxacillin Resistance in Clinical Bloodstream Isolates of <i>Staphylococcus aureus</i> . <i>Journal of Clinical Microbiology</i> , 2001, 39, 53-56.	1.8	50
288	Initial results from a longitudinal international surveillance programme for anidulafungin (2003). <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 54, 1051-1056.	1.3	50

#	ARTICLE	IF	CITATIONS
289	Trends in <i>Klebsiella pneumoniae</i> carbapenemase-positive <i>K. pneumoniae</i> in US hospitals: report from the 2007–2009 SENTRY Antimicrobial Surveillance Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 356-360.	0.8	50
290	Antifungal susceptibilities of <i>Candida</i> , <i>Cryptococcus neoformans</i> and <i>Aspergillus fumigatus</i> from the Asia and Western Pacific region: data from the SENTRY antifungal surveillance program (2010–2012). <i>Journal of Antibiotics</i> , 2015, 68, 556-561.	1.0	50
291	Comparative Antimicrobial Activity and Kill-Curve Investigations of Novel Ketolide Antimicrobial Agents (HMR 3004 and HMR 3647) Tested against <i>Haemophilus influenzae</i> and <i>Moraxella catarrhalis</i> Strains. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 349-353.	0.8	49
292	Global patterns of susceptibility for 21 commonly utilized antimicrobial agents tested against 48,440 Enterobacteriaceae in the SENTRY Antimicrobial Surveillance Program (1997-2001). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 47, 361-364.	0.8	49
293	Analysis of <i>Salmonella</i> spp. with resistance to extended-spectrum cephalosporins and fluoroquinolones isolated in North America and Latin America: report from the SENTRY Antimicrobial Surveillance Program (1997–2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 54, 13-21.	0.8	49
294	Antimicrobial activity of daptomycin tested against Gram-positive pathogens collected in Europe, Latin America, and selected countries in the Asia-Pacific Region (2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 417-422.	0.8	49
295	The Importance of Antimicrobial Resistance Monitoring Worldwide and the Origins of SENTRY Antimicrobial Surveillance Program. <i>Open Forum Infectious Diseases</i> , 2019, 6, S1-S4.	0.4	49
296	Application of an In Vitro Infection Model and Simulation for Reevaluation of Fluoroquinolone Breakpoints for <i>Salmonella enterica</i> Serotype Typhi. <i>Antimicrobial Agents and Chemotherapy</i> , 2005, 49, 1775-1781.	1.4	48
297	CEM-101, a novel fluoroketolide: antimicrobial activity against a diverse collection of Gram-positive and Gram-negative bacteria. <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 66, 393-401.	0.8	48
298	Unmet Needs and Prospects for Oritavancin in the Management of Vancomycin-Resistant Enterococcal Infections. <i>Clinical Infectious Diseases</i> , 2012, 54, S233-S238.	2.9	48
299	Linezolid Surveillance Results for the United States: LEADER Surveillance Program 2011. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 1077-1081.	1.4	48
300	Antimicrobial activity of cefpirome an update compared to five third-generation cephalosporins against nearly 6000 recent clinical isolates from five medical centers. <i>Diagnostic Microbiology and Infectious Disease</i> , 1991, 14, 361-364.	0.8	47
301	The E-Test applied to susceptibility tests of gonococci, multiply-resistant enterococci, and enterobacteriaceae producing potent β -lactamases. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 459-463.	0.8	47
302	Prevalence of important pathogens and the antimicrobial activity of parenteral drugs at numerous medical centers in the united states II. Study of the intra- and interlaboratory dissemination of extended-spectrum β -lactamase-producing Enterobacteriaceae. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 20, 203-208.	0.8	47
303	Practical approach to the identification of clinically relevant <i>Enterococcus</i> species. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 165-171.	0.8	47
304	Two-year assessment of the pathogen frequency and antimicrobial resistance patterns among organisms isolated from skin and soft tissue infections in latin American hospitals: Results from the SENTRY antimicrobial surveillance program, 1997-98. <i>International Journal of Infectious Diseases</i> , 2000, 4, 75-84.	1.5	47
305	Molecular and Biochemical Characterization of OXA-45, an Extended-Spectrum Class 2 β -Lactamase in <i>Pseudomonas aeruginosa</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 2859-2863.	1.4	47
306	Prevalence and Significance of a Negative Extended-Spectrum β -Lactamase (ESBL) Confirmation Test Result after a Positive ESBL Screening Test Result for Isolates of <i>Escherichia coli</i> and <i>Klebsiella pneumoniae</i> : Results from the SENTRY Asia-Pacific Surveillance Program. <i>Journal of Clinical Microbiology</i> , 2007, 45, 1478-1482.	1.8	47

#	ARTICLE	IF	CITATIONS
307	Contemporary activity of meropenem and comparator broad-spectrum agents: MYSTIC program report from the United States component (2005). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 207-215.	0.8	47
308	Pathogen occurrence and antimicrobial resistance trends among urinary tract infection isolates in the Asia-Western Pacific Region: report from the SENTRY Antimicrobial Surveillance Program, 1998-1999. <i>International Journal of Antimicrobial Agents</i> , 2002, 20, 10-17.	1.1	46
309	Pre-clinical development of antifungal susceptibility test methods for the testing of the novel antifungal agent E1210 versus <i>Candida</i> : comparison of CLSI and European Committee on Antimicrobial Susceptibility Testing methods. <i>Journal of Antimicrobial Chemotherapy</i> , 2011, 66, 2581-2584.	1.3	46
310	Antimicrobial activity of cefepime tested against bush group I β -lactamase-producing strains resistant to ceftazidime a multilaboratory national and international clinical isolate study. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 19, 33-38.	0.8	45
311	Fluoroquinolone-resistant <i>Haemophilus influenzae</i> : Frequency of occurrence and analysis of confirmed strains in the SENTRY antimicrobial surveillance program (North and Latin America). <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 36, 255-259.	0.8	45
312	Five-year analysis of <i>Haemophilus influenzae</i> isolates with reduced susceptibility to fluoroquinolones: prevalence results from the SENTRY antimicrobial surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 46, 55-61.	0.8	45
313	First Isolation of bla VIM-2 in Latin America: Report from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 1433-1434.	1.4	45
314	Antimicrobial Susceptibility Patterns of KPC-Producing or CTX-M-Producing Enterobacteriaceae. <i>Microbial Drug Resistance</i> , 2010, 16, 61-65.	0.9	45
315	LEADER Surveillance program results for 2010: an activity and spectrum analysis of linezolid using 6801 clinical isolates from the United States (61 medical centers). <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 74, 54-61.	0.8	45
316	Avibactam reverts the ceftazidime MIC90 of European Gram-negative bacterial clinical isolates to the epidemiological cut-off value. <i>Journal of Chemotherapy</i> , 2014, 26, 333-338.	0.7	45
317	Serotype distribution and antimicrobial susceptibility of USA <i>Streptococcus pneumoniae</i> isolates collected prior to and post introduction of 13-valent pneumococcal conjugate vaccine. <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 80, 19-25.	0.8	45
318	Application of Next-Generation Sequencing for Characterization of Surveillance and Clinical Trial Isolates: Analysis of the Distribution of β -lactamase Resistance Genes and Lineage Background in the United States. <i>Open Forum Infectious Diseases</i> , 2019, 6, S69-S78.	0.4	45
319	First Description of <i>Klebsiella pneumoniae</i> Harboring CTX-M β -Lactamases (CTX-M-14 and CTX-M-3) in Taiwan. <i>Antimicrobial Agents and Chemotherapy</i> , 2002, 46, 1098-1100.	1.4	44
320	Bactericidal activity of BAL9141, a novel parenteral cephalosporin against contemporary Gram-positive and Gram-negative isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 50, 73-75.	0.8	44
321	Evaluation of alternative disk diffusion methods for detecting <i>mecA</i> -mediated oxacillin resistance in an international collection of staphylococci: Validation report from the SENTRY antimicrobial surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 51, 57-62.	0.8	44
322	Comparative activities of cefepime and piperacillin/tazobactam tested against a global collection of <i>Escherichia coli</i> and <i>Klebsiella</i> spp. with an ESBL phenotype. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 341-344.	0.8	44
323	Antimicrobial activity of omiganan pentahydrochloride tested against contemporary bacterial pathogens commonly responsible for catheter-associated infections. <i>Journal of Antimicrobial Chemotherapy</i> , 2008, 61, 1092-1098.	1.3	44
324	Antimicrobial susceptibility of daptomycin and comparator agents tested against methicillin-resistant <i>Staphylococcus aureus</i> and vancomycin-resistant enterococci: trend analysis of a 6-year period in US medical centers (2005-2010). <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 70, 412-416.	0.8	44

#	ARTICLE	IF	CITATIONS
325	Ceftazidime-avibactam and comparator agents tested against urinary tract isolates from a global surveillance program (2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 80, 233-238.	0.8	44
326	Summary of Linezolid Activity and Resistance Mechanisms Detected during the 2012 LEADER Surveillance Program for the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 1243-1247.	1.4	44
327	Use of Anidulafungin as a Surrogate Marker To Predict Susceptibility and Resistance to Caspofungin among 4,290 Clinical Isolates of <i>Candida</i> by Using CLSI Methods and Interpretive Criteria. <i>Journal of Clinical Microbiology</i> , 2014, 52, 3223-3229.	1.8	44
328	<i>In Vitro</i> Activity of Dalbavancin against Drug-Resistant <i>Staphylococcus aureus</i> Isolates from a Global Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 5007-5009.	1.4	44
329	Evaluation of Reference Dilution Test Methods for Antimicrobial Susceptibility Testing of <i>Pseudomonas aeruginosa</i> Strains Isolated from Patients with Cystic Fibrosis. <i>Journal of Clinical Microbiology</i> , 1999, 37, 2987-2991.	1.8	44
330	In vitro evaluation of a novel orally administered cephalosporin (cefditoren) tested against 1249 recent clinical isolates of <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i> , and <i>Streptococcus pneumoniae</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 573-578.	0.8	43
331	Molecular epidemiology of selected multidrug-resistant bacteria: A global report from the SENTRY Antimicrobial Surveillance Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 231-236.	0.8	43
332	Trends in linezolid susceptibility patterns in 2002: Report from the worldwide Zyvox Annual Appraisal of Potency and Spectrum Program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 53-58.	0.8	43
333	Occurrence of plasmidic AmpC type β -lactamase-mediated resistance in <i>Escherichia coli</i> : report from the SENTRY Antimicrobial Surveillance Program (North America, 2004). <i>International Journal of Antimicrobial Agents</i> , 2006, 28, 578-581.	1.1	43
334	Antimicrobial Activity of Omigaman Pentahydrochloride against Contemporary Fungal Pathogens Responsible for Catheter-Associated Infections. <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 1187-1189.	1.4	43
335	Antimicrobial characterisation of CEM-101 activity against respiratory tract pathogens, including multidrug-resistant pneumococcal serogroup 19A isolates. <i>International Journal of Antimicrobial Agents</i> , 2010, 35, 537-543.	1.1	43
336	Comparative ceftaroline activity tested against pathogens associated with community-acquired pneumonia: results from an international surveillance study. <i>Journal of Antimicrobial Chemotherapy</i> , 2011, 66, iii69-iii80.	1.3	43
337	Antimicrobial Activity of Ceftaroline Tested against Staphylococci with Reduced Susceptibility to Linezolid, Daptomycin, or Vancomycin from U.S. Hospitals, 2008 to 2011. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 3178-3181.	1.4	43
338	In vitro activity of three carbapenem antibiotics comparative studies with biapenem (L-627), imipenem, and meropenem against aerobic pathogens isolated worldwide. <i>Diagnostic Microbiology and Infectious Disease</i> , 1993, 17, 299-305.	0.8	42
339	Antimicrobial activity of 12 broad-spectrum agents tested against 270 nosocomial blood stream infection isolates caused by non-Enteric Gram-negative bacilli: Occurrence of resistance, molecular epidemiology, and screening for metallo-enzymes. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 187-192.	0.8	42
340	Antimicrobial activity of gatifloxacin (AM-1155, CG5501), and four other fluoroquinolones tested against 2,284 recent clinical strains of <i>Streptococcus pneumoniae</i> from Europe, Latin America, Canada, and the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 315-320.	0.8	42
341	Molecular Epidemiology of Macrolide Resistance in Neonatal Bloodstream Isolates of Group B <i>Streptococci</i> . <i>Journal of Clinical Microbiology</i> , 2003, 41, 2659-2661.	1.8	42
342	Factors Influencing Broth Microdilution Antimicrobial Susceptibility Test Results for Dalbavancin, a New Glycopeptide Agent. <i>Journal of Clinical Microbiology</i> , 2007, 45, 3151-3154.	1.8	42

#	ARTICLE	IF	CITATIONS
343	Update of the in vitro activity of daptomycin tested against 6710 Gram-positive cocci isolated in North America (2006). <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 61, 235-239.	0.8	42
344	Revised Reference Broth Microdilution Method for Testing Telavancin: Effect on MIC Results and Correlation with Other Testing Methodologies. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 5547-5551.	1.4	42
345	Isavuconazole and Nine Comparator Antifungal Susceptibility Profiles for Common and Uncommon <i>Candida</i> Species Collected in 2012: Application of New CLSI Clinical Breakpoints and Epidemiological Cutoff Values. <i>Mycopathologia</i> , 2014, 178, 1-9.	1.3	42
346	Update of the telavancin activity in vitro tested against a worldwide collection of Gram-positive clinical isolates (2013), when applying the revised susceptibility testing method. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 81, 275-279.	0.8	42
347	BAY 12-8039, a novel fluoroquinolone. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 32, 45-50.	0.8	41
348	Zyvox® Annual Appraisal of Potency and Spectrum Program Results for 2006: an activity and spectrum analysis of linezolid using clinical isolates from 16 countries. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 59, 199-209.	0.8	41
349	Antimicrobial activity of tigecycline against community-acquired methicillin-resistant <i>Staphylococcus aureus</i> isolates recovered from North American medical centers. <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 60, 433-436.	0.8	41
350	Antipseudomonal activity of piperacillin/tazobactam: more than a decade of experience from the SENTRY Antimicrobial Surveillance Program (1997-2007). <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 65, 331-334.	0.8	41
351	Comprehensive assessment of tigecycline activity tested against a worldwide collection of <i>Acinetobacter</i> spp. (2005-2009). <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 68, 307-311.	0.8	41
352	Variation in Potency and Spectrum of Tigecycline Activity against Bacterial Strains from U.S. Medical Centers since Its Approval for Clinical Use (2006 to 2012). <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 2274-2280.	1.4	41
353	Fluoroquinolone resistance an evolving national problem or just a problem for some physicians?. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 177-179.	0.8	40
354	The comparative antimicrobial activity of levofloxacin tested against 350 clinical isolates of streptococci. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 25, 47-51.	0.8	40
355	Comparative activity and spectrum of broad-spectrum β -lactams (cefepime, ceftazidime, ceftriaxone,) Tj ETQq1 1 0.784314 rgBT /Overl antimicrobial surveillance program (North America: 2001-2002). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 47, 435-440.	0.8	40
356	Evolution and dissemination of extended-spectrum β -lactamase-producing <i>Klebsiella pneumoniae</i> : Epidemiology and molecular report from the SENTRY Antimicrobial Surveillance Program (1997-2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 51, 1-7.	0.8	40
357	Characterization of Baseline Methicillin-Resistant <i>Staphylococcus aureus</i> Isolates Recovered from Phase IV Clinical Trial for Linezolid. <i>Journal of Clinical Microbiology</i> , 2010, 48, 568-574.	1.8	40
358	Dissemination of a pSCFS3-Like <i>cfrc</i> -Carrying Plasmid in <i>Staphylococcus aureus</i> and <i>Staphylococcus epidermidis</i> Clinical Isolates Recovered from Hospitals in Ohio. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 2923-2928.	1.4	40
359	Pharmacokinetics-Pharmacodynamics of Tazobactam in Combination with Piperacillin in an In Vitro Infection Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 2075-2080.	1.4	40
360	Evaluation of Vitek and API 20S for species identification of enterococci. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 22, 315-319.	0.8	39

#	ARTICLE	IF	CITATIONS
361	Antimicrobial Activity of SCH 27899, Oligosaccharide Member of the Everninomycin Class with a Wide Gram-Positive Spectrum. <i>Clinical Microbiology and Infection</i> , 1995, 1, 35-43.	2.8	39
362	Stability of linezolid activity in an era of mobile oxazolidinone resistance determinants: results from the 2009 Zyvox® Annual Appraisal of Potency and Spectrum program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 68, 459-467.	0.8	39
363	Oritavancin Microbiologic Features and Activity Results From the Surveillance Program in the United States. <i>Clinical Infectious Diseases</i> , 2012, 54, S203-S213.	2.9	39
364	Expansion of Clonal Complex 258 KPC-2-Producing <i>Klebsiella pneumoniae</i> in Latin American Hospitals: Report of the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 1668-1669.	1.4	39
365	Relationship between Ceftolozane-Tazobactam Exposure and Selection for <i>Pseudomonas aeruginosa</i> Resistance in a Hollow-Fiber Infection Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 6024-6031.	1.4	39
366	Cefmenoxime (SCE-1365), a new Cephalosporin: In Vitro Activity, Comparison with Other Antimicrobial Agents, Beta-Lactamase Stability, and Disk Diffusion Testing with Tentative Interpretive Criteria. <i>Antimicrobial Agents and Chemotherapy</i> , 1981, 20, 747-759.	1.4	38
367	Antimicrobial susceptibility testing (AST): A review of changing trends, quality control guidelines, test accuracy, and recommendation for the testing of β -lactam drugs. <i>Diagnostic Microbiology and Infectious Disease</i> , 1983, 1, 1-24.	0.8	38
368	Cefquinome (HR 111V) in vitro evaluation of a broad-spectrum cephalosporin indicated for infections in animals. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 20, 49-55.	0.8	38
369	Phenotypic detection of mec A-positive staphylococcal blood stream isolates: High accuracy of simple disk diffusion tests. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 25, 107-112.	0.8	38
370	Antimicrobial activity and in vitro susceptibility test development for cefditoren against <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i> , and <i>Streptococcus</i> species. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 37, 99-105.	0.8	38
371	In Vitro Antimicrobial Findings for Fusidic Acid Tested Against Contemporary (2008-2009) Gram-Positive Organisms Collected in the United States. <i>Clinical Infectious Diseases</i> , 2011, 52, S477-S486.	2.9	38
372	Polymyxins: Wisdom Does Not Always Come With Age. <i>Clinical Infectious Diseases</i> , 2013, 57, 877-883.	2.9	38
373	Relationship between Ceftolozane-Tazobactam Exposure and Drug Resistance Amplification in a Hollow-Fiber Infection Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 4134-4138.	1.4	38
374	Studies to optimize the in vitro testing of piperacillin combined with tazobactam (YTR 830). <i>Diagnostic Microbiology and Infectious Disease</i> , 1989, 12, 495-510.	0.8	37
375	In vitro activity evaluations of cefdinir (FK482, CI-983, and PD134393). <i>Diagnostic Microbiology and Infectious Disease</i> , 1991, 14, 425-434.	0.8	37
376	International multicenter investigation of LB203W, a new fluoronaphthyridone. <i>Clinical Microbiology and Infection</i> , 1998, 4, 280-284.	2.8	37
377	Antimicrobial activity of gatifloxacin compared to seven other compounds tested against Gram-positive organisms isolated at 10 cancer-treatment centers. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 37-43.	0.8	37
378	Carbapenem-resistant <i>Serratia marcescens</i> isolates producing Bush group 2f β -lactamase (SME-1) in the United States: results from the MYSTIC Programme. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 39, 125-127.	0.8	37

#	ARTICLE	IF	CITATIONS
379	Antimicrobial susceptibility patterns of unusual nonfermentative gram-negative bacilli isolated from Latin America: report from the SENTRY Antimicrobial Surveillance Program (1997-2002). <i>Memorias Do Instituto Oswaldo Cruz</i> , 2005, 100, 571-577.	0.8	37
380	TR-700 in vitro activity against and resistance mutation frequencies among Gram-positive pathogens. <i>Journal of Antimicrobial Chemotherapy</i> , 2009, 63, 716-720.	1.3	37
381	CEM-101 Activity against Gram-Positive Organisms. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 2182-2187.	1.4	37
382	Antimicrobial Activity of Ceftaroline-Avibactam Tested against Clinical Isolates Collected from U.S. Medical Centers in 2010-2011. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 1982-1988.	1.4	37
383	Potency and Spectrum of Activity of AN3365, a Novel Boron-Containing Protein Synthesis Inhibitor, Tested against Clinical Isolates of Enterobacteriaceae and Nonfermentative Gram-Negative Bacilli. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 2849-2857.	1.4	37
384	Antimicrobial activity, spectrum and pharmacokinetics of old and new orally administered cepheems. <i>Antimicrobial Newsletter</i> , 1988, 5, 1-7.	0.9	36
385	A review of cephalosporin metabolism: A lesson to be learned for future chemotherapy. <i>Diagnostic Microbiology and Infectious Disease</i> , 1989, 12, 25-31.	0.8	36
386	Antimicrobial activity of gatifloxacin tested against 1676 strains of ciprofloxacin-resistant Gram-positive cocci isolated from patient infections in North and South America. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 32, 247-252.	0.8	36
387	Cefepime MIC as a Predictor of the Extended-Spectrum β -Lactamase Type in <i>Klebsiella pneumoniae</i> , Taiwan. <i>Emerging Infectious Diseases</i> , 2002, 8, 522-524.	2.0	36
388	An overview of the Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) Program: 1997-2004. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 53, 247-256.	0.8	36
389	Activity of meropenem as serine carbapenemases evolve in US Medical Centers: monitoring report from the MYSTIC Program (2006). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 59, 425-432.	0.8	36
390	The Genetic Environment of the <i>cfr</i> Gene and the Presence of Other Mechanisms Account for the Very High Linezolid Resistance of <i>Staphylococcus epidermidis</i> Isolate 426-3147L. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 1173-1179.	1.4	36
391	In vitro efficacy of six cephalosporins tested against Enterobacteriaceae isolated at 38 North American medical centres participating in the SENTRY Antimicrobial Surveillance Program, 1997-1998. <i>International Journal of Antimicrobial Agents</i> , 2000, 15, 111-118.	1.1	35
392	In vitro activity of 11 antimicrobial agents, including gatifloxacin and GAR936, tested against clinical isolates of <i>Mycobacterium marinum</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 42, 145-147.	0.8	35
393	Potency and antimicrobial spectrum update for piperacillin/tazobactam (2000): emphasis on its activity against resistant organism populations and generally untested species causing community-acquired respiratory tract infections. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 49-60.	0.8	35
394	Geographic variations in garenoxacin (BMS284756) activity tested against pathogens associated with skin and soft tissue infections: report from the SENTRY Antimicrobial Surveillance Program (2000). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 303-309.	0.8	35
395	The Activity of Daptomycin Against Wild-Type <i>Staphylococcus aureus</i> and Strains with Reduced Susceptibility to Vancomycin. <i>Clinical Infectious Diseases</i> , 2006, 43, 798-799.	2.9	35
396	Comparison of 2002-2006 OPTAMA Programs for US Hospitals: Focus on Gram-Negative Resistance. <i>Annals of Pharmacotherapy</i> , 2009, 43, 220-227.	0.9	35

#	ARTICLE	IF	CITATIONS
397	Telavancin activity against Gram-positive bacteria isolated from respiratory tract specimens of patients with nosocomial pneumonia. <i>Journal of Antimicrobial Chemotherapy</i> , 2010, 65, 2396-2404.	1.3	35
398	Worldwide Appraisal and Update (2010) of Telavancin Activity Tested against a Collection of Gram-Positive Clinical Pathogens from Five Continents. <i>Antimicrobial Agents and Chemotherapy</i> , 2012, 56, 3999-4004.	1.4	35
399	Activity of cefepime (BMY-28142) and cefpirome (HR 810) against Gram-negative bacilli resistant to cefotaxime or ceftaxidime. <i>Journal of Antimicrobial Chemotherapy</i> , 1989, 23, 163-165.	1.3	34
400	The impact of antimicrobial resistance: changing epidemiology of community-acquired respiratory-tract infections. <i>American Journal of Health-System Pharmacy</i> , 1999, 56, S4-S11.	0.5	34
401	Results from the Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) Programme: report of the 2001 data from 15 United States medical centres. <i>International Journal of Antimicrobial Agents</i> , 2004, 23, 52-59.	1.1	34
402	Comprehensive in vitro evaluation of cefepime combined with aztreonam or ampicillin/sulbactam against multi-drug resistant <i>Pseudomonas aeruginosa</i> and <i>Acinetobacter</i> spp.. <i>International Journal of Antimicrobial Agents</i> , 2005, 25, 380-384.	1.1	34
403	Characterization of Methicillin-Resistant <i>Staphylococcus aureus</i> Strains Recovered from a Phase IV Clinical Trial for Linezolid versus Vancomycin for Treatment of Nosocomial Pneumonia. <i>Journal of Clinical Microbiology</i> , 2012, 50, 3694-3702.	1.8	34
404	Frequency of occurrence and antimicrobial susceptibility of Gram-negative bacteremia isolates in patients with urinary tract infection: results from United States and European hospitals (2009-2011). <i>Journal of Chemotherapy</i> , 2014, 26, 133-138.	0.7	34
405	In vitro antifungal susceptibilities of isolates of <i>Candida</i> spp. and <i>Aspergillus</i> spp. from China to nine systemically active antifungal agents: data from the SENTRY antifungal surveillance program, 2010 through 2012. <i>Mycoses</i> , 2015, 58, 209-214.	1.8	34
406	Surrogate analysis of vancomycin to predict susceptible categorization of dalbavancin. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 82, 73-77.	0.8	34
407	Multicenter in vitro evaluation of lomefloxacin (NY-198, SC-47111), including tests against nearly 7,000 bacterial isolates and preliminary recommendations for susceptibility testing. <i>Diagnostic Microbiology and Infectious Disease</i> , 1988, 10, 221-240.	0.8	33
408	Vancomycin-resistant <i>Enterococcus raffinosus</i> : Molecular epidemiology, species identification error, and frequency of occurrence in a national resistance surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 43-49.	0.8	33
409	Inducible β -lactamase-mediated resistance to third-generation cephalosporins. <i>Clinical Microbiology and Infection</i> , 1997, 3, S7-S20.	2.8	33
410	Evaluation of the in vitro activity of six broad-spectrum β -lactam antimicrobial agents tested against over 2,000 clinical isolates from 22 medical centers in Japan. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 123-134.	0.8	33
411	Antimicrobial susceptibility profile among β -haemolytic <i>Streptococcus</i> spp. collected in the SENTRY antimicrobial surveillance program—North America, 2001. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 46, 291-294.	0.8	33
412	Contemporary in vitro synergy rates for aztreonam combined with newer fluoroquinolones and β -lactams tested against gram-negative bacilli. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 47, 547-550.	0.8	33
413	Evaluation of PPI-0903M (T91825), a novel cephalosporin: bactericidal activity, effects of modifying in vitro testing parameters and optimization of disc diffusion tests. <i>Journal of Antimicrobial Chemotherapy</i> , 2005, 56, 1047-1052.	1.3	33
414	Evaluation of dalbavancin in combination with nine antimicrobial agents to detect enhanced or antagonistic interactions. <i>International Journal of Antimicrobial Agents</i> , 2006, 27, 557-560.	1.1	33

#	ARTICLE	IF	CITATIONS
415	Antimicrobial activity of cefepime tested against ceftazidime-resistant Gram-negative clinical strains from North American Hospitals: report from the SENTRY Antimicrobial Surveillance Program (1998â€“2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 63-68.	0.8	33
416	Influence of patient age on the frequency of occurrence and antimicrobial resistance patterns of isolates from hematology/oncology patients: Report from the Chemotherapy Alliance for Neutropenics and the Control of Emerging Resistance Program (North America). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 75-82.	0.8	33
417	Activity of Dalbavancin Tested against <i>Staphylococcus</i> spp. and β -Hemolytic <i>Streptococcus</i> spp. Isolated from 52 Geographically Diverse Medical Centers in the United States. <i>Journal of Clinical Microbiology</i> , 2007, 45, 998-1004.	1.8	33
418	The in vitro evaluation of solithromycin (CEM-101) against pathogens isolated in the United States and Europe (2009). <i>Journal of Infection</i> , 2010, 61, 476-483.	1.7	33
419	Differences in potency and categorical agreement between colistin and polymyxin B when testing 15,377 clinical strains collected worldwide. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 83, 379-381.	0.8	33
420	Multicenter evaluation of antimicrobial resistance to six broad-spectrum β -lactams in Colombia using the estest method. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 265-272.	0.8	32
421	Comparative antimicrobial activity of trovafloxacin tested against 3049 streptococcus pneumoniae isolates from the 1997â€“1998 respiratory infection season. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 32, 119-126.	0.8	32
422	In vitro evaluation of cefepime and other broad-spectrum β -lactams in 22 medical centers in Japan: a phase II trial comparing two annual organism samples. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 307-315.	0.8	32
423	Detection of emerging resistance patterns within longitudinal surveillance systems: data sensitivity and microbial susceptibility. <i>Journal of Antimicrobial Chemotherapy</i> , 2000, 46, 1-8.	1.3	32
424	Determination of epidemic clonality among multidrug-resistant strains of <i>Acinetobacter</i> spp. and <i>Pseudomonas aeruginosa</i> in the MYSTIC Programme (USA, 1999â€“2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 211-216.	0.8	32
425	Daptomycin tested against 915 bloodstream isolates of viridans group streptococci (eight species) and <i>Streptococcus bovis</i> . <i>Journal of Antimicrobial Chemotherapy</i> , 2005, 55, 574-578.	1.3	32
426	Potency and spectrum of garenoxacin tested against an international collection of skin and soft tissue infection pathogens: report from the SENTRY antimicrobial surveillance program (1999â€“2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 58, 19-26.	0.8	32
427	Activities of Tedizolid and Linezolid Determined by the Reference Broth Microdilution Method against 3,032 Gram-Positive Bacterial Isolates Collected in Asia-Pacific, Eastern Europe, and Latin American Countries in 2014. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 5393-5399.	1.4	32
428	Antimicrobial susceptibility patterns of community- and hospital-acquired methicillin-resistant <i>Staphylococcus aureus</i> from United States Hospitals: results from the AWARE Ceftaroline Surveillance Program (2012â€“2014). <i>Diagnostic Microbiology and Infectious Disease</i> , 2016, 86, 76-79.	0.8	32
429	Polymyxin Susceptibility Testing and Interpretive Breakpoints: Recommendations from the United States Committee on Antimicrobial Susceptibility Testing (USCAST). <i>Antimicrobial Agents and Chemotherapy</i> , 2020, 64, .	1.4	32
430	In vitro evaluation of cefixime (FK027, FR17027, CL284635): Spectrum against recent clinical isolates, comparative antimicrobial activity, β -lactamase stability, and preliminary susceptibility testing criteria. <i>Diagnostic Microbiology and Infectious Disease</i> , 1986, 5, 151-162.	0.8	31
431	Comparative in vitro assessment of sparfloxacin activity and spectrum using results from over 14,000 pathogens isolated at 190 medical centers in the USA. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 173-186.	0.8	31
432	Fluoroquinolone-resistant <i>Moraxella catarrhalis</i> in a patient with pneumonia: report from the SENTRY antimicrobial surveillance program (1998). <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 32, 131-135.	0.8	31

#	ARTICLE	IF	CITATIONS
433	A comparison of the antimicrobial activity of meropenem and selected broad-spectrum antimicrobials tested against multi-drug resistant Gram-negative bacilli including bacteraemic <i>Salmonella</i> spp.: International Journal of Antimicrobial Agents, 2002, 20, 426-431.	1.1	31
434	Influence of patient age on the susceptibility patterns of <i>Streptococcus pneumoniae</i> isolates in North America (2000-2001): report from the SENTRY Antimicrobial Surveillance Program. Diagnostic Microbiology and Infectious Disease, 2003, 46, 77-80.	0.8	31
435	CANCER Resistance Surveillance Program: Initial Results from Hematology/Oncology Centers in North America. Annals of Pharmacotherapy, 2003, 37, 47-56.	0.9	31
436	Worldwide summary of telavancin spectrum and potency against Gram-positive pathogens: 2007 to 2008 surveillance results. Diagnostic Microbiology and Infectious Disease, 2010, 67, 359-368.	0.8	31
437	Tigecycline activity tested against multidrug-resistant Enterobacteriaceae and <i>Acinetobacter</i> spp. isolated in US medical centers (2005-2009). Diagnostic Microbiology and Infectious Disease, 2011, 69, 223-227.	0.8	31
438	Oritavancin Activity against Vancomycin-Susceptible and Vancomycin-Resistant Enterococci with Molecularly Characterized Glycopeptide Resistance Genes Recovered from Bacteremic Patients, 2009-2010. Antimicrobial Agents and Chemotherapy, 2012, 56, 1639-1642.	1.4	31
439	Method Preferences and Test Accuracy of Antimicrobial Susceptibility Testing. Archives of Pathology and Laboratory Medicine, 2001, 125, 1285-1289.	1.2	31
440	National survey of the in vitro spectrum of piperacillin-tazobactam tested against more than 40,000 aerobic clinical isolates from 236 medical centers. Diagnostic Microbiology and Infectious Disease, 1995, 21, 141-151.	0.8	30
441	Spectrum and activity of three contemporary fluoroquinolones tested against <i>Pseudomonas aeruginosa</i> isolates from urinary tract infections in the SENTRY Antimicrobial Surveillance Program (Europe and the Americas; 2000): More alike than different!. Diagnostic Microbiology and Infectious Disease, 2001, 41, 161-163.	0.8	30
442	Contemporary antimicrobial activity of triple antibiotic ointment: a multiphased study of recent clinical isolates in the United States and Australia. Diagnostic Microbiology and Infectious Disease, 2006, 54, 63-71.	0.8	30
443	Delayed resistance selection for doripenem when passaging <i>Pseudomonas aeruginosa</i> isolates with doripenem plus an aminoglycoside. Diagnostic Microbiology and Infectious Disease, 2006, 55, 241-243.	0.8	30
444	Multicenter Evaluation of the Etest and Disk Diffusion Methods for Differentiating Daptomycin-Susceptible from Non-Daptomycin-Susceptible <i>Staphylococcus aureus</i> Isolates. Journal of Clinical Microbiology, 2006, 44, 3098-3104.	1.8	30
445	In vitro potency evaluations of various piperacillin/tazobactam generic products compared with the contemporary branded (Zosyn®, Wyeth) formulation. Diagnostic Microbiology and Infectious Disease, 2008, 61, 76-79.	0.8	30
446	Pharmacokinetic and pharmacodynamic profile of ceftobiprole. Diagnostic Microbiology and Infectious Disease, 2008, 61, 96-102.	0.8	30
447	Activity Analyses of Staphylococcal Isolates From Pediatric, Adult, and Elderly Patients: AWARE Ceftaroline Surveillance Program. Clinical Infectious Diseases, 2012, 55, S181-S186.	2.9	30
448	Spectrum and potency of ceftaroline tested against leading pathogens causing skin and soft-tissue infections in Europe (2010). International Journal of Antimicrobial Agents, 2013, 41, 337-342.	1.1	30
449	Oritavancin Activity against <i>Staphylococcus aureus</i> Causing Invasive Infections in U.S. and European Hospitals: a 5-Year International Surveillance Program. Antimicrobial Agents and Chemotherapy, 2014, 58, 2921-2924.	1.4	30
450	Activity of Debio1452, a FabI Inhibitor with Potent Activity against <i>Staphylococcus aureus</i> and Coagulase-Negative <i>Staphylococcus</i> spp., Including Multidrug-Resistant Strains. Antimicrobial Agents and Chemotherapy, 2015, 59, 2583-2587.	1.4	30

#	ARTICLE	IF	CITATIONS
451	<i>In Vitro</i> Activity of Ceftazidime-Avibactam against Contemporary <i>Pseudomonas aeruginosa</i> Isolates from U.S. Medical Centers by Census Region, 2014. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 2537-2541.	1.4	30
452	In vitro activity of ceftazidime/avibactam against Gram-negative pathogens isolated from pneumonia in hospitalised patients, including ventilated patients. <i>International Journal of Antimicrobial Agents</i> , 2016, 47, 235-242.	1.1	30
453	Comparative antimicrobial activity of piperacillin-tazobactam tested against more than 5000 recent clinical isolates from five medical centers a reevaluation after five years. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 153-168.	0.8	29
454	Multicenter Evaluation of the Antimicrobial Activity for Six Broad-Spectrum β -Lactams in Venezuela Using the Etest Method. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 30, 45-52.	0.8	29
455	Comparison of Activities of Broad-Spectrum β -Lactam Compounds against 1,128 Gram-Positive Cocci Recently Isolated in Cancer Treatment Centers. <i>Antimicrobial Agents and Chemotherapy</i> , 1999, 43, 940-943.	1.4	29
456	Disk diffusion susceptibility test development for the new glycycline, GAR-936. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 249-252.	0.8	29
457	Initial Quality Control Evaluations for Susceptibility Testing of Dalbavancin (BI397), an Investigational Glycopeptide with Potent Gram-Positive Activity. <i>Journal of Clinical Microbiology</i> , 2003, 41, 2795-2796.	1.8	29
458	Antimicrobial activity of tigecycline (GAR-936) tested against 3498 recent isolates of <i>Staphylococcus aureus</i> recovered from nosocomial and community-acquired infections. <i>International Journal of Antimicrobial Agents</i> , 2004, 24, 567-571.	1.1	29
459	Oxazolidinone susceptibility patterns for 2005: International Report from the Zyvox [®] Annual Appraisal of Potency and Spectrum Study. <i>International Journal of Antimicrobial Agents</i> , 2007, 29, 295-301.	1.1	29
460	Potency of Anidulafungin Compared to Nine Other Antifungal Agents Tested against <i>Candida</i> spp., <i>Cryptococcus</i> spp., and <i>Aspergillus</i> spp.: Results from the Global SENTRY Antimicrobial Surveillance Program (2008). <i>Journal of Clinical Microbiology</i> , 2010, 48, 2984-2987.	1.8	29
461	Evaluation of the activity of fusidic acid tested against contemporary Gram-positive clinical isolates from the USA and Canada. <i>International Journal of Antimicrobial Agents</i> , 2010, 35, 282-287.	1.1	29
462	Surveillance of dalbavancin potency and spectrum in the United States (2012). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 122-123.	0.8	29
463	Pharmacodynamic considerations in the treatment of moderate to severe pseudomonal infections with cefepime. <i>Journal of Antimicrobial Chemotherapy</i> , 2002, 49, 445-453.	1.3	28
464	Antimicrobial spectrum and activity of NVP PDF-713, a novel peptide deformylase inhibitor, tested against 1,837 recent gram-positive clinical isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 63-65.	0.8	28
465	The in vitro evaluation of tigecycline tested against pathogens isolated in eight countries in the Asia-Western Pacific region (2008). <i>Journal of Infection</i> , 2010, 60, 440-451.	1.7	28
466	Minocycline activity tested against <i>Acinetobacter baumannii</i> complex, <i>Stenotrophomonas maltophilia</i> , and <i>Burkholderia cepacia</i> species complex isolates from a global surveillance program (2013). <i>Diagnostic Microbiology and Infectious Disease</i> , 2016, 85, 352-355.	0.8	28
467	Results from the Solithromycin International Surveillance Program (2014). <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 3662-3668.	1.4	28
468	In vitro activity and disk susceptibility of timentin: current status. <i>American Journal of Medicine</i> , 1985, 79, 25-32.	0.6	27

#	ARTICLE	IF	CITATIONS
469	Comparative activity of garenoxacin (BMS 284756), a novel desfluoroquinolone, tested against 8,331 isolates from community-acquired respiratory tract infections: North American results from the SENTRY Antimicrobial Surveillance Program (1999–2001). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 45, 273-278.	0.8	27
470	Quinupristin-Dalfopristin Resistance in <i>Streptococcus pneumoniae</i> : Novel L22 Ribosomal Protein Mutation in Two Clinical Isolates from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 2696-2698.	1.4	27
471	Evaluation of the contemporary occurrence rates of metallo- β -lactamases in multidrug-resistant Gram-negative bacilli in Japan: Report from the SENTRY Antimicrobial Surveillance Program (1998–2002). <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 289-294.	0.8	27
472	<i>In Vitro</i> Activity of Telavancin against a Contemporary Worldwide Collection of <i>Staphylococcus aureus</i> Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 2704-2706.	1.4	27
473	<i>Escherichia coli</i> resistance to quinolones at a comprehensive cancer center. <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 67, 266-269.	0.8	27
474	Antimicrobial activity of ceftaroline and comparator agents tested against bacterial isolates causing skin and soft tissue infections and community-acquired respiratory tract infections isolated from the Asia-Pacific region and South Africa (2010). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 61-68.	0.8	27
475	Antimicrobial activity of the novel polymyxin derivative NAB739 tested against Gram-negative pathogens. <i>Journal of Antimicrobial Chemotherapy</i> , 2013, 68, 636-639.	1.3	27
476	Ceftaroline Activity against Bacterial Pathogens Frequently Isolated in U.S. Medical Centers: Results from Five Years of the AWARE Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 2458-2461.	1.4	27
477	Interpretive errors using an automated system for the susceptibility testing of imipenem and aztreonam. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 57-60.	0.8	26
478	Antimicrobial Susceptibility Testing of Clinical Isolates of <i>Bordetella pertussis</i> from Northern California: Report from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2001, 45, 3599-3600.	1.4	26
479	Confirmation of extended-spectrum β -lactamase-producing <i>Serratia marcescens</i> : preliminary report from Taiwan. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 45, 221-224.	0.8	26
480	Reevaluation of Enterobacteriaceae MIC/disk diffusion zone diameter regression scattergrams for 9 β -lactams: adjustments of breakpoints for strains producing extended spectrum β -lactamases. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 235-246.	0.8	26
481	Antimicrobial activity of daptomycin tested against clinical strains of indicated species isolated in North American medical centers (2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 53, 329-332.	0.8	26
482	Comparisons of parenteral broad-spectrum cephalosporins tested against bacterial isolates from pediatric patients: report from the SENTRY Antimicrobial Surveillance Program (1998–2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 109-116.	0.8	26
483	Update of contemporary antimicrobial resistance rates across China: reference testing results for 12 medical centers (2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 77, 258-266.	0.8	26
484	Arbekacin Activity against Contemporary Clinical Bacteria Isolated from Patients Hospitalized with Pneumonia. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 3263-3270.	1.4	26
485	Antimicrobial Activities of Ceftazidime-Avibactam and Comparator Agents against Gram-Negative Organisms Isolated from Patients with Urinary Tract Infections in U.S. Medical Centers, 2012 to 2014. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 4355-4360.	1.4	26
486	The antimicrobial activity of A-56268 (TE-031) and roxithromycin (RU965) against <i>Legionella</i> using broth microdilution method. <i>Journal of Antimicrobial Chemotherapy</i> , 1987, 19, 841-842.	1.3	25

#	ARTICLE	IF	CITATIONS
487	Comparative antimicrobial activity of gatifloxacin tested against Streptococcus spp. including quality control guidelines and etest method validation. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 91-98.	0.8	25
488	Antimicrobial resistance rates and clonality results from the Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) Programme: Report of year five (2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 273-281.	0.8	25
489	In vitro activity of tigecycline, a new glycolcycline, tested against 1,326 clinical bacterial strains isolated from Latin America. <i>Brazilian Journal of Infectious Diseases</i> , 2005, 9, 348-356.	0.3	25
490	Comparison of Dalbavancin MIC Values Determined by Etest (AB BIODISK) and Reference Dilution Methods Using Gram-Positive Organisms. <i>Journal of Clinical Microbiology</i> , 2006, 44, 2988-2990.	1.8	25
491	Antimicrobial activity of a novel peptide deformylase inhibitor, LBM415, tested against respiratory tract and cutaneous infection pathogens: a global surveillance report (2003-2004). <i>Journal of Antimicrobial Chemotherapy</i> , 2006, 57, 914-923.	1.3	25
492	Ceftazidime-avibactam activity when tested against ceftazidime-nonsusceptible <i>Citrobacter</i> spp., <i>Enterobacter</i> spp., <i>Serratia marcescens</i> , and <i>Pseudomonas aeruginosa</i> from United States medical centers (2011-2014). <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 83, 389-394.	0.8	25
493	In vitro activity of linezolid as assessed through the 2013 LEADER surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 81, 283-289.	0.8	25
494	Oxacillin- and Quinolone-Resistant <i>Staphylococcus aureus</i> in Sao Paulo, Brazil: A Multicenter Molecular Epidemiology Study. <i>Infection Control and Hospital Epidemiology</i> , 1993, 14, 260-264.	1.0	24
495	Emerging multiply resistant enterococci among clinical isolates II. Validation of the Etest to recognize glycopeptide-resistant strains. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 95-100.	0.8	24
496	Activity of BMS284756 against 2,681 recent clinical isolates of <i>Haemophilus influenzae</i> and <i>Moraxella catarrhalis</i> : report from The SENTRY Antimicrobial Surveillance Program (2000) in Europe, Canada and the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 39, 245-250.	0.8	24
497	Cefditoren in vitro activity and spectrum: a review of international studies using reference methods. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 41, 1-14.	0.8	24
498	Contemporary evaluation of the in vitro activity and spectrum of cefdinir compared with other orally administered antimicrobials tested against common respiratory tract pathogens (2000-2002). <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 47, 515-525.	0.8	24
499	Potential utility of a peptide deformylase inhibitor (NVP PDF-713) against oxazolidinone-resistant or streptogramin-resistant Gram-positive organism isolates. <i>Journal of Antimicrobial Chemotherapy</i> , 2004, 53, 804-807.	1.3	24
500	Susceptibility patterns of <i>Streptococcus pneumoniae</i> isolates in North America (2002-2003): contemporary in vitro activities of amoxicillin/clavulanate and 15 other antimicrobial agents. <i>International Journal of Antimicrobial Agents</i> , 2005, 25, 282-289.	1.1	24
501	Oral β -lactams applied to uncomplicated infections of skin and skin structures. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, S55-S65.	0.8	24
502	Antimicrobial Activities of Piperacillin-Tazobactam against <i>Haemophilus influenzae</i> Isolates, Including β -Lactamase-Negative Ampicillin-Resistant and β -Lactamase-Positive Amoxicillin-Clavulanate-Resistant Isolates, and Mutations in Their Quinolone Resistance-Determining Regions. <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 4225-4230.	1.4	24
503	Telavancin In Vitro Activity against a Collection of Methicillin-Resistant <i>Staphylococcus aureus</i> Isolates, Including Resistant Subsets, from the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 1811-1814.	1.4	24
504	β -Lactamase Characterization of Gram-Negative Pathogens Recovered from Patients Enrolled in the Phase 2 Trials for Ceftazidime-Avibactam: Clinical Efficacies Analyzed against Subsets of Molecularly Characterized Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 1328-1335.	1.4	24

#	ARTICLE	IF	CITATIONS
505	Antimicrobial Activity of High-Proportion Cefepime-Tazobactam (WCK 4282) against a Large Number of Gram-Negative Isolates Collected Worldwide in 2014. <i>Antimicrobial Agents and Chemotherapy</i> , 2017, 61, .	1.4	24
506	Clarithromycin in vitro activity enhanced by its major metabolite, 14-hydroxyclearithromycin. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 259-266.	0.8	23
507	Species identification and determination of high-level aminoglycoside resistance among enterococci comparison study of sterile body fluid isolates, 1985-1991. <i>Diagnostic Microbiology and Infectious Disease</i> , 1993, 16, 119-122.	0.8	23
508	North American (United States and Canada) comparative susceptibility of two fluoroquinolones: ofloxacin and ciprofloxacin. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 18, 49-56.	0.8	23
509	Development, characterization, and initial evaluations of S1 a new chromogenic cephalosporin for β -lactamase detection. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 1-8.	0.8	23
510	Evaluation of the Vitek System to accurately test the susceptibility of <i>Pseudomonas aeruginosa</i> clinical isolates against cefepime. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 32, 107-110.	0.8	23
511	Tiamulin Activity against Fastidious and Nonfastidious Veterinary and Human Bacterial Isolates: Initial Development of In Vitro Susceptibility Test Methods. <i>Journal of Clinical Microbiology</i> , 2002, 40, 461-465.	1.8	23
512	Microbiology of newer fluoroquinolones: focus on respiratory pathogens. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 213-220.	0.8	23
513	Multicentre evaluation of the in vitro activity of linezolid in the Western Pacific. <i>Journal of Antimicrobial Chemotherapy</i> , 2003, 51, 339-345.	1.3	23
514	Validation of commercial dry-form broth microdilution panels and test reproducibility for susceptibility testing of dalbavancin, a new very long-acting glycopeptide. <i>International Journal of Antimicrobial Agents</i> , 2004, 23, 197-199.	1.1	23
515	Activity of Linezolid against 3,251 Strains of Uncommonly Isolated Gram-Positive Organisms: Report from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2007, 51, 1491-1493.	1.4	23
516	Tigecycline (GAR-936) activity against <i>Streptococcus gallolyticus</i> (bovis) and viridans group streptococci. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 333-336.	0.8	23
517	Complete Sequence of p07-406, a 24,179-Base-Pair Plasmid Harboring the <i>bla</i> _{VIM-7} Metallo- β -Lactamase Gene in a <i>Pseudomonas aeruginosa</i> Isolate from the United States. <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 3099-3105.	1.4	23
518	Multicenter evaluation of the in vitro activity of dalbavancin tested against staphylococci and streptococci in 5 European countries: results from the DECIDE Surveillance Program (2007). <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 64, 177-184.	0.8	23
519	Molecular characterization of vancomycin-resistant <i>Enterococcus</i> spp. clinical isolates recovered from hospitalized patients among several medical institutions in China. <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 74, 399-403.	0.8	23
520	Ceftaroline activity against bacterial organisms isolated from acute bacterial skin and skin structure infections in United States medical centers (2009-2011). <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 78, 422-428.	0.8	23
521	Mechanisms of Resistance, Clonal Expansion, and Increasing Prevalence of <i>Acinetobacter baumannii</i> Strains Displaying Elevated Tigecycline MIC Values in Latin America. <i>Microbial Drug Resistance</i> , 2016, 22, 253-258.	0.9	23
522	Historical overview of the cephalosporin spectrum: Four generations of structural evolution. <i>Antimicrobial Newsletter</i> , 1992, 8, 75-82.	0.9	22

#	ARTICLE	IF	CITATIONS
523	Epidemiology, Laboratory Detection, and Therapy of Penicillin-resistant Streptococcal Infections. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 453-459.	0.8	22
524	Comparative activity of clinafloxacin and nine other compounds tested against 2000 contemporary clinical isolates from patients in United States hospitals. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 81-88.	0.8	22
525	Can antimicrobial susceptibility testing results for ciprofloxacin or levofloxacin predict susceptibility to a newer fluoroquinolone, gatifloxacin?: Report from The SENTRY Antimicrobial Surveillance Program (1997-99). <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 39, 237-243.	0.8	22
526	Activities of BMS 284756 (T-3811) against <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i> , and <i>Streptococcus pneumoniae</i> Isolates from SENTRY Antimicrobial Surveillance Program Medical Centers in Latin America (1999). <i>Antimicrobial Agents and Chemotherapy</i> , 2001, 45, 1463-1466.	1.4	22
527	Comparative activity of oral and parenteral cephalosporins tested against multidrug-resistant <i>Streptococcus pneumoniae</i> : report from the SENTRY Antimicrobial Surveillance Program (1997-2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 51, 147-150.	0.8	22
528	Carbapenem Susceptibility Discords among <i>Acinetobacter</i> Isolates. <i>Clinical Infectious Diseases</i> , 2006, 42, 158-158.	2.9	22
529	A global evaluation of voriconazole activity tested against recent clinical isolates of <i>Candida</i> spp.. <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 63, 233-236.	0.8	22
530	Update on the telavancin activity tested against European staphylococcal clinical isolates (2009-2010). <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 71, 93-97.	0.8	22
531	Challenges in Assessing Microbial Susceptibility and Predicting Clinical Response to Newer-Generation Fluoroquinolones. <i>Journal of Ocular Pharmacology and Therapeutics</i> , 2012, 28, 3-11.	0.6	22
532	Antimicrobial activity of daptomycin in comparison to glycopeptides and other antimicrobials when tested against numerous species of coagulase-negative <i>Staphylococcus</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 73, 212-214.	0.8	22
533	Activity of ceftobiprole against methicillin-resistant <i>Staphylococcus aureus</i> strains with reduced susceptibility to daptomycin, linezolid or vancomycin, and strains with defined SCCmec types. <i>International Journal of Antimicrobial Agents</i> , 2014, 43, 323-327.	1.1	22
534	Activity of Fusidic Acid Tested against <i>Staphylococci</i> Isolated from Patients in U.S. Medical Centers in 2014. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 3827-3831.	1.4	22
535	Ofloxacin, a new broad-spectrum fluoroquinolone results from a multicenter, national comparative activity surveillance study. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 425-434.	0.8	21
536	The north american component (the united states and canada) of an international comparative MIC trial monitoring ofloxacin resistance. <i>Diagnostic Microbiology and Infectious Disease</i> , 1993, 17, 157-161.	0.8	21
537	Pulsed-field gel electrophoresis of restriction-digested genomic DNA and antimicrobial susceptibility of <i>Xanthomonas maltophilia</i> strains from Brazil, Switzerland and the USA. <i>Journal of Antimicrobial Chemotherapy</i> , 1994, 33, 615-618.	1.3	21
538	Multicenter evaluation of antimicrobial resistance to six broad-spectrum β -lactams in Colombia: comparison of data from 1997 and 1998 using the Etest method. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 235-241.	0.8	21
539	<i>Salmonella</i> bloodstream infections: report from the SENTRY Antimicrobial Surveillance Program (1997-2001). <i>International Journal of Antimicrobial Agents</i> , 2003, 22, 395-405.	1.1	21
540	Contemporary Prevalence of BRO β -Lactamases in <i>Moraxella catarrhalis</i> : Report from the SENTRY Antimicrobial Surveillance Program (North America, 1997 to 2004). <i>Journal of Clinical Microbiology</i> , 2006, 44, 3775-3777.	1.8	21

#	ARTICLE	IF	CITATIONS
541	In vitro activity of omiganan pentahydrochloride tested against vancomycin-tolerant, -intermediate, and -resistant <i>Staphylococcus aureus</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 60, 399-403.	0.8	21
542	JNJ-Q2, a New Fluoroquinolone with Potent In Vitro Activity against <i>Staphylococcus aureus</i> , Including Methicillin- and Fluoroquinolone-Resistant Strains. <i>Antimicrobial Agents and Chemotherapy</i> , 2011, 55, 3631-3634.	1.4	21
543	Telavancin activity tested against a contemporary collection of Gram-positive pathogens from USA Hospitals (2007-2009). <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 72, 113-117.	0.8	21
544	Update on the prevalence and genetic characterization of NDM-1-producing Enterobacteriaceae in Indian hospitals during 2010. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 210-213.	0.8	21
545	Antimicrobial Activity of Ceftaroline Tested against Drug-Resistant Subsets of <i>Streptococcus pneumoniae</i> from U.S. Medical Centers. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 2468-2471.	1.4	21
546	Ceftaroline: clinical and microbiology experience with focus on methicillin-resistant <i>Staphylococcus aureus</i> after regulatory approval in the USA. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 81, 183-188.	0.8	21
547	Baseline Activity of Telavancin against Gram-Positive Clinical Isolates Responsible for Documented Infections in U.S. Hospitals (2011-2012) as Determined by the Revised Susceptibility Testing Method. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 702-706.	1.4	21
548	Cefdinir (FK482), an orally administered cephalosporin in vitro activity comparison against recent clinical isolates from five medical centers and determination of MIC quality control guidelines. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 537-543.	0.8	20
549	Can antimicrobial activity be sustained? an appraisal of orally administered drugs used for respiratory tract infections. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 27, 21-28.	0.8	20
550	Antimicrobial susceptibility of bacteria causing urinary tract infections in Latin American hospitals: results from the SENTRY Antimicrobial Surveillance Program (1997). <i>Clinical Microbiology and Infection</i> , 1999, 5, 478-487.	2.8	20
551	Multicenter evaluation of the antimicrobial activity for seven broad-spectrum β -lactams in Turkey using the Etest method. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 65-73.	0.8	20
552	Molecular Typing and Antimicrobial Susceptibility of Vancomycin-Resistant <i>Enterococcus faecium</i> in Brazil. <i>Infection Control and Hospital Epidemiology</i> , 2002, 23, 19-22.	1.0	20
553	Comparative spectrum and activity of NVP-PDF386 (VRC4887), a new peptide deformylase inhibitor. <i>Journal of Antimicrobial Chemotherapy</i> , 2003, 51, 157-161.	1.3	20
554	Clonal occurrences of multidrug-resistant Gram-negative bacilli: report from the Meropenem Yearly Susceptibility Test Information Collection Surveillance Program in the United States (2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 54, 249-257.	0.8	20
555	Molecular Characterization of <i>Staphylococcus aureus</i> Isolates from a 2005 Clinical Trial of Uncomplicated Skin and Skin Structure Infections. <i>Antimicrobial Agents and Chemotherapy</i> , 2007, 51, 3381-3384.	1.4	20
556	Increased Antimicrobial Susceptibility Profiles among Polymyxin-Resistant <i>Acinetobacter baumannii</i> Clinical Isolates. <i>Clinical Infectious Diseases</i> , 2008, 46, 1324-1326.	2.9	20
557	Spectrum of activity, mutation rates, synergistic interactions, and the effects of pH and serum proteins for fusidic acid (CEM-102). <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 66, 301-307.	0.8	20
558	Effects of Breakpoint Changes on Carbapenem Susceptibility Rates of Enterobacteriaceae: Results from the SENTRY Antimicrobial Surveillance Program, United States, 2008 to 2012. <i>Canadian Journal of Infectious Diseases and Medical Microbiology</i> , 2014, 25, 285-287.	0.7	20

#	ARTICLE	IF	CITATIONS
559	Ceftaroline Activity Tested Against Bacterial Isolates From Pediatric Patients. <i>Pediatric Infectious Disease Journal</i> , 2014, 33, 837-842.	1.1	20
560	Update on dalbavancin activity tested against Gram-positive clinical isolates responsible for documented skin and skin-structure infections in US and European hospitals (2011-13): Table 1.. <i>Journal of Antimicrobial Chemotherapy</i> , 2016, 71, 276-278.	1.3	20
561	In vitro activity of cefepime and ceftazidime against 197 nosocomial blood stream isolates of streptococci: A multicenter sample. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 273-276.	0.8	19
562	Evaluation of Low-Dose, Extended-Interval Clindamycin Regimens against <i>Staphylococcus aureus</i> and <i>Streptococcus pneumoniae</i> Using a Dynamic In Vitro Model of Infection. <i>Antimicrobial Agents and Chemotherapy</i> , 1999, 43, 2005-2009.	1.4	19
563	Antimicrobial activity and spectrum of SCH27899 (ziracin®) tested against gram-positive species including recommendations for routine susceptibility testing methods and quality control. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 103-110.	0.8	19
564	Multicenter evaluation of the antimicrobial activity for six broad-spectrum β -lactams in Venezuela: comparison of data from 1997 and 1998 using the Etest method. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 153-158.	0.8	19
565	Cefditoren activity against nearly 1000 non-fastidious bacterial isolates and the development of in vitro susceptibility test methods. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 37, 143-146.	0.8	19
566	Gatifloxacin used for therapy of outpatient community-acquired pneumonia caused by <i>Streptococcus pneumoniae</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 93-100.	0.8	19
567	Emergence and epidemiology of fluoroquinolone-resistant <i>Streptococcus pneumoniae</i> strains from Italy: report from the SENTRY Antimicrobial Surveillance Program (2001-2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 54, 157-164.	0.8	19
568	Activity of garenoxacin, an investigational des-F(6)-quinolone, tested against pathogens from community-acquired respiratory tract infections, including those with elevated or resistant-level fluoroquinolone MIC values. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 58, 9-17.	0.8	19
569	Expanded studies of piperacillin/tazobactam formulations: variations among branded product lots and assessment of 46 generic lots. <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 65, 319-322.	0.8	19
570	Antimicrobial activity of daptomycin tested against <i>Staphylococcus aureus</i> with vancomycin MIC of 2 μ g/mL isolated in the United States and European hospitals (2006-2008). <i>Diagnostic Microbiology and Infectious Disease</i> , 2010, 66, 329-331.	0.8	19
571	Contemporary tetracycline susceptibility testing: doxycycline MIC methods and interpretive criteria (CLSI and EUCAST) performance when testing Gram-positive pathogens. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 69-72.	0.8	19
572	Quality Control MIC Ranges Used for Telavancin with Application of a Revised CLSI Reference Broth Microdilution Method. <i>Journal of Clinical Microbiology</i> , 2014, 52, 3399-3401.	1.8	19
573	Decreased Ceftriaxone Susceptibility in Emerging (35B and 6C) and Persisting (19A) <i>Streptococcus pneumoniae</i> Serotypes in the United States, 2011-2012: Ceftaroline Remains Active <i>In Vitro</i> among β -Lactam Agents. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 4923-4927.	1.4	19
574	Antimicrobial activity of ceftaroline combined with avibactam tested against bacterial organisms isolated from acute bacterial skin and skin structure infections in United States medical centers (2010-2012). <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 78, 449-456.	0.8	19
575	Ceftaroline Activity Tested Against Bacterial Isolates Causing Community-acquired Respiratory Tract Infections and Skin and Skin Structure Infections in Pediatric Patients From United States Hospitals. <i>Pediatric Infectious Disease Journal</i> , 2017, 36, 486-491.	1.1	19
576	Assessment of Laboratory Performance With <i>Streptococcus pneumoniae</i> Antimicrobial Susceptibility Testing in the United States. <i>Archives of Pathology and Laboratory Medicine</i> , 1999, 123, 285-289.	1.2	19

#	ARTICLE	IF	CITATIONS
577	Antimicrobial Susceptibility Tests with Cefotaxime and Correlation with Clinical Bacteriologic Response. <i>Clinical Infectious Diseases</i> , 1982, 4, S316-S324.	2.9	18
578	Should novobiocin be clinically re-evaluated?. <i>Diagnostic Microbiology and Infectious Disease</i> , 1989, 12, 363-365.	0.8	18
579	Frequency of occurrence and antimicrobial susceptibility patterns for pathogens isolated from Latin American patients with a diagnosis of pneumonia: results from the SENTRY antimicrobial surveillance program (1998). <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 37, 63-74.	0.8	18
580	In vitro activity of selected cephalosporins and erythromycin against staphylococci and pneumococci isolated at 38 North American medical centers participating in the SENTRY Antimicrobial Surveillance Program, 1997-1998. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 37, 93-98.	0.8	18
581	Activity of BMS284756 (T-3811) tested against anaerobic bacteria, <i>Campylobacter jejuni</i> , <i>Helicobacter pylori</i> and <i>Legionella</i> spp.. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 40, 45-49.	0.8	18
582	Antimicrobial susceptibility of inducible AmpC β -lactamase-producing Enterobacteriaceae from the Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) Programme, Europe 1997-2000. <i>International Journal of Antimicrobial Agents</i> , 2002, 19, 383-388.	1.1	18
583	Selection of a Surrogate Agent (Vancomycin or Teicoplanin) for Initial Susceptibility Testing of Dalbavancin: Results from an International Antimicrobial Surveillance Program. <i>Journal of Clinical Microbiology</i> , 2006, 44, 2622-2625.	1.8	18
584	Plasmid-borne vga(A)-encoding gene in methicillin-resistant <i>Staphylococcus aureus</i> ST398 recovered from swine and a swine farmer in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 71, 177-180.	0.8	18
585	ZAAPS Program results for 2010: an activity and spectrum analysis of linezolid using clinical isolates from 75 medical centres in 24 countries. <i>Journal of Chemotherapy</i> , 2012, 24, 328-337.	0.7	18
586	Comprehensive update of dalbavancin activity when tested against uncommonly isolated streptococci, <i>Corynebacterium</i> spp., <i>Listeria monocytogenes</i> , and <i>Micrococcus</i> spp. (1357 strains). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 239-240.	0.8	18
587	Ceftaroline activity tested against contemporary Latin American bacterial pathogens (2011). <i>Brazilian Journal of Infectious Diseases</i> , 2014, 18, 187-195.	0.3	18
588	Antimicrobial activity of cefotaxime tested against infrequently isolated pathogenic species (unusual) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf</i>	0.8	17
589	Comparative antistreptococcal activity of two newer fluoroquinolones, levofloxacin and sparfloxacin. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 199-201.	0.8	17
590	Antimicrobial activity evaluations of gatifloxacin, a new fluoroquinolone: contemporary pathogen results from a global antimicrobial resistance surveillance program (SENTRY, 1997). <i>Clinical Microbiology and Infection</i> , 1999, 5, 540-546.	2.8	17
591	Antimicrobial susceptibility of quinupristin/dalfopristin tested against gram-positive cocci from Latin America: results from the Global SMART (GSMART) surveillance study. <i>Brazilian Journal of Infectious Diseases</i> , 2001, 5, 21-31.	0.3	17
592	In vitro evaluation of AZD2563, a new oxazolidinone, tested against unusual gram-positive species. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 42, 119-122.	0.8	17
593	Antimicrobial activity of broad-spectrum agents tested against gram-negative bacilli resistant to ceftazidime: report from the SENTRY Antimicrobial Surveillance Program (North America, 2001). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 379-382.	0.8	17
594	Emergence of Two <i>Klebsiella pneumoniae</i> Isolates Harboring Plasmid-Mediated CTX-M-15 β -Lactamase in Taiwan. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 362-363.	1.4	17

#	ARTICLE	IF	CITATIONS
595	Case Report of <i>Aurantimonas altamirensis</i> Bloodstream Infection. <i>Journal of Clinical Microbiology</i> , 2009, 47, 514-515.	1.8	17
596	Update on Linezolid <i>In Vitro</i> Activity through the Zyvox Annual Appraisal of Potency and Spectrum Program, 2013. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 2454-2457.	1.4	17
597	Ceftriaxone: A summary of in vitro antibacterial qualities including recommendations for susceptibility tests with 30- μ g disks. <i>Diagnostic Microbiology and Infectious Disease</i> , 1983, 1, 295-311.	0.8	16
598	Fluoroquinolone (Lomefloxacin) International Surveillance Trial: A report of 30 months of monitoring in vitro activity. <i>American Journal of Medicine</i> , 1992, 92, S52-S57.	0.6	16
599	Cross-resistance analysis for clinafloxacin compared with ciprofloxacin, fleroxacin, ofloxacin, and sparfloxacin using a predictor panel of ciprofloxacin-resistant bacteria. <i>Journal of Antimicrobial Chemotherapy</i> , 1995, 36, 431-434.	1.3	16
600	Antimicrobial activity of SCH27899 (Ziracin [®]), a novel everninomicin derivative, tested against <i>Streptococcus</i> spp.: disk diffusion/Etest method evaluations and quality control guidelines. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 33, 19-25.	0.8	16
601	Summation: β -lactam resistance surveillance in the Asia-Western Pacific Region. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 333-338.	0.8	16
602	Comparison of the activity of two broad-spectrum cephalosporins tested against 2,299 strains of <i>Pseudomonas aeruginosa</i> isolated at 38 North American medical centers participating in the SENTRY antimicrobial surveillance program, 1997-1998. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 36, 125-129.	0.8	16
603	Antimicrobial Activity of DC-159a, a New Fluoroquinolone, against 1,149 Recently Collected Clinical Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 3763-3775.	1.4	16
604	Codetection of <i>bla</i> -OXA-23-Like Gene (<i>bla</i> -OXA-133) and <i>bla</i> -OXA-58 in <i>Acinetobacter radioresistens</i> : Report from the SENTRY Antimicrobial Surveillance Program. <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 843-844.	1.4	16
605	Susceptibility of <i>Klebsiella</i> spp. to colistin and polymyxin B: results from the SENTRY Antimicrobial Surveillance Program (2006-2009). <i>International Journal of Antimicrobial Agents</i> , 2011, 37, 174-175.	1.1	16
606	Ceftaroline Potency Among 9 US Census Regions: Report From the 2010 AWARE Program. <i>Clinical Infectious Diseases</i> , 2012, 55, S194-S205.	2.9	16
607	<i>Streptococcus sanguinis</i> Isolate Displaying a Phenotype with Cross-Resistance to Several rRNA-Targeting Agents. <i>Journal of Clinical Microbiology</i> , 2013, 51, 2728-2731.	1.8	16
608	In vitro activity of dalbavancin against multidrug-resistant <i>Staphylococcus aureus</i> and streptococci from patients with documented infections in Europe and surrounding regions (2011-2013). <i>International Journal of Antimicrobial Agents</i> , 2016, 47, 495-499.	1.1	16
609	New insights into the activity of third-generation cephalosporins against pneumonia-causing bacteria. <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 73-80.	0.8	15
610	Recent trends in the college of American pathologists proficiency results for antimicrobial susceptibility testing: Preparing for CLIA '88. <i>Clinical Microbiology Newsletter</i> , 1992, 14, 33-37.	0.4	15
611	In-vitro comparison of DU-6859a, a novel fluoroquinolone, with other quinolones and oral cephalosporins tested against 5086 recent clinical isolates. <i>Journal of Antimicrobial Chemotherapy</i> , 1993, 32, 877-884.	1.3	15
612	Antimicrobial activity of FK-037, a new broad-spectrum cephalosporin international in vitro comparison with cefepime and ceftazidime. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 18, 167-173.	0.8	15

#	ARTICLE	IF	CITATIONS
613	Vitek GPS card susceptibility testing accuracy using direct inoculation from Bactec 9240 blood culture bottles. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 24, 109-112.	0.8	15
614	Clindamycin resistance among erythromycin-resistant <i>Streptococcus pneumoniae</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 25, 201-204.	0.8	15
615	In Vitro Evaluation of Sparfloxacin Activity and Spectrum Against 24,940 Pathogens Isolated in the United States and Canada, the Final Analysis. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 31, 313-325.	0.8	15
616	Phenotypic and genotypic characterizations of chinese strains of <i>Escherichia coli</i> producing extended-spectrum β -lactamases. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 34, 159-164.	0.8	15
617	Gatifloxacin Phase IV surveillance trial (TeqCES Study) utilizing 5000 primary care physician practices: report of pathogens isolated and susceptibility patterns in community-acquired respiratory tract infections. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 44, 77-84.	0.8	15
618	Multicenter assessment of the linezolid spectrum and activity using the disk diffusion and Etest methods: report of the Zyvox [®] Antimicrobial Potency Study in Latin America (LA-ZAPS). <i>Brazilian Journal of Infectious Diseases</i> , 2002, 6, 100-9.	0.3	15
619	Quality Control Guidelines for MIC Susceptibility Testing of Omiganan Pentahydrochloride (MBI 226), a Novel Antimicrobial Peptide. <i>Journal of Clinical Microbiology</i> , 2004, 42, 1386-1387.	1.8	15
620	Activity of gatifloxacin tested against isolates from pediatric patients: report from the SENTRY Antimicrobial Surveillance Program (North America, 1998-2003). <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 55, 157-164.	0.8	15
621	Antimicrobial activity and spectrum of daptomycin: results from the surveillance program in Australia and New Zealand (2008). <i>Pathology</i> , 2010, 42, 470-473.	0.3	15
622	Clonal Dissemination of <i>Klebsiella pneumoniae</i> Carbapenemase KPC-3 in Long Beach, California. <i>Journal of Clinical Microbiology</i> , 2010, 48, 623-625.	1.8	15
623	Evaluation of quinolone resistance-determining region mutations and efflux pump expression in <i>Neisseria meningitidis</i> resistant to fluoroquinolones. <i>Diagnostic Microbiology and Infectious Disease</i> , 2012, 72, 263-266.	0.8	15
624	Spectrum and potency of ceftaroline tested against leading pathogens causing community-acquired respiratory tract infections in Europe (2010). <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 86-88.	0.8	15
625	Activity of Ceftaroline-Avibactam Tested Against Contemporary Enterobacteriaceae Isolates Carrying β -Lactamases Prevalent in the United States. <i>Microbial Drug Resistance</i> , 2014, 20, 436-440.	0.9	15
626	Ceftaroline activity against organisms isolated from respiratory tract infections in USA hospitals: results from the AWARE program, 2009-2011. <i>Diagnostic Microbiology and Infectious Disease</i> , 2014, 78, 437-442.	0.8	15
627	<i>Klebsiella pneumoniae</i> Isolate from a New York City Hospital Belonging to Sequence Type 258 and Carrying <i>bla</i> KPC-2 and <i>bla</i> VIM-4. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 1924-1927.	1.4	15
628	Performance of BD Max StaphSR for Screening of Methicillin-Resistant <i>Staphylococcus aureus</i> Isolates among a Contemporary and Diverse Collection from 146 Institutions Located in Nine U.S. Census Regions: Prevalence of <i>mecA</i> Dropout Mutants. <i>Journal of Clinical Microbiology</i> , 2016, 54, 204-207.	1.8	15
629	Automation of polymerase chain reaction tests. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 181-185.	0.8	14
630	Ciprofloxacin as broad-spectrum empiric therapy—are fluoroquinolones still viable monotherapeutic agents compared with β -lactams: Data from the MYSTIC Program (US)?. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 42, 213-215.	0.8	14

#	ARTICLE	IF	CITATIONS
631	Microbiologic characterization of isolates from a dalbavancin clinical trial for catheter-related bloodstream infections. <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 54, 83-87.	0.8	14
632	Selection of a surrogate β -lactam testing agent for initial susceptibility testing of doripenem, a new carbapenem. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 59, 467-472.	0.8	14
633	Daptomycin Activity Tested Against Linezolid-Nonsusceptible Gram-Positive Clinical Isolates. <i>Microbial Drug Resistance</i> , 2009, 15, 245-249.	0.9	14
634	Antimicrobial activity of doripenem tested against prevalent Gram-positive pathogens: results from a global surveillance study (2003-2007). <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 63, 440-446.	0.8	14
635	Analysis of 5-year trends in daptomycin activity tested against <i>Staphylococcus aureus</i> and enterococci from European and US hospitals (2009-2013). <i>Journal of Global Antimicrobial Resistance</i> , 2015, 3, 161-165.	0.9	14
636	Results from Oritavancin Resistance Surveillance Programs (2011 to 2014): Clarification for Using Vancomycin as a Surrogate To Infer Oritavancin Susceptibility. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 3174-3177.	1.4	14
637	Activity of Plazomicin Tested against <i>Enterobacteriales</i> Isolates Collected from U.S. Hospitals in 2016-2017: Effect of Different Breakpoint Criteria on Susceptibility Rates among Aminoglycosides. <i>Antimicrobial Agents and Chemotherapy</i> , 2020, 64, .	1.4	14
638	Cefaclor and Cefatrizine, New Investigational Orally Administered Cephalosporins: In-vitro Collaborative Evaluation Against Clinical Bacterial Isolates and Comparison with Related Antimicrobics. <i>American Journal of Clinical Pathology</i> , 1979, 72, 578-585.	0.4	13
639	Cross-resistance analysis for DU-6859a, a new fluoroquinolone, compared to six structurally similar compounds (ciprofloxacin, clinafloxacin, fleroxacin, levofloxacin, ofloxacin, and sparfloxacin). <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 21, 51-54.	0.8	13
640	Anti- <i>Legionella</i> activity of trovafloxacin compared with seven other antimicrobial agents including an intermethod evaluation. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 55-57.	0.8	13
641	Multicenter Evaluation of the In Vitro Activity of Six Broad-Spectrum β -Lactam Antimicrobial Agents in Puerto Rico. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 30, 113-119.	0.8	13
642	BMS284756 (formerly T-3811, a des-fluoroquinolone) potency and spectrum tested against over 10 000 bacterial bloodstream infection isolates from the SENTRY antimicrobial surveillance programme (2000). <i>Journal of Antimicrobial Chemotherapy</i> , 2002, 49, 851-855.	1.3	13
643	Performance of Fusidic Acid (CEM-102) Susceptibility Testing Reagents: Broth Microdilution, Disk Diffusion, and Etest Methods as Applied to <i>Staphylococcus aureus</i> . <i>Journal of Clinical Microbiology</i> , 2010, 48, 972-976.	1.8	13
644	Disk Diffusion and MIC Quality Control Ranges for BC-3205 and BC-3781, Two Novel Pleuromutilin Antibiotics. <i>Journal of Clinical Microbiology</i> , 2012, 50, 3361-3364.	1.8	13
645	Treatment of Methicillin-Susceptible <i>Staphylococcus aureus</i> Osteoarticular and Prosthetic Joint Infections: Using the Oxacillin Minimum Inhibitory Concentration to Guide Appropriate Ceftriaxone Use. <i>Clinical Infectious Diseases</i> , 2013, 57, 161-162.	2.9	13
646	Daptomycin Activity against Uncommonly Isolated Streptococcal and Other Gram-Positive Species Groups. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 6378-6380.	1.4	13
647	Ceftazidime-Avibactam Activity against Aerobic Gram Negative Organisms Isolated from Intra-Abdominal Infections in United States Hospitals, 2012-2014. <i>Surgical Infections</i> , 2016, 17, 473-478.	0.7	13
648	Activity of telavancin against Gram-positive pathogens isolated from bone and joint infections in North American, Latin American, European and Asia-Pacific nations. <i>Diagnostic Microbiology and Infectious Disease</i> , 2017, 88, 184-187.	0.8	13

#	ARTICLE	IF	CITATIONS
649	Use of the E-test to assess macrolide-lincosamide resistance patterns among peptostreptococcus species. Antimicrobial Newsletter, 1992, 8, 45-49.	0.9	12
650	The fourth-generation cephalosporins: Antimicrobial activity and spectrum definitions using cefpirome as an example. Antimicrobial Newsletter, 1993, 9, 9-16.	0.9	12
651	Relationships between Patient- and Institution-Specific Variables and Decreased Antimicrobial Susceptibility of Gram-Negative Pathogens. Clinical Infectious Diseases, 2003, 37, 344-350.	2.9	12
652	Garenoxacin activity against isolates from patients hospitalized with community-acquired pneumonia and multidrug-resistant Streptococcus pneumoniae. Diagnostic Microbiology and Infectious Disease, 2007, 58, 1-7.	0.8	12
653	Activity of daptomycin against Gram-positive bacterial isolates from Indian medical centres (2006-2007). International Journal of Antimicrobial Agents, 2009, 34, 497-499.	1.1	12
654	Comment on: Role of changes in the L3 loop of the active site in the evolution of enzymatic activity of VIM-type metallo- β -lactamases. Journal of Antimicrobial Chemotherapy, 2011, 66, 684-685.	1.3	12
655	Activity of JNJ-Q2, a new fluoroquinolone, tested against contemporary pathogens isolated from patients with community-acquired bacterial pneumonia. International Journal of Antimicrobial Agents, 2012, 39, 321-325.	1.1	12
656	Activity of ceftaroline and comparator agents tested against contemporary Gram-positive and -negative (2011) isolates collected in Europe, Turkey, and Israel. Journal of Chemotherapy, 2014, 26, 202-210.	0.7	12
657	Antimicrobial Activity of Ceftaroline Tested against Staphylococcus aureus from Surgical Skin and Skin Structure Infections in US Medical Centers. Surgical Infections, 2016, 17, 443-447.	0.7	12
658	In-vitro activity of FK-037, a new parenteral cephalosporin. Journal of Antimicrobial Chemotherapy, 1994, 33, 137-144.	1.3	11
659	Validation of NCCLS macrolide (azithromycin, clarithromycin, and erythromycin) interpretive criteria for Haemophilus influenzae tested with the Haemophilus test medium. Diagnostic Microbiology and Infectious Disease, 1994, 18, 243-249.	0.8	11
660	In vitro evaluation of contemporary β -lactam drugs tested against viridans group and β -haemolytic streptococci. Diagnostic Microbiology and Infectious Disease, 1997, 27, 151-154.	0.8	11
661	In vitro evaluation of cefepime and other broad-spectrum β -lactams for isolates in Malaysia and Singapore medical centers. Diagnostic Microbiology and Infectious Disease, 1999, 35, 277-283.	0.8	11
662	In vitro evaluation of broad-spectrum β -lactams in the Philippines medical centers: role of fourth-generation cephalosporins. Diagnostic Microbiology and Infectious Disease, 1999, 35, 291-297.	0.8	11
663	In vitro evaluation of cefepime and other broad-spectrum β -lactams in eight medical centers in Thailand. Diagnostic Microbiology and Infectious Disease, 1999, 35, 325-331.	0.8	11
664	Characterization of extended spectrum β -lactamase-producing Klebsiella pneumoniae from Beijing, China. International Journal of Antimicrobial Agents, 2001, 18, 185-188.	1.1	11
665	In Vitro Evaluation of AZD2563, a Novel Oxazolidinone, against 603 Recent Staphylococcal Isolates. Antimicrobial Agents and Chemotherapy, 2002, 46, 2662-2664.	1.4	11
666	Genotypic Characterization of Carbapenem-Nonsusceptible Acinetobacterspp. Isolated in Latin America. Microbial Drug Resistance, 2004, 10, 286-291.	0.9	11

#	ARTICLE	IF	CITATIONS
667	Potency and spectrum reevaluation of cefdinir tested against pathogens causing skin and soft tissue infections: A sample of North American isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 49, 283-287.	0.8	11
668	Quality Control Guidelines for Susceptibility Testing of Retapamulin (SB-275833) by Reference and Standardized Methods. <i>Journal of Clinical Microbiology</i> , 2005, 43, 6212-6213.	1.8	11
669	Use of in vitro susceptibility and pathogen prevalence data to model the expected clinical success rates of tigecycline and other commonly used antimicrobials for empirical treatment of complicated skin and skin-structure infections. <i>International Journal of Antimicrobial Agents</i> , 2007, 30, 514-520.	1.1	11
670	Contemporary potencies of minocycline and tetracycline HCL tested against Gram-positive pathogens: SENTRY Program results using CLSI and EUCAST breakpoint criteria. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 402-405.	0.8	11
671	In vitro spectrum of pexiganan activity; bactericidal action and resistance selection tested against pathogens with elevated MIC values to topical agents. <i>Diagnostic Microbiology and Infectious Disease</i> , 2016, 86, 66-69.	0.8	11
672	Ceftaroline Activity Against Multidrug-Resistant <i>Streptococcus pneumoniae</i> from U.S. Medical Centers (2014) and Molecular Characterization of a Single Ceftaroline Nonsusceptible Isolate. <i>Microbial Drug Resistance</i> , 2017, 23, 571-579.	0.9	11
673	1-N-(S-3-amino-2-hydroxypropionyl) gentamicin B (Sch 21420): A collaborative in vitro susceptibility comparison with amikacin and gentamicin against 12,984 clinical bacterial isolates. <i>Current Microbiology</i> , 1978, 1, 359-364.	1.0	10
674	Antimicrobial activity and spectrum of ceftibuten (7432-S, SCH 39720) a review of United States and Canadian results. <i>Diagnostic Microbiology and Infectious Disease</i> , 1991, 14, 37-43.	0.8	10
675	Activity of the quinupristin-dalfopristin combination (RP-59500; Synercid) tested against vancomycin-resistant <i>Enterococcus</i> species. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 24, 59-60.	0.8	10
676	Quality control guidelines for amoxicillin, amoxicillin-clavulanate, azithromycin, piperacillin-tazobactam, roxithromycin, ticarcillin, ticarcillin-clavulanate, trovafloxacin (CP 99,219), U-100592, and U-100766 for various National Committee for Clinical Laboratory Standards susceptibility testing methods Results from multicenter trials. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 24, 87-91.	0.8	10
677	Antimicrobial activity and spectrum of sparfloxacin tested against erythromycin-resistant <i>Streptococcus pneumoniae</i> isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 24, 113-116.	0.8	10
678	Susceptibility testing interpretive criteria for levofloxacin when testing respiratory pathogens, <i>Haemophilus influenzae</i> and <i>Moraxella catarrhalis</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 24, 155-160.	0.8	10
679	Antimicrobial characteristics of quinupristin/dalfopristin (Synercid® at 30:70 ratio) compared to alternative ratios for in vitro testing. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 27, 129-138.	0.8	10
680	Avoparcin, a glycopeptide used in animal foods: Antimicrobial spectrum and potency tested against human isolates from the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 29, 241-248.	0.8	10
681	In vitro evaluation of cefepime and other broad-spectrum β -lactams against bacteria from Indonesian medical centers. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 285-290.	0.8	10
682	Surveillance in Taiwan Using Molecular Epidemiology for Extended-Spectrum Beta-Lactamase-Producing <i>Klebsiella pneumoniae</i> . <i>Infection Control and Hospital Epidemiology</i> , 2004, 25, 812-818.	1.0	10
683	Cefdinir activity against contemporary North American isolates from community-acquired urinary tract infections. <i>International Journal of Antimicrobial Agents</i> , 2005, 25, 89-92.	1.1	10
684	Pharmacodynamics of cefprozil against <i>Haemophilus influenzae</i> in an in vitro pharmacodynamic model. <i>Diagnostic Microbiology and Infectious Disease</i> , 2006, 56, 379-386.	0.8	10

#	ARTICLE	IF	CITATIONS
685	Interim susceptibility testing for ceftaroline, a new MRSA-active cephalosporin: selecting potent surrogate β -lactam markers to predict ceftaroline activity against clinically indicated species. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 89-93.	0.8	10
686	Retrospective Molecular Analysis of DIM-1 Metallo- β -Lactamase Discovered in <i>Pseudomonas stutzeri</i> from India in 2000. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 596-598.	1.4	10
687	Antimicrobial Resistance Surveillance and New Drug Development. <i>Open Forum Infectious Diseases</i> , 2019, 6, S5-S13.	0.4	10
688	Bactericidal activity of cefotaxime, desacetylcefotaxime, rifampin, and various combinations tested at cerebrospinal fluid levels against penicillin-resistant <i>Streptococcus pneumoniae</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 22, 119-123.	0.8	9
689	In vitro antimicrobial activity and MIC quality control guidelines of RPR 106972 (RPR 112808/RPR106950): A novel orally administered streptogramin combination. <i>Diagnostic Microbiology and Infectious Disease</i> , 1997, 28, 139-147.	0.8	9
690	In vitro evaluation of broad-spectrum β -lactams tested in medical centers in Korea: role of fourth-generation cephalosporins. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 317-323.	0.8	9
691	Serious streptococcal infections produced by isolates resistant to streptogramins (quinupristin/dalfopristin): Case reports from the SENTRY antimicrobial surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2000, 36, 269-272.	0.8	9
692	Changing Antimicrobial Susceptibility Patterns among <i>Streptococcus pneumoniae</i> and <i>Haemophilus influenzae</i> from Brazil: Report from the SENTRY Antimicrobial Surveillance Program (1998-2004). <i>Microbial Drug Resistance</i> , 2006, 12, 91-98.	0.9	9
693	Antimicrobial activity of cefditoren tested against contemporary (2004-2006) isolates of <i>Haemophilus influenzae</i> and <i>Moraxella catarrhalis</i> responsible for community-acquired respiratory tract infections in the United States. <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 61, 240-244.	0.8	9
694	In vitro potency of doripenem tested against an international collection of rarely isolated bacterial pathogens. <i>Diagnostic Microbiology and Infectious Disease</i> , 2009, 63, 434-439.	0.8	9
695	Determination of disk diffusion and MIC quality control guidelines for GSK2251052: a novel boron-containing antibacterial. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 437-439.	0.8	9
696	Educational antimicrobial susceptibility testing as a critical component of microbiology laboratory proficiency programs: American Proficiency Institute results for 2007-2011. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 75, 357-360.	0.8	9
697	Telavancin activity tested against Gram-positive clinical isolates from European, Russian and Israeli hospitals (2011-2013) using a revised broth microdilution testing method: redefining the baseline activity of telavancin. <i>Journal of Chemotherapy</i> , 2016, 28, 83-88.	0.7	9
698	Perfil de sensibilidade a antimicrobianos de bactérias isoladas do trato respiratório baixo de pacientes com pneumonia internados em hospitais brasileiros: resultados do Programa SENTRY, 1997 e 1998. <i>Jornal De Pneumologia</i> , 2001, 27, 59-67.	0.1	9
699	Antimicrobial activity of Ro 19-5247 (T-2525), a new oral cephalosporin, tested against 7,745 recent clinical isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 1987, 6, 193-198.	0.8	8
700	Antimicrobial activity of FK-037 against class I β -lactamase producing species resistant to ceftazidime: a multi-laboratory clinical isolate sample. <i>Journal of Antimicrobial Chemotherapy</i> , 1993, 32, 654-656.	1.3	8
701	Two investigational glycolcyclines, DMG-DMDOT and DMG-MINO antimicrobial activity studied against gram-positive species. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 24, 53-57.	0.8	8
702	Comparative in-vitro activity of quinupristin/ dalfopristin (RP 59500) tested against penicillin-and macrolide-resistant pneumococci by the Etest. <i>Journal of Antimicrobial Chemotherapy</i> , 1996, 38, 553-555.	1.3	8

#	ARTICLE	IF	CITATIONS
703	In vitro activity of linezolid (U-100766) against <i>Haemophilus influenzae</i> measured by three different susceptibility testing methods. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 39, 49-53.	0.8	8
704	Quality control guidelines for MIC susceptibility testing of NVP PDF-713: a novel peptide deformylase inhibitor. <i>International Journal of Antimicrobial Agents</i> , 2003, 22, 84-86.	1.1	8
705	Tigecycline Disk Diffusion Breakpoints of <i>Acinetobacter</i> spp.: a Clinical Point of View. <i>Journal of Clinical Microbiology</i> , 2007, 45, 2095-2096.	1.8	8
706	In71, an <i>Enterobacter cloacae</i> bla _{VIM-1} -Carrying Integron Related to In70.2 from Italian <i>Pseudomonas aeruginosa</i> Isolates: A SENTRY Antimicrobial Surveillance Program Report. <i>Microbial Drug Resistance</i> , 2007, 13, 130-134.	0.9	8
707	In vitro activity of garenoxacin tested against a worldwide collection of ciprofloxacin-susceptible and ciprofloxacin-resistant Enterobacteriaceae strains (1999–2004). <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 58, 27-32.	0.8	8
708	Antimicrobial activity of daptomycin and selected comparators tested against bloodstream <i>Staphylococcus aureus</i> isolates from hemodialysis patients. <i>International Journal of Infectious Diseases</i> , 2009, 13, 291-295.	1.5	8
709	Surveillance of JNJ-Q2 activity tested against <i>Staphylococcus aureus</i> and beta-hemolytic streptococci as a component of the 2010 sentry antimicrobial surveillance program. <i>Diagnostic Microbiology and Infectious Disease</i> , 2011, 71, 415-420.	0.8	8
710	<i>Candida glabrata</i> : Multidrug Resistance and Increased Virulence in a Major Opportunistic Fungal Pathogen. <i>Current Fungal Infection Reports</i> , 2012, 6, 154-164.	0.9	8
711	Post- ¹² -Lactamase-Inhibitor Effect of Tazobactam in Combination with Ceftolozane on Extended-Spectrum- ¹² -Lactamase-Producing Strains. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 2434-2437.	1.4	8
712	Validation of a Commercial Dry-Form Broth Microdilution Device (Sensititre) for Testing Tedizolid, a New Oxazolidinone. <i>Journal of Clinical Microbiology</i> , 2015, 53, 657-659.	1.8	8
713	Microbiological Assessment of Polymyxin B Components Tested Alone and in Combination. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 7823-7825.	1.4	8
714	Analysis of Vancomycin Susceptibility Testing Results for Presumptive Categorization of Telavancin. <i>Journal of Clinical Microbiology</i> , 2015, 53, 2727-2730.	1.8	8
715	Prevalence of macrolide–lincosamide resistance and multidrug resistance phenotypes in streptococcal isolates causing infections in European hospitals: Evaluation of the in vitro activity of oritavancin and comparator agents. <i>Journal of Global Antimicrobial Resistance</i> , 2017, 8, 28-32.	0.9	8
716	Old In Vitro Antimicrobial Breakpoints Are Misleading Stewardship Efforts, Delaying Adoption of Innovative Therapies, and Harming Patients. <i>Open Forum Infectious Diseases</i> , 2020, 7, ofaa084.	0.4	8
717	Evaluations of Antimicrobial Susceptibility Test Proficiency by the College of American Pathologists Survey Program: A Clarification of Quality Control Recommendations. <i>American Journal of Clinical Pathology</i> , 1982, 78, 168-172.	0.4	7
718	In vitro activity of RU29246 the metabolite of a new HR916 cephalosporin ester. <i>Diagnostic Microbiology and Infectious Disease</i> , 1991, 14, 473-483.	0.8	7
719	Activity of two novel fluoroquinolones (DU-6859a and DV-7751a) tested against glycopeptide-resistant enterococcal isolates. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 23, 123-127.	0.8	7
720	In vitro evaluation of cefepime and other broad-spectrum ¹² -lactams in Taiwan medical centers. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 299-305.	0.8	7

#	ARTICLE	IF	CITATIONS
721	Preliminary susceptibility testing guidelines for AZD2563, a long-acting oxazolidinone. <i>International Journal of Antimicrobial Agents</i> , 2004, 23, 6-10.	1.1	7
722	<i>Klebsiella pneumoniae</i> Carbapenemase-Producing Enterobacteriaceae Testing Susceptible to Cefepime by Reference Methods. <i>Journal of Clinical Microbiology</i> , 2013, 51, 2388-2390.	1.8	7
723	Validation of Sensititre Dry-Form Broth Microdilution Panels for Susceptibility Testing of Ceftazidime-Avibactam, a Broad-Spectrum-β-Lactamase Inhibitor Combination. <i>Antimicrobial Agents and Chemotherapy</i> , 2015, 59, 5036-5039.	1.4	7
724	Ability of the Modified Vitek Card To Detect Coagulase-Negative Staphylococci with <i>mecA</i> and Oxacillin-Resistant Phenotypes. <i>Journal of Clinical Microbiology</i> , 1999, 37, 2122-2123.	1.8	7
725	Susceptibility testing interpretive criteria and drug stability for cefdinir, cefetamet, and cefpodoxime against <i>Neisseria gonorrhoeae</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 1992, 15, 685-691.	0.8	6
726	Prediction of piperacillin-tazobactam susceptibility among Enterobacteriaceae, <i>Pseudomonas aeruginosa</i> , and other bacteria using ticarcillin-clavulanic acid, ceftazidime, and other broad-spectrum antimicrobial in vitro test results. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 20, 143-149.	0.8	6
727	Nosocomial Transmission of <i>Serratia odorifera</i> Biogroup 2: Case Report Demonstration by Macrorestriction Analysis of Chromosomal DNA Using Pulsed-Field Gel Electrophoresis. <i>Infection Control and Hospital Epidemiology</i> , 1994, 15, 390-393.	1.0	6
728	Comparative activity of twelve beta-lactam drugs tested against penicillin-resistant <i>Streptococcus pneumoniae</i> from five medical centers: Effects of serum protein and capsular material on potency and spectrum as measured by reference tests. <i>Diagnostic Microbiology and Infectious Disease</i> , 1996, 25, 137-141.	0.8	6
729	Development of in vitro susceptibility testing methods for gemifloxacin (formerly LB20304a or) Tj ETQq1 1 0.784314 rgBT /Overlock. <i>Journal of Clinical Microbiology</i> , 1999, 35, 227-234.	0.8	6
730	Proposed quality control guidelines for National Committee for Clinical Laboratory Standards Susceptibility Tests using the veterinary antimicrobial agent tiamulin. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 40, 67-70.	0.8	6
731	A pragmatic approach to identify extended-spectrum β-lactamase-producing <i>Klebsiella pneumoniae</i> in Taiwan: in vitro activity of newer and established antimicrobial agents. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 48, 277-282.	0.8	6
732	Disk diffusion quality control guidelines for NVP-PDF 713: a novel peptide deformylase inhibitor. <i>Diagnostic Microbiology and Infectious Disease</i> , 2004, 48, 55-57.	0.8	6
733	Ceftaroline activity tested against viridans group streptococci from US hospitals. <i>Diagnostic Microbiology and Infectious Disease</i> , 2016, 84, 232-235.	0.8	6
734	Telavancin activity tested against a collection of <i>Staphylococcus aureus</i> isolates causing pneumonia in hospitalized patients in the United States (2013-2014). <i>Diagnostic Microbiology and Infectious Disease</i> , 2016, 86, 300-302.	0.8	6
735	Antimicrobial activity of ceftazidime-avibactam and comparator agents when tested against bacterial isolates causing infection in cancer patients (2013-2014). <i>Diagnostic Microbiology and Infectious Disease</i> , 2017, 87, 261-265.	0.8	6
736	Cefpirome (HR810) disk diffusion susceptibility tests: Confirmation of interpretive criteria using cefotaxime and cefoperazone-resistant strains and effects of blood supplemented media. <i>Diagnostic Microbiology and Infectious Disease</i> , 1986, 4, 345-349.	0.8	5
737	Critical assessment of the newer non-quinolone oral antimicrobial agents. <i>Antimicrobial Newsletter</i> , 1989, 6, 53-60.	0.9	5
738	Preliminary interpretive criteria for disk diffusion susceptibility testing of SCH 27899, a compound in the everninomicin class of antimicrobial agents. <i>Diagnostic Microbiology and Infectious Disease</i> , 1995, 23, 157-160.	0.8	5

#	ARTICLE	IF	CITATIONS
739	In Vitro Activity of Cefepime and Other Broad-Spectrum β -Lactams Tested Against 129 mec A-negative Staphylococcus spp. Isolates: A Multicenter Sample. <i>Diagnostic Microbiology and Infectious Disease</i> , 1998, 30, 65-69.	0.8	5
740	Evaluation of the in vitro antimicrobial activity of cefepime compared to other broad-spectrum β -lactams tested against recent clinical isolates from 10 Chinese hospitals. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 35, 135-142.	0.8	5
741	Comparative antimicrobial activity of ABT-773, a novel ketolide, tested against drug-resistant Gram-positive cocci and Haemophilus influenzae. <i>International Journal of Antimicrobial Agents</i> , 2001, 17, 451-455.	1.1	5
742	Accuracy of broth microdilution and E test methods for detecting chloramphenicol acetyl transferase mediated resistance in Streptococcus pneumoniae: Geographic variations in the prevalence of resistance in The SENTRY Antimicrobial Surveillance Program (1999). <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 39, 267-269.	0.8	5
743	In vitro evaluation of AZD2563, a new oxazolidinone, tested against beta-haemolytic and viridans group streptococci. <i>Journal of Antimicrobial Chemotherapy</i> , 2002, 49, 1019-1021.	1.3	5
744	Pharmacodynamics in the Evaluation of Drug Regimens. <i>Annals of Pharmacotherapy</i> , 2002, 36, 530-532.	0.9	5
745	Are Enterococcus faecalis Strains with vat (E) in Poultry a Reservoir for Human Streptogramin Resistance? vat (E) Occurrence in Human Enterococcal Bloodstream Infections in North America (SENTRY Antimicrobial Surveillance Program, 2002). <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 360-361.	1.4	5
746	Development of anidulafungin for disk diffusion susceptibility testing against Candida spp.. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 58, 371-374.	0.8	5
747	More potency assay results for generic non-USA lots of piperacillin/tazobactam and initial reports for generic meropenem compounds marketed in the USA. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 110-112.	0.8	5
748	IMP-33, a New IMP Variant Detected in Pseudomonas aeruginosa from Sicily. <i>Antimicrobial Agents and Chemotherapy</i> , 2013, 57, 6401-6403.	1.4	5
749	Genotypic Characterization of Methicillin-Resistant Staphylococcus aureus Recovered at Baseline from Phase 3 Pneumonia Clinical Trials for Ceftobiprole. <i>Microbial Drug Resistance</i> , 2016, 22, 53-58.	0.9	5
750	Sensibilidade a antimicrobianos de bactérias isoladas do trato respiratório de pacientes com infecções respiratórias adquiridas na comunidade: resultados brasileiros do Programa SENTRY de Vigilância de Resistência a Antimicrobianos dos anos de 1997 e 1998. <i>Jornal De Pneumologia</i> , 2001, 27, 25-34.	0.1	5
751	Interpretive criteria for DU-6859a disk diffusion tests using 5- $\frac{1}{4}$ g disks. <i>Diagnostic Microbiology and Infectious Disease</i> , 1994, 18, 125-127.	0.8	4
752	Clinical use of β -lactamase inhibitors in combination with extended-spectrum penicillins. <i>American Journal of Health-System Pharmacy</i> , 1995, 52, S29-S33.	0.5	4
753	Comparative in vitro activity of apacillin alone and combined with Ro 48-1220, a novel penam β -lactamase inhibitor. <i>Clinical Microbiology and Infection</i> , 1995, 1, 86-100.	2.8	4
754	Comparative in vitro activity of alexomycin (U-82127) tested against Escherichia coli, Salmonella spp., and enterococci of animal and human origin. <i>Clinical Microbiology and Infection</i> , 1998, 4, 601-603.	2.8	4
755	Comment on: Linezolid resistance in coagulase-negative staphylococci. <i>Journal of Antimicrobial Chemotherapy</i> , 2006, 58, 899-900.	1.3	4
756	Comparison of Censored Regression and Standard Regression Analyses for Modeling Relationships between Antimicrobial Susceptibility and Patient- and Institution-Specific Variables. <i>Antimicrobial Agents and Chemotherapy</i> , 2006, 50, 62-67.	1.4	4

#	ARTICLE	IF	CITATIONS
757	Comparative activity of linezolid against respiratory tract infection isolates of <i>Staphylococcus aureus</i> : an 11-year report from the SENTRY Antimicrobial Surveillance Program. <i>International Journal of Antimicrobial Agents</i> , 2011, 37, 584-585.	1.1	4
758	Definitions and Epidemiology of <i>Candida</i> Species not Susceptible to Echinocandins. <i>Current Fungal Infection Reports</i> , 2011, 5, 120-127.	0.9	4
759	Detection of NDM-1-producing <i>Enterobacteriaceae</i> in Romania: report of the SENTRY Antimicrobial Surveillance Program. <i>Journal of Medical Microbiology</i> , 2014, 63, 483-484.	0.7	4
760	Accuracy of the Thermo Fisher Scientific (Sensititre [®] , [®]) dry-form broth microdilution MIC product when testing ceftaroline. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015, 81, 280-282.	0.8	4
761	Determination of Disk Diffusion and MIC Quality Control Guidelines for Solithromycin, a Novel Fluoroketolide Antibacterial, against <i>Neisseria gonorrhoeae</i> . <i>Journal of Clinical Microbiology</i> , 2015, 53, 3888-3890.	1.8	4
762	Dalbavancin Activity When Tested against <i>Streptococcus pneumoniae</i> Isolated in Medical Centers on Six Continents (2011 to 2014). <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 3419-3425.	1.4	4
763	Reproducibility of dalbavancin MIC test results and an updated surrogate accuracy analysis of vancomycin MIC values to infer dalbavancin susceptibility (2014). <i>Diagnostic Microbiology and Infectious Disease</i> , 2016, 86, 249-251.	0.8	4
764	The application of in vitro surveillance data for antibacterial dose selection. <i>Current Opinion in Pharmacology</i> , 2017, 36, 130-138.	1.7	4
765	β -Lactam antibiotics: The first- and second-generation cephalosporins. <i>Antimicrobial Newsletter</i> , 1985, 2, 25-28.	0.9	3
766	Microbiological Conclusions. <i>Drugs</i> , 1991, 42, 22-24.	4.9	3
767	Reevaluation of contemporary laboratory methods for detection of antimicrobial resistance among enterococci. <i>Clinical Microbiology and Infection</i> , 1996, 1, 190-194.	2.8	3
768	In vitro assessment of gatifloxacin spectrum and potency tested against <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i> , and <i>Streptococcus pneumoniae</i> isolates from the Asia-Western Pacific component of the SENTRY antimicrobial surveillance program (1998-1999). <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 315-318.	0.8	3
769	Reevaluation of the cefepime minimal inhibitory concentrations and disk diffusion test zone diameter relationship for a worldwide collection of <i>Enterobacteriaceae</i> enriched for extended-spectrum β -lactamase-producing organisms. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 95-99.	0.8	3
770	Reproducibility of daptomycin MIC results using dry-form commercial trays with appropriate supplemental calcium content. <i>International Journal of Antimicrobial Agents</i> , 2005, 25, 274-275.	1.1	3
771	Influence of polysorbate-80 when determining the tigecycline MIC by the reference method. <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 58, 145-146.	0.8	3
772	Fixed-Ratio Combination Testing of an Echinocandin, Anidulafungin, and an Azole, Voriconazole, against 1,467 <i>Candida</i> Species Isolates. <i>Antimicrobial Agents and Chemotherapy</i> , 2010, 54, 4041-4043.	1.4	3
773	Comparative potencies of contemporary generic vancomycin lot: in vitro assay results from nine products and a reference reagent-grade sample. <i>Diagnostic Microbiology and Infectious Disease</i> , 2013, 76, 237-238.	0.8	3
774	Commentary. <i>Pediatric Infectious Disease Journal</i> , 2013, 32, 970-971.	1.1	3

#	ARTICLE	IF	CITATIONS
775	Telavancin activity when tested by a revised susceptibility testing method against uncommonly isolated Gram-positive pathogens responsible for documented infections in hospitals worldwide (2011-2013). <i>Journal of Global Antimicrobial Resistance</i> , 2015, 3, 36-39.	0.9	3
776	Comparison of Amino Acid Decarboxylase and Dihydrolase Results by Moeller, Rapid, and Replicator Plate Methods. <i>Journal of Clinical Microbiology</i> , 1976, 3, 75-76.	1.8	3
777	Spectrum and antimicrobial activity of Alexomycin (PNU-82, 127), a peptide compound projected for use in animal health. <i>Diagnostic Microbiology and Infectious Disease</i> , 1999, 33, 181-186.	0.8	2
778	MYSTIC: closing summary. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 41, 197-198.	0.8	2
779	Evaluation of in vitro susceptibility testing criteria for gemifloxacin when tested against <i>Haemophilus influenzae</i> strains with reduced susceptibility to ciprofloxacin and ofloxacin. <i>Diagnostic Microbiology and Infectious Disease</i> , 2002, 43, 323-326.	0.8	2
780	Determination of disk diffusion and MIC quality control parameters for AZD2563, a novel long-acting oxazolidinone. <i>Diagnostic Microbiology and Infectious Disease</i> , 2003, 45, 73-76.	0.8	2
781	MIC Quality Control Guidelines and Disk Diffusion Test Optimization for CEM-101, a Novel Fluoroketolide. <i>Journal of Clinical Microbiology</i> , 2010, 48, 1470-1473.	1.8	2
782	Oritavancin Activity Tested against Molecularly Characterized Staphylococci and Enterococci Displaying Elevated Linezolid MIC Results. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 3817-3820.	1.4	2
783	Validation of cefditoren MIC quality control ranges by a multi-laboratory study (2001). <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 40, 71-73.	0.8	1
784	MYSTIC preface. <i>Diagnostic Microbiology and Infectious Disease</i> , 2001, 41, 169.	0.8	1
785	Re-evaluations of disk diffusion quality control ranges for 11 drugs: report from the QC working group of the national committee for clinical laboratory standards. <i>International Journal of Antimicrobial Agents</i> , 2002, 20, 289-292.	1.1	1
786	Activity of DX-619 Compared to Other Agents against Viridans Group Streptococci, <i>Streptococcus bovis</i> , and <i>Cardiobacterium hominis</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2006, 50, 4191-4194.	1.4	1
787	Bactericidal activity of cefepime and ceftriaxone tested against <i>Streptococcus pneumoniae</i> . <i>Diagnostic Microbiology and Infectious Disease</i> , 2007, 57, 345-349.	0.8	1
788	Determination of Disk Diffusion and MIC Quality Control Guidelines for JNJ-Q2, a Novel Quinolone. <i>Journal of Clinical Microbiology</i> , 2011, 49, 3009-3011.	1.8	1
789	Reply to Nation et al. <i>Clinical Infectious Diseases</i> , 2013, 57, 1657-1658.	2.9	1
790	Reproducibility assessment of tigecycline MIC results by broth microdilution methods using commercially prepared dry-form panels. <i>Diagnostic Microbiology and Infectious Disease</i> , 2005, 52, 67-69.	0.8	0
791	Proposed quality control ranges for 2-1/4g anidulafungin disk diffusion testing and ability of the method to detect strains with elevated echinocandin MIC values. <i>Diagnostic Microbiology and Infectious Disease</i> , 2008, 62, 474-477.	0.8	0
792	<i>Enterococcus</i> , 0, 953-960.		0

#	ARTICLE	IF	CITATIONS
793	Regional pooling of national data from a small number of sites can be misleading: maybe yes? But data can be complimentary to other studies and valuable to infectious disease physicians!. Diagnostic Microbiology and Infectious Disease, 2014, 80, 91-92.	0.8	0
794	Enterococcus. , 0, , 895-900.		0