

# Masatsune Kainosho

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10815705/publications.pdf>

Version: 2024-02-01

163  
papers

6,131  
citations

81900

39  
h-index

88630

70  
g-index

176  
all docs

176  
docs citations

176  
times ranked

4180  
citing authors

#	ARTICLE	IF	CITATIONS
1	Conformational features and ionization states of Lys side chains in a protein studied using the stereo-array isotope labeling (SAIL) method. <i>Magnetic Resonance</i> , 2021, 2, 223-237.	1.9	0
2	Stereo-Array Isotope Labeling (SAIL) and Related Methods. , 2021, , 1-3.		0
3	Recent developments in isotope-aided NMR methods for supramolecular protein complexes –SAIL aromatic TROSY. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2020, 1864, 129439.	2.4	7
4	Pressure dependence of side chain <sup>1</sup> H and <sup>15</sup> N-chemical shifts in the model peptides Ac-Gly-Gly-Xxx-Ala-NH <sub>2</sub> . <i>Journal of Biomolecular NMR</i> , 2020, 74, 381-399.	2.8	1
5	Aromatic Ring Dynamics, Thermal Activation, and Transient Conformations of a 468 kDa Enzyme by Specific <sup>1</sup> H- <sup>13</sup> C Labeling and Fast Magic-Angle Spinning NMR. <i>Journal of the American Chemical Society</i> , 2019, 141, 11183-11195.	13.7	43
6	Isotope-Aided Methods for Biological NMR Spectroscopy: Past, Present, and Future. , 2018, , 37-61.		5
7	A Numb-Mdm2 fuzzy complex reveals an isoform-specific involvement of Numb in breast cancer. <i>Journal of Cell Biology</i> , 2018, 217, 745-762.	5.2	33
8	Perspective: next generation isotope-aided methods for protein NMR spectroscopy. <i>Journal of Biomolecular NMR</i> , 2018, 71, 119-127.	2.8	14
9	Stable-Isotope-Aided NMR Spectroscopy. , 2018, , 469-486.		0
10	Evolution and diversification of the plant gibberellin receptor GID1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, E7844-E7853.	7.1	51
11	Structural and Functional Analysis of the C-Terminal Region of FliG, an Essential Motor Component of Vibrio Na <sup>+</sup> -Driven Flagella. <i>Structure</i> , 2017, 25, 1540-1548.e3.	3.3	13
12	Pressure dependence of side chain <sup>13</sup> C chemical shifts in model peptides Ac-Gly-Gly-Xxx-Ala-NH <sub>2</sub> . <i>Journal of Biomolecular NMR</i> , 2017, 69, 53-67.	2.8	8
13	<sup>13</sup> C-NMR studies on disulfide bond isomerization in bovine pancreatic trypsin inhibitor (BPTI). <i>Journal of Biomolecular NMR</i> , 2016, 66, 37-53.	2.8	7
14	Highly efficient residue-selective labeling with isotope-labeled Ile, Leu, and Val using a new auxotrophic E. coli strain. <i>Journal of Biomolecular NMR</i> , 2016, 65, 109-119.	2.8	29
15	Stable-Isotope-Aided NMR Spectroscopy. , 2016, , 1-18.		1
16	Differential Large-Amplitude Breathing Motions in the Interface of FKBP12-Drug Complexes. <i>Biochemistry</i> , 2015, 54, 6983-6995.	2.5	24
17	Nano-mole scale sequential signal assignment by <sup>1</sup> H-detected protein solid-state NMR. <i>Chemical Communications</i> , 2015, 51, 15055-15058.	4.1	39
18	Nano-Mole Scale Side-Chain Signal Assignment by <sup>1</sup> H-Detected Protein Solid-State NMR by Ultra-Fast Magic-Angle Spinning and Stereo-Array Isotope Labeling. <i>PLoS ONE</i> , 2015, 10, e0122714.	2.5	16

#	ARTICLE	IF	CITATIONS
19	Automated resonance assignment of the 21 kDa stereo-array isotope labeled thioldisulfide oxidoreductase DsbA. <i>Journal of Magnetic Resonance</i> , 2014, 249, 88-93.	2.1	7
20	Use of H/D isotope effects to gather information about hydrogen bonding and hydrogen exchange rates. <i>Journal of Magnetic Resonance</i> , 2014, 241, 148-154.	2.1	26
21	Expression and purification of a GRAS domain of SLR1, the rice DELLA protein. <i>Protein Expression and Purification</i> , 2014, 95, 248-258.	1.3	21
22	Differential isotope-labeling for Leu and Val residues in a protein by <i>E. coli</i> cellular expression using stereo-specifically methyl labeled amino acids. <i>Journal of Biomolecular NMR</i> , 2013, 57, 237-249.	2.8	35
23	Cell-Free Protein Synthesis Using <i>E. coli</i> Cell Extract for NMR Studies. <i>Advances in Experimental Medicine and Biology</i> , 2012, 992, 167-177.	1.6	14
24	Stereo-Array Isotope Labeling Method for Studying Protein Structure and Dynamics. <i>Advances in Experimental Medicine and Biology</i> , 2012, 992, 83-93.	1.6	10
25	Cell-Free Protein Production for NMR Studies. <i>Methods in Molecular Biology</i> , 2012, 831, 71-84.	0.9	21
26	1B1424 Solution NMR analysis of FUG C-terminal domain derived from Na <sup>+</sup> -driven motor of <i>Vibrio</i> (Proteins: Structure & Function I, Oral Presentation, The 50th Annual Meeting of the Biophysical Society, 2012, 10.1117/1.3610101)		
27	Conformational analysis by quantitative NOE measurements of the $\beta^2$ -proton pairs across individual disulfide bonds in proteins. <i>Journal of Biomolecular NMR</i> , 2012, 52, 127-139.	2.8	13
28	Synthesis of Stereoarray Isotope Labeled (SAIL) Lysine via the $\alpha$ -Head-to-Tail Conversion of SAIL Glutamic Acid. <i>Organic Letters</i> , 2011, 13, 161-163.	4.6	12
29	Hydrogen Exchange Study on the Hydroxyl Groups of Serine and Threonine Residues in Proteins and Structure Refinement Using NOE Restraints with Polar Side-Chain Groups. <i>Journal of the American Chemical Society</i> , 2011, 133, 17420-17427.	13.7	24
30	Exclusively NOESY-based automated NMR assignment and structure determination of proteins. <i>Journal of Biomolecular NMR</i> , 2011, 50, 137-146.	2.8	26
31	Alternative SAIL-Trp for robust aromatic signal assignment and determination of the $\beta^2$ conformation by intra-residue NOEs. <i>Journal of Biomolecular NMR</i> , 2011, 51, 425-435.	2.8	24
32	Hydrogen exchange during cell-free incorporation of deuterated amino acids and an approach to its inhibition. <i>Journal of Biomolecular NMR</i> , 2011, 51, 467-476.	2.8	26
33	Solution NMR Structure of Proteorhodopsin. <i>Angewandte Chemie - International Edition</i> , 2011, 50, 11942-11946.	13.8	162
34	Construction and performance of an NMR tube with a sample cavity formed within magnetic susceptibility-matched glass. <i>Journal of Magnetic Resonance</i> , 2011, 209, 167-173.	2.1	27
35	Application of SAIL phenylalanine and tyrosine with alternative isotope-labeling patterns for protein structure determination. <i>Journal of Biomolecular NMR</i> , 2010, 46, 45-49.	2.8	38
36	$^1\text{H}$ -detected $^1\text{H}$ - $^1\text{H}$ correlation spectroscopy of a stereo-array isotope labeled amino acid under fast magic-angle spinning. <i>Journal of Magnetic Resonance</i> , 2010, 203, 253-256.	2.1	8

#	ARTICLE	IF	CITATIONS
37	Detection of the Sulfhydryl Groups in Proteins with Slow Hydrogen Exchange Rates and Determination of Their Proton/Deuteron Fractionation Factors Using the Deuterium-Induced Effects on the $^{13}\text{C}$ NMR Signals. <i>Journal of the American Chemical Society</i> , 2010, 132, 6254-6260.	13.7	29
38	Automated NMR structure determination of stereo-array isotope labeled ubiquitin from minimal sets of spectra using the SAIL-FLYA system. <i>Journal of Biomolecular NMR</i> , 2009, 44, 261-272.	2.8	27
39	Assymmetric synthesis of (2S,3R)- and (2S,3S)-[2- $^{13}\text{C}$ ;3- $^2\text{H}$ ] glutamic acid. <i>Tetrahedron Letters</i> , 2009, 50, 1482-1484.	1.4	12
40	Hydrogen Exchange Rate of Tyrosine Hydroxyl Groups in Proteins As Studied by the Deuterium Isotope Effect on $^{13}\text{C}$ Chemical Shifts. <i>Journal of the American Chemical Society</i> , 2009, 131, 18556-18562.	13.7	48
41	SAIL " stereo-array isotope labeling. <i>Quarterly Reviews of Biophysics</i> , 2009, 42, 247-300.	5.7	64
42	Protein NMR Study Expanded by the SAIL Method. <i>Seibutsu Butsuri</i> , 2009, 49, 206-209.	0.1	0
43	Recent Developments in Stable-Isotope-Aided Methods for Protein NMR Spectroscopy. , 2008, , 215-222.		0
44	Stable isotope labeling methods for protein NMR spectroscopy. <i>Progress in Nuclear Magnetic Resonance Spectroscopy</i> , 2008, 53, 208-226.	7.5	85
45	Structure of the putative 32 kDa myrosinase-binding protein from <i>Arabidopsis</i> (At3g16450.1) determined by SAIL-NMR. <i>FEBS Journal</i> , 2008, 275, 5873-5884.	4.7	28
46	Solution Structure of the C-terminal Dimerization Domain of SARS Coronavirus Nucleocapsid Protein Solved by the SAIL-NMR Method. <i>Journal of Molecular Biology</i> , 2008, 380, 608-622.	4.2	111
47	Structural Basis of the Role of the NikA Ribbon-Helix-Helix Domain in Initiating Bacterial Conjugation. <i>Journal of Molecular Biology</i> , 2008, 384, 690-701.	4.2	21
48	Stereoselective Synthesis of Triply Isotope-Labeled Ser, Cys, and Ala: Amino Acids for Stereoarray Isotope Labeling Technology. <i>Organic Letters</i> , 2008, 10, 2785-2787.	4.6	18
49	Automated structure determination of proteins with the SAIL-FLYA NMR method. <i>Nature Protocols</i> , 2007, 2, 2896-2902.	12.0	48
50	Biosynthesis of Quinolactacin A, a TNF Production Inhibitor. <i>Journal of Antibiotics</i> , 2006, 59, 418-427.	2.0	19
51	Optimal isotope labelling for NMR protein structure determinations. <i>Nature</i> , 2006, 440, 52-57.	27.8	442
52	Evaluation of stereo-array isotope labeling (SAIL) patterns for automated structural analysis of proteins with CYANA. <i>Magnetic Resonance in Chemistry</i> , 2006, 44, S152-S157.	1.9	32
53	Carbon-13 NMR Method for the Detection of Correlated Hydrogen Exchange at Adjacent Backbone Peptide Amides and Its Application to Hydrogen Exchange in Five Antiparallel $^2$ Strands within the Hydrophobic Core of Streptomyces Subtilisin Inhibitor (SSI). <i>Biochemistry</i> , 2005, 44, 11811-11820.	2.5	15
54	NMR Assignment Methods for the Aromatic Ring Resonances of Phenylalanine and Tyrosine Residues in Proteins. <i>Journal of the American Chemical Society</i> , 2005, 127, 12620-12626.	13.7	46

#	ARTICLE	IF	CITATIONS
55	Structural Basis of a Myosin Phosphatase Inhibitory Protein, CPI-17. <i>Seibutsu Butsuri</i> , 2005, 45, 72-77.	0.1	0
56	Efficient production of isotopically labeled proteins by cell-free synthesis: A practical protocol. <i>Journal of Biomolecular NMR</i> , 2004, 30, 311-325.	2.8	124
57	Letter to the editor: <sup>1</sup> H, <sup>13</sup> C and <sup>15</sup> N backbone assignment of a 32 kDa hypothetical protein from <i>Arabidopsis thaliana</i> , At3g16450.1. <i>Journal of Biomolecular NMR</i> , 2004, 30, 357-358.	2.8	2
58	Phosphorylation-induced conformational change responsible for the function of a myosin phosphatase inhibitor, CPI-17. <i>Science and Technology of Advanced Materials</i> , 2004, 5, 383-386.	6.1	1
59	A New Stable-Isotope-Aided NMR Method for Structural Determinations of Proteins: The SAIL Method. <i>Seibutsu Butsuri</i> , 2004, 44, 200-205.	0.1	2
60	Rotational diffusion tensor of nucleic acids from <sup>13</sup> C NMR relaxation. <i>Journal of Biomolecular NMR</i> , 2003, 27, 133-142.	2.8	49
61	Distinctive Solution Conformation of Phosphatase Inhibitor CPI-17 Substituted with Aspartate at the Phosphorylation-site Threonine Residue. <i>Journal of Molecular Biology</i> , 2003, 326, 1539-1547.	4.2	16
62	Solid-Phase Synthesis of Selectively Labeled DNA: Applications for Multidimensional Nuclear Magnetic Resonance Spectroscopy. <i>Methods in Enzymology</i> , 2002, 338, 261-283.	1.0	14
63	Characterization of the ATP-Binding Domain of the Sarco(endoplasmic Reticulum Ca <sup>2+</sup> -ATPase: Probing Nucleotide Binding by Multidimensional NMR. <i>Biochemistry</i> , 2002, 41, 1156-1164.	2.5	32
64	NMR structure of <i>Streptomyces</i> killer toxin-like protein, SKLP: further evidence for the wide distribution of single-domain <sup>12</sup> I <sup>3</sup> -crystallin superfamily proteins. <i>Journal of Molecular Biology</i> , 2001, 305, 109-120.	4.2	26
65	Solution NMR structure of the myosin phosphatase inhibitor protein CPI-17 shows phosphorylation-induced conformational changes responsible for activation 1 Edited by P. E. Wright. <i>Journal of Molecular Biology</i> , 2001, 314, 839-849.	4.2	38
66	Synthesis of <sup>13</sup> C/D Doubly Labeled Leucines: Probes for Conformational Analysis of the Leucine Side-chain. <i>Journal of Organic Chemistry</i> , 2001, 66, 5919-5922.	3.2	17
67	[ <sup>13</sup> C, <sup>13</sup> C]- and [ <sup>13</sup> C, <sup>1</sup> H]-TROSY in a Triple Resonance Experiment for Ribose Base and Intra-base Correlations in Nucleic Acids. <i>Journal of the American Chemical Society</i> , 2001, 123, 658-664.	13.7	61
68	Developing model systems for the NMR study of substituent effects on the N-H...N hydrogen bond in duplex DNA. <i>Magnetic Resonance in Chemistry</i> , 2001, 39, S159-S165.	1.9	29
69	H...N hydrogen bond lengths in double stranded DNA from internucleotide dipolar couplings. <i>Journal of Biomolecular NMR</i> , 2001, 19, 361-365.	2.8	21
70	Sugar conformation of a stereospecific 2'-R or 2'-S deuterium-labeled DNA decamer studied with proton-proton J coupling constants. <i>Journal of Biomolecular NMR</i> , 2001, 19, 19-31.	2.8	7
71	Backbone <sup>1</sup> H, <sup>13</sup> C, and <sup>15</sup> N resonance assignments of an 18.2 kDa protein, <i>E. coli</i> peptidyl-prolyl cis-trans isomerase b (EPPIb). <i>Journal of Biomolecular NMR</i> , 2000, 18, 75-76.	2.8	15
72	Determination of <sup>2</sup> J(HN) and <sup>1</sup> J(HN) coupling constants across Watson-Crick base pairs in the Antennapedia homeodomain-DNA complex using TROSY. <i>Journal of Biomolecular NMR</i> , 2000, 16, 39-46.	2.8	43

#	ARTICLE	IF	CITATIONS
73	Three-dimensional structure determination of a uniformly labeled molecule by frequency-selective dipolar recoupling under magic-angle spinning. <i>Journal of Biomolecular NMR</i> , 2000, 17, 111-123.	2.8	37
74	Structural comparison between wild-type and P25S human cystatin A by NMR spectroscopy. Does this mutation affect the alpha-helix conformation?. <i>Journal of Structural and Functional Genomics</i> , 2000, 1, 26-42.	1.2	7
75	Studies of physicochemical properties of N-H...N hydrogen bonds in DNA, using selective <sup>15</sup> N-labeling and direct <sup>15</sup> N 1D NMR. <i>Journal of Biomolecular NMR</i> , 2000, 18, 269-277.	2.8	22
76	Trends in Structure and Growth of Higher Fullerenes Isomer Structure of C <sub>86</sub> and C <sub>88</sub> . <i>Molecular Crystals and Liquid Crystals</i> , 2000, 340, 553-558.	0.3	49
77	The NMR Structure of a DNA Dodecamer in an Aqueous Dilute Liquid Crystalline Phase. <i>Journal of the American Chemical Society</i> , 2000, 122, 6190-6200.	13.7	201
78	Direct Observation of Hydrogen Bonding in Biomolecules by NMR. <i>Seibutsu Butsuri</i> , 2000, 40, 379-384.	0.1	1
79	<sup>13</sup> C-NMR Relaxation Analysis of Nucleic Acid Structure and Dynamics. <i>Seibutsu Butsuri</i> , 2000, 40, 191-194.	0.1	0
80	Backbone <sup>1</sup> H, <sup>13</sup> C, and <sup>15</sup> N resonance assignments of <i>Streptomyces subtilisin</i> inhibitor. <i>Journal of Biomolecular NMR</i> , 1999, 14, 285-286.	2.8	0
81	Quantitative Measurement of Transverse and Longitudinal Cross-Correlation between <sup>13</sup> C- <sup>1</sup> H Dipolar Interaction and <sup>13</sup> C Chemical Shift Anisotropy: Application to a <sup>13</sup> C-Labeled DNA Duplex. <i>Journal of Magnetic Resonance</i> , 1999, 136, 169-175.	2.1	17
82	The 2D { <sup>31</sup> P} Spin-Echo-Difference Constant-Time [ <sup>13</sup> C, <sup>1</sup> H]-HMQC Experiment for Simultaneous Determination of <sup>31</sup> H <sup>31</sup> P and <sup>31</sup> C <sup>31</sup> P in <sup>13</sup> C-Labeled Nucleic Acids and Their Protein Complexes. <i>Journal of Magnetic Resonance</i> , 1999, 140, 491-494.	2.1	20
83	Stereodivergent Synthesis of (2S,3S,4R,5R)- and (2S,3S,4R,5S)-[3,4,5-D <sup>3</sup> ]Proline Depending on the Substituent of the <sup>13</sup> -Lactam Ring. <i>Journal of Organic Chemistry</i> , 1999, 64, 9275-9278.	3.2	17
84	Determination of the Complete Structure of a Uniformly Labeled Molecule by Rotational Resonance Solid-State NMR in the Tilted Rotating Frame. <i>Journal of the American Chemical Society</i> , 1999, 121, 4064-4065.	13.7	87
85	Analysis of the relationship between enzyme activity and its internal motion using nuclear magnetic resonance: <sup>15</sup> N relaxation studies of wild-type and mutant lysozyme 1 Edited by P. E. Wright. <i>Journal of Molecular Biology</i> , 1999, 286, 1547-1565.	4.2	53
86	Conformational Changes of the BS2 Operator DNA upon Complex Formation with the Antennapedia Homeodomain Studied by NMR with <sup>13</sup> C/ <sup>15</sup> N-labeled DNA. <i>Journal of Molecular Biology</i> , 1999, 292, 609-617.	4.2	14
87	Differential isotope labeling strategy for determining the structure of myristoylated recoverin by NMR spectroscopy. <i>Journal of Biomolecular NMR</i> , 1998, 11, 135-152.	2.8	25
88	Dual amino acid-selective and site-directed stable-isotope labeling of the human c-Ha-Ras protein by cell-free synthesis. <i>Journal of Biomolecular NMR</i> , 1998, 11, 295-306.	2.8	126
89	NMR with ( <sup>13</sup> C, ( <sup>15</sup> N)-doubly-labeled DNA: The shape Antennapedia homeodomain complex with a 14-mer DNA duplex. <i>Journal of Biomolecular NMR</i> , 1998, 12, 25-37.	2.8	26
90	Stereospecific assignment of H5' and H5 <sup>3</sup> in a (5'R)-/(5'S)-deuterium- labeled DNA decamer for (3) J (HH) determination and unambiguous NOE assignments. <i>Journal of Biomolecular NMR</i> , 1998, 11, 103-109.	2.8	14

#	ARTICLE	IF	CITATIONS
91	NMR structure of the histidine kinase domain of the E. coli osmosensor EnvZ. <i>Nature</i> , 1998, 396, 88-92.	27.8	248
92	DNA Duplex Dynamics: NMR Relaxation Studies of a Decamer with Uniformly <sup>13</sup> C-Labeled Purine Nucleotides. <i>Journal of Magnetic Resonance</i> , 1998, 135, 310-333.	2.1	55
93	Systematic synthesis of specifically <sup>13</sup> C/ <sup>2</sup> H - labeled nucleosides from [ <sup>13</sup> C 6 ]- d -glucose. <i>Tetrahedron Letters</i> , 1998, 39, 2793-2796.	1.4	15
94	Synthesis of [ <sup>2</sup> H 1 ]-nucleosides with defined ( <sup>5</sup> S)/( <sup>5</sup> R) - ratios. <i>Tetrahedron Letters</i> , 1998, 39, 2873-2876.	1.4	7
95	Determination of peptide $\tau$ angles in solids by relayed anisotropy correlation NMR. <i>Solid State Nuclear Magnetic Resonance</i> , 1998, 11, 169-175.	2.3	32
96	NMR structure of the Streptomyces metalloproteinase inhibitor, SMPI, isolated from Streptomyces nigrescens TK-23: another example of an ancestral <sup>12</sup> [ <sup>3</sup> -crystallin precursor structure 1 1 Edited by P. E. Wright. <i>Journal of Molecular Biology</i> , 1998, 282, 421-433.	4.2	39
97	Collision-Induced Dissociation Spectra Obtained by Fourier Transform Ion Cyclotron Resonance Mass Spectrometry Using a <sup>13</sup> C, <sup>15</sup> N-Doubly Depleted Protein. <i>Analytical Chemistry</i> , 1998, 70, 3333-3336.	6.5	10
98	Measurement of Deoxyribose <sup>3</sup> JHHS scalar Couplings Reveals Protein Binding-Induced Changes in the Sugar Puckers of the DNA. <i>Journal of the American Chemical Society</i> , 1998, 120, 821-822.	13.7	18
99	Elucidation of the mode of interaction of thermolysin with a proteinaceous metalloproteinase inhibitor, SMPI, based on a model complex structure and a structural dynamics analysis 1 1 Edited by P. E. Wright. <i>Journal of Molecular Biology</i> , 1998, 282, 435-446.	4.2	19
100	Measurement of <sup>3</sup> JC2â€P scalar Couplings in a 17 kDa Protein Complex with <sup>13</sup> C, <sup>15</sup> N-Labeled DNA Distinguishes the Bland BIIPhosphate Conformations of the DNA. <i>Journal of the American Chemical Society</i> , 1997, 119, 9901-9902.	13.7	33
101	Synthesis of ( <sup>5</sup> S)-[ <sup>5</sup> - <sup>2</sup> H1;1â€2,2â€3â€4â€5â€- <sup>13</sup> C5]-Thymidine via stereoselective deuteration of a 5-oxoribose derivative. <i>Tetrahedron Letters</i> , 1997, 38, 395-398.	1.4	14
102	Novel approach to diastereoselective synthesis of 2'-deoxy[5'- <sup>2</sup> H1]ribonucleoside derivatives by reduction of the corresponding 5'-O-acetyl-2'-deoxy-5'-phenylselenoribonucleoside derivatives with a Bu <sub>3</sub> Sn <sup>2</sup> H-Et <sub>3</sub> B system. <i>Chirality</i> , 1997, 9, 435-442.	2.6	14
103	<sup>5</sup> Methylene Proton Signal Assignment of DNA/RNA Oligomers Labeled with <sup>5</sup> -Monodeuterated Nucleosides by <sup>1</sup> H- <sup>31</sup> P HSQC Spectroscopy. <i>Magnetic Resonance in Chemistry</i> , 1996, 34, S40-S46.	1.9	22
104	Relayed anisotropy correlation NMR: determination of dihedral angles in solids. <i>Chemical Physics Letters</i> , 1996, 256, 133-140.	2.6	100
105	Motion of Scandium Ions in Sc <sub>2</sub> C <sub>8</sub> O <sub>4</sub> Observed by <sup>45</sup> Sc Solution NMR. <i>The Journal of Physical Chemistry</i> , 1996, 100, 9579-9581.	2.9	61
106	Biosynthesis of Lactacystin... <i>Journal of Antibiotics</i> , 1995, 48, 1015-1020.	2.0	20
107	Sequence-Specific DNA Recognition of the Escherichia coli Ada Protein Associated with the Methylation-Dependent Functional Switch for Transcriptional Regulation. <i>Journal of Biochemistry</i> , 1995, 118, 1184-1191.	1.7	9
108	Significance of the Highly Conserved Gly-4 Residue in Human Cystatin A1. <i>Journal of Biochemistry</i> , 1995, 118, 635-642.	1.7	14

#	ARTICLE	IF	CITATIONS
109	Novel synthesis of 2'-deoxy[5'- <sup>2</sup> H]ribonucleoside derivatives from 5'-O-Ac-2'-deoxy-5'-PhSe-ribonucleoside derivatives. <i>Tetrahedron Letters</i> , 1995, 36, 6699-6700.	1.4	14
110	Highly Diastereoselective Synthesis of (2'S)-[2'- <sup>2</sup> H]-2'-Deoxyribonucleosides from the Corresponding Ribonucleosides. <i>Nucleosides, Nucleotides and Nucleic Acids</i> , 1995, 14, 333-336.	1.1	13
111	Solution structure of a human cystatin A variant, cystatin A2-98 M65L by NMR spectroscopy. A possible role of the interactions between the N- and C-termini to maintain the inhibitory active form of cystatin A. <i>Biochemistry</i> , 1995, 34, 14637-14648.	2.5	43
112	Stereospecific measurements of the vicinal <sup>1</sup> H- <sup>31</sup> P coupling constants for the diastereotopic C5' methylene protons in a DNA dodecamer with a <sup>13</sup> C/ <sup>2</sup> H doubly labeled residue. Conformational analysis of the torsion angle $\beta$ . <i>Journal of the American Chemical Society</i> , 1995, 117, 7277-7278.	13.7	24
113	Synthesis of L-threo- and L-erythro-[1- <sup>13</sup> C, 2,3- <sup>2</sup> H <sub>2</sub> ]amino acids: novel probes for conformational analysis of peptide side chains. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1995, , 1603-1609.	0.9	22
114	Sonochemical and Triethylborane-Induced Tin Deuteride Reduction for the Highly Diastereoselective Synthesis of (2'R)-2'-Deoxy[2'- <sup>2</sup> H]ribonucleoside Derivatives. <i>Journal of Organic Chemistry</i> , 1995, 60, 6980-6986.	3.2	39
115	Preparation and heteronuclear 2D NMR spectroscopy of a DNA dodecamer containing a thymidine residue with a uniformly <sup>13</sup> C-labeled deoxyribose ring. <i>Journal of Biomolecular NMR</i> , 1994, 4, 581-586.	2.8	39
116	Biosynthesis of lactacystin. Origin of the carbons and stereospecific NMR assignment of the two diastereotopic methyl groups. <i>Tetrahedron Letters</i> , 1994, 35, 5009-5012.	1.4	20
117	Synthesis of phenylalanines regiospecifically labelled with deuterium in the aromatic ring. <i>Journal of Labelled Compounds and Radiopharmaceuticals</i> , 1994, 34, 831-837.	1.0	11
118	Methylation Dependent Functional Switch Mechanism Newly Found in the Escherichia coli Ada Protein. <i>Journal of the American Chemical Society</i> , 1994, 116, 6035-6036.	13.7	39
119	An Alternative Triple-Resonance Method for the Through-Bond Correlation of Intranucleotide H1' and H8 NMR Signals of Purine Nucleotides. Application to a DNA Dodecamer with Fully <sup>13</sup> C/ <sup>15</sup> N-Labeled Deoxyadenosine Residues. <i>Journal of the American Chemical Society</i> , 1994, 116, 5977-5978.	13.7	48
120	Sonochemical and triethylborane-induced tin deuteride reduction for the highly stereoselective synthesis of (2â€²R)-[2â€²- <sup>2</sup> H]-2â€²-deoxyribonucleosides from 2â€²-functionalized ribonucleosides. <i>Tetrahedron Letters</i> , 1993, 34, 1317-1320.	1.4	27
121	Localisation of methionine residues in bacteriorhodopsin by carbonyl <sup>13</sup> C-NMR with sequence-specific assignments. <i>FEBS Letters</i> , 1993, 327, 7-12.	2.8	15
122	NMR characterization of isomers of C78, C82 and C84 fullerenes. <i>Nature</i> , 1992, 357, 142-145.	27.8	519
123	Synthesis of [1,1â€²- <sup>13</sup> C <sub>2</sub> ]-L-cystine. <i>Journal of Labelled Compounds and Radiopharmaceuticals</i> , 1991, 29, 867-874.	1.0	5
124	Reductive cleavage and regeneration of the disulfide bonds in Streptomyces subtilisin inhibitor (SSI) as studied by the carbonyl <sup>13</sup> C NMR resonances of cysteinyl residues. <i>Journal of Biomolecular NMR</i> , 1991, 1, 49-64.	2.8	9
125	Application of <sup>13</sup> C Nuclear Magnetic Resonance Spectroscopy to Molecular Structural Analyses of Antibody Molecules. <i>Journal of Biochemistry</i> , 1989, 105, 867-869.	1.7	34
126	Internal motion of a tryptophan residue in Streptomyces subtilisin inhibitor: Deuterium nuclear magnetic resonance in solution. <i>Proteins: Structure, Function and Bioinformatics</i> , 1988, 4, 131-136.	2.6	8



#	ARTICLE	IF	CITATIONS
127	Correlation of carbon-13 and nitrogen-15 chemical shifts in selectively and uniformly labeled proteins by heteronuclear two-dimensional NMR spectroscopy. <i>Journal of the American Chemical Society</i> , 1988, 110, 6256-6258.	13.7	104
128	Local structural features around the C-terminal segment of <i>Streptomyces subtilisin</i> inhibitor studied by the carbonyl carbon nuclear magnetic resonances of three phenylalanyl residues. <i>Biochemistry</i> , 1987, 26, 1068-1075.	2.5	42
129	Assignment of the three methionyl carbonyl carbon resonances in <i>Streptomyces subtilisin</i> inhibitor by a carbon-13 and nitrogen-15 double-labeling technique. A new strategy for structural studies of proteins in solution. <i>Biochemistry</i> , 1982, 21, 6273-6279.	2.5	174
130	Dihedral-angle dependence of the vicinal $^{15}\text{N}$ , $^{13}\text{C}$ spin-coupling constants. A new NMR parameter for the conformational analysis of amino acids and peptides. <i>Magnetic Resonance in Chemistry</i> , 1981, 17, 46-49.	0.7	3
131	Detection of Proton-acceptor Sites of Hydrogen Bonding in Adenine-Uracil Base Pairs by the Use of $^{15}\text{N}$ Magnetic Resonance. <i>FEBS Journal</i> , 1981, 117, 553-558.	0.2	18
132	Caution in using nitrogen-15-carbon-13 spin-spin coupling for determining (bio)synthetic pathways. <i>Journal of the American Chemical Society</i> , 1979, 101, 1031-1032.	13.7	12
133	ASSIGNMENT OF $^1\text{H}$ -PROTON RESONANCES OF L-HISTIDINE BY STEREOSELECTIVE DEUTERIUM SUBSTITUTION. <i>Chemistry Letters</i> , 1979, 8, 395-396.	1.3	4
134	A NEW STRATEGY OF FOOD ANALYSIS USING LIQUID CHROMATOGRAPHY AND $^{13}\text{C}$ NMR SPECTROSCOPY. , 1979, , 59-80.		0
135	Carbon-13 nuclear magnetic resonance spectra of gross plant tissues containing starch. <i>Tetrahedron Letters</i> , 1978, 19, 1563-1566.	1.4	11
136	Ion permeation across the bilayer of annealed phosphatidylcholine vesicles at elevated temperatures. Concentration dependence and the micelle-bilayer dynamic equilibrium. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1977, 468, 411-422.	2.6	15
137	In situ analysis of the microbial fermentation process by natural abundance $^{13}\text{C}$ and $^{31}\text{P}$ NMR spectroscopy. Production of adenosine-5'-triphosphate from adenosine. <i>FEBS Letters</i> , 1977, 80, 385-389.	2.8	21
138	The formation and annealing of structural defects in lipid bilayer vesicles. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1976, 443, 313-330.	2.6	112
139	Proton magnetic resonance studies of lipid bilayer membranes Experimental determination of inter- and intramolecular nuclear relaxation rates in sonicated phosphatidylcholine bilayer vesicles. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1976, 433, 282-293.	2.6	57
140	The formation and annealing of structural defects in lipid bilayer vesicles. <i>Nucleic Acids and Protein Synthesis</i> , 1976, 443, 313-330.	1.7	42
141	$^{13}\text{C}$ nuclear magnetic resonance spectrum of dried fruits and its histological implications. <i>Tetrahedron Letters</i> , 1976, 17, 4757-4760.	1.4	9
142	$^{13}\text{C}$ NMR studies of the intact plant tissues. Cytoplasmic aucubin and sucrose in a single seed of <i>aucuba japonica</i> . <i>Tetrahedron Letters</i> , 1976, 17, 4279-4282.	1.4	7
143	Thermal phase transitions in deuterated lecithin bilayers. <i>Chemistry and Physics of Lipids</i> , 1975, 14, 343-349.	3.2	58
144	State of molecular motion of cholesterol in lecithin bilayers. <i>Nature</i> , 1975, 256, 582-584.	27.8	69

#	ARTICLE	IF	CITATIONS
145	Effects of structural defects in sonicated phospholipid vesicles on fusion and ion permeability. <i>Nature</i> , 1975, 256, 584-586.	27.8	61
146	Conformational analysis of amino acids and peptides using specific isotope substitution. II. Conformation of serine, tyrosine, phenylalanine, aspartic acid, asparagine, and aspartic acid .beta.-methyl ester in various ionization states. <i>Journal of the American Chemical Society</i> , 1975, 97, 5630-5631.	13.7	84
147	Diastereomeric interaction of partially resolved amines facilitated by lanthanide chelates. Evidence for dynamic equilibrium between seven-coordinate and eight-coordinate alkylamine-lanthanide chelate adducts. <i>Journal of the American Chemical Society</i> , 1975, 97, 1761-1765.	13.7	14
148	Tris(6,6,7,7,8,8,8-heptafluoro-2,2-dimethyl-3,5-octamedionato)gadolinium (Gd(fod) <sub>3</sub> )-induced contact shifts. Versatile new method to estimate contact and pseudocontact shift contributions to observed lanthanide-induced shifts. <i>Journal of the American Chemical Society</i> , 1975, 97, 330-334.	13.7	46
149	Conformational study of cyclic nucleotides. Lanthanide ion assisted analysis of the hydrogen-1 nuclear magnetic resonance spectra. <i>Journal of the American Chemical Society</i> , 1975, 97, 6839-6843.	13.7	39
150	Conformational analysis of amino acids and peptides using specific isotope substitution. I. Conformation of L-phenylalanyl-glycine. <i>Biochemical and Biophysical Research Communications</i> , 1975, 64, 425-432.	2.1	27
151	Evidence for the presence of contact term contribution to lanthanide induced shifts in <sup>1</sup> H and <sup>13</sup> C NMR spectra of pyridine N-oxides. <i>Tetrahedron Letters</i> , 1973, 14, 1573-1576.	1.4	13
152	Importance of complex formation and contact shifts in the application of lanthanide shift reagents to <sup>1</sup> H and <sup>13</sup> C nmr spectra of aromatic compounds. <i>Tetrahedron Letters</i> , 1973, 14, 3127-3130.	1.4	16
153	EVIDENCE FOR THE PRESENCE OF CONTACT TERM CONTRIBUTION TO LANTHANIDE-INDUCED ISOTROPIC SHIFTS IN <sup>13</sup> C AND <sup>19</sup> F NMR SPECTRA OF ALIPHATIC COMPOUNDS: CAUTION FOR APPLICATIONS OF LANTHANIDE SHIFT REAGENTS. <i>Chemistry Letters</i> , 1972, 1, 1061-1064.	1.3	11
154	Use of chiral solvents or lanthanide shift reagents to distinguish meso from d or l diastereomers. <i>Journal of the American Chemical Society</i> , 1972, 94, 5924-5926.	13.7	65
155	High-resolution proton and phosphorus nuclear magnetic resonance spectra of flavine adenine dinucleotide and its conformation in aqueous solution. <i>Biochemistry</i> , 1972, 11, 741-752.	2.5	86
156	Studies of Phosphorylation. V. The Synthesis of Inosine-5- <sup>32</sup> P-thiophosphates. <i>Bulletin of the Chemical Society of Japan</i> , 1971, 44, 460-463.	3.2	10
157	Conformation of $\hat{\pm}$ -Aminobutyric Acid in Aqueous Solution. <i>Bulletin of the Chemical Society of Japan</i> , 1971, 44, 2577-2582.	3.2	4
158	The Crystal Structure of 5,5-Dimethyl-2-oxo-2-hydroxy-1,3,2-dioxaphosphorinane Monohydrate. <i>Bulletin of the Chemical Society of Japan</i> , 1969, 42, 1819-1824.	3.2	19
159	Phosphorus-proton Spin-spin Coupling in the P=O-C-H Group. A Comparison of Cyclic and Acyclic Systems. <i>Bulletin of the Chemical Society of Japan</i> , 1969, 42, 1713-1718.	3.2	42
160	Conformational Dependence of P=O Stretching Vibration Frequency in Six-membered Cyclic Phosphates. <i>Bulletin of the Chemical Society of Japan</i> , 1969, 42, 845-845.	3.2	26
161	Conformational changes in the sulfur analogs of 11- and 12-membered metacyclophanes. <i>Tetrahedron Letters</i> , 1968, 9, 4185-4189.	1.4	27
162	Medium-Sized Cyclophanes. VI. High-Temperature NMR Spectra of [2.2]Metacyclophane and Optical Resolution of 4,14-Disubstituted Derivatives. <i>Bulletin of the Chemical Society of Japan</i> , 1968, 41, 218-221.	3.2	26

#	ARTICLE	IF	CITATIONS
163	Medium-sized Cyclophanes. II. The Stereoselective Synthesis and Optical Resolution of 4, 14-Dimethyl[2.2]metacyclophane. Bulletin of the Chemical Society of Japan, 1966, 39, 856-856.	3.2	36