

# Ronald Bottlender

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10844392/publications.pdf>

Version: 2024-02-01

47  
papers

5,283  
citations

109321

35  
h-index

197818

49  
g-index

49  
all docs

49  
docs citations

49  
times ranked

5933  
citing authors

#	ARTICLE	IF	CITATIONS
1	Predictors of treatment response to psychological interventions in people at clinical high risk of first-episode psychosis. <i>Microbial Biotechnology</i> , 2019, 13, 120-127.	1.7	5
2	Coping as a predictor of treatment outcome in people at clinical high risk of psychosis. <i>Microbial Biotechnology</i> , 2016, 10, 17-27.	1.7	19
3	Accelerated Brain Aging in Schizophrenia and Beyond: A Neuroanatomical Marker of Psychiatric Disorders. <i>Schizophrenia Bulletin</i> , 2014, 40, 1140-1153.	4.3	369
4	Association between psychopathology and problems of psychosocial functioning in the long-term outcome of patients diagnosed with schizophrenic, schizoaffective and affective disorders. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2013, 263, 85-92.	3.2	20
5	Early Recognition and Disease Prediction in the At-Risk Mental States for Psychosis Using Neurocognitive Pattern Classification. <i>Schizophrenia Bulletin</i> , 2012, 38, 1200-1215.	4.3	121
6	Disease Prediction in the At-Risk Mental State for Psychosis Using Neuroanatomical Biomarkers: Results From the FePsy Study. <i>Schizophrenia Bulletin</i> , 2012, 38, 1234-1246.	4.3	139
7	Multivariate patterns of brain-cognition associations relating to vulnerability and clinical outcome in the at-risk mental states for psychosis. <i>Human Brain Mapping</i> , 2012, 33, 2104-2124.	3.6	23
8	Differences in hippocampal volume between major depression and schizophrenia: a comparative neuroimaging study. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010, 260, 127-137.	3.2	53
9	The Munich 15-year follow-up study (MUFUSSAD) on first-hospitalized patients with schizophrenic or affective disorders: comparison of psychopathological and psychosocial course and outcome and prediction of chronicity. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010, 260, 367-384.	3.2	56
10	Social disability in schizophrenic, schizoaffective and affective disorders 15years after first admission. <i>Schizophrenia Research</i> , 2010, 116, 9-15.	2.0	73
11	Neuroanatomical correlates of executive dysfunction in the at-risk mental state for psychosis. <i>Schizophrenia Research</i> , 2010, 123, 160-174.	2.0	46
12	Use of neuroanatomical pattern regression to predict the structural brain dynamics of vulnerability and transition to psychosis. <i>Schizophrenia Research</i> , 2010, 123, 175-187.	2.0	58
13	Neuroanatomical correlates of different vulnerability states for psychosis and their clinical outcomes. <i>British Journal of Psychiatry</i> , 2009, 195, 218-226.	2.8	85
14	Prediction of symptom remission in schizophrenia during inpatient treatment. <i>World Journal of Biological Psychiatry</i> , 2009, 10, 426-434.	2.6	18
15	Use of Neuroanatomical Pattern Classification to Identify Subjects in At-Risk Mental States of Psychosis and Predict Disease Transition. <i>Archives of General Psychiatry</i> , 2009, 66, 700.	12.3	382
16	Anterior cingulate cortex does not differ between patients with major depression and healthy controls, but relatively large anterior cingulate cortex predicts a good clinical course. <i>Psychiatry Research - Neuroimaging</i> , 2008, 163, 76-83.	1.8	49
17	Depression during an acute episode of schizophrenia or schizophreniform disorder and its impact on treatment response. <i>Psychiatry Research</i> , 2008, 158, 297-305.	3.3	24
18	Structural correlates of psychopathological symptom dimensions in schizophrenia: A voxel-based morphometric study. <i>NeuroImage</i> , 2008, 39, 1600-1612.	4.2	166

#	ARTICLE	IF	CITATIONS
19	Depression-Related Variation in Brain Morphology Over 3 Years. Archives of General Psychiatry, 2008, 65, 1156.	12.3	329
20	Effect of hippocampal and amygdala volumes on clinical outcomes in major depression: a 3-year prospective magnetic resonance imaging study. Journal of Psychiatry and Neuroscience, 2008, 33, 423-30.	2.4	211
21	Association of the Brain-Derived Neurotrophic Factor Val66Met Polymorphism With Reduced Hippocampal Volumes in Major Depression. Archives of General Psychiatry, 2007, 64, 410.	12.3	357
22	Randomized controlled multicentre trial of cognitive behaviour therapy in the early initial prodromal state: effects on social adjustment post treatment. Microbial Biotechnology, 2007, 1, 71-78.	1.7	54
23	Psychopathological characteristics and treatment response of first episode compared with multiple episode schizophrenic disorders. European Archives of Psychiatry and Clinical Neuroscience, 2007, 257, 47-53.	3.2	58
24	Reduced hippocampal volume correlates with executive dysfunctioning in major depression. Journal of Psychiatry and Neuroscience, 2006, 31, 316-23.	2.4	178
25	Interventions in the initial prodromal states of psychosis in Germany: concept and recruitment. British Journal of Psychiatry, 2005, 187, s45-s48.	2.8	57
26	Fifteen-year follow-up of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition depressive disorders: the prognostic significance of psychotic features. Comprehensive Psychiatry, 2005, 46, 322-327.	3.1	16
27	Classification of Functional Psychoses and Its Implication for Prognosis: Comparison between ICD-10 and DSM-IV. Psychopathology, 2004, 37, 110-117.	1.5	30
28	Mixed depressive features predict maniform switch during treatment of depression in bipolar I disorder. Journal of Affective Disorders, 2004, 78, 149-152.	4.1	79
29	Early detection and secondary prevention of psychosis: facts and visions*. European Archives of Psychiatry and Clinical Neuroscience, 2004, 254, 117-128.	3.2	183
30	Pharmacological long-term treatment strategies in first episode schizophrenia. European Archives of Psychiatry and Clinical Neuroscience, 2004, 254, 129-140.	3.2	52
31	Psychopathology of early-onset versus late-onset schizophrenia revisited: an observation of 473 neuroleptic-naive patients before and after first-admission treatments. Schizophrenia Research, 2004, 67, 175-183.	2.0	59
32	Reduced Hippocampal Volumes Associated With the Long Variant of the Serotonin Transporter Polymorphism in Major Depression. Archives of General Psychiatry, 2004, 61, 177.	12.3	164
33	Hippocampal and Amygdala Changes in Patients With Major Depressive Disorder and Healthy Controls During a 1-Year Follow-Up. Journal of Clinical Psychiatry, 2004, 65, 492-499.	2.2	188
34	Larger amygdala volumes in first depressive episode as compared to recurrent major depression and healthy control subjects. Biological Psychiatry, 2003, 53, 338-344.	1.3	238
35	The impact of the duration of untreated psychosis prior to first psychiatric admission on the 15-year outcome in schizophrenia. Schizophrenia Research, 2003, 62, 37-44.	2.0	127
36	The impact of the duration of untreated psychosis on short- and long-term outcome in schizophrenia. Current Opinion in Psychiatry, 2003, 16, S39-S43.	6.3	18

#	ARTICLE	IF	CITATIONS
37	On the Descriptive Validity of ICD-10 Schizophrenia: Empirical Analyses in the Spectrum of Non-Affective Functional Psychoses. <i>Psychopathology</i> , 2003, 36, 152-159.	1.5	36
38	Negative Symptoms in Depressed and Schizophrenic Patients. <i>Journal of Clinical Psychiatry</i> , 2003, 64, 954-958.	2.2	39
39	Hippocampal Changes in Patients With a First Episode of Major Depression. <i>American Journal of Psychiatry</i> , 2002, 159, 1112-1118.	7.2	464
40	Enlargement of the amygdala in patients with a first episode of major depression. <i>Biological Psychiatry</i> , 2002, 51, 708-714.	1.3	285
41	The Kraepelinian dichotomy: preliminary results of a 15-year follow-up study on functional psychoses: focus on negative symptoms. <i>Schizophrenia Research</i> , 2002, 56, 87-94.	2.0	70
42	The impact of duration of untreated psychosis and premorbid functioning on outcome of first inpatient treatment in schizophrenic and schizoaffective patients. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2002, 252, 226-231.	3.2	39
43	Deficit states in schizophrenia and their association with the length of illness and gender. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2001, 251, 272-278.	3.2	19
44	Mood-stabilisers reduce the risk of developing antidepressant-induced maniform states in acute treatment of bipolar I depressed patients. <i>Journal of Affective Disorders</i> , 2001, 63, 79-83.	4.1	91
45	Long-term course of schizophrenic, affective and schizoaffective psychosis: focus on negative symptoms and their impact on global indicators of outcome. <i>Acta Psychiatrica Scandinavica</i> , 2000, 102, 54-57.	4.5	52
46	Dipole localization of P300 and normal aging. <i>Brain Topography</i> , 2000, 13, 3-9.	1.8	19
47	Impact of duration of symptoms prior to first hospitalization on acute outcome in 998 schizophrenic patients. <i>Schizophrenia Research</i> , 2000, 44, 145-150.	2.0	50