

Peter Van Der Geer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10849114/publications.pdf>

Version: 2024-02-01

32
papers

6,741
citations

331670

21
h-index

414414

32
g-index

33
all docs

33
docs citations

33
times ranked

5736
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of STS-1 as a novel ShcA-binding protein. <i>Biochemical and Biophysical Research Communications</i> , 2017, 490, 1334-1339.	2.1	11
2	Identification and mutagenesis of the TACE and β -secretase cleavage sites in the colony-stimulating factor 1 receptor. <i>Biochemical and Biophysical Research Communications</i> , 2014, 450, 782-787.	2.1	15
3	Analysis of Protein-Protein Interactions by Coimmunoprecipitation. <i>Methods in Enzymology</i> , 2014, 541, 35-47.	1.0	9
4	Wolfram Syndrome protein, Miner1, regulates sulphhydryl redox status, the unfolded protein response, and Ca^{2+} homeostasis. <i>EMBO Molecular Medicine</i> , 2013, 5, 904-918.	6.9	101
5	Engineering NGF receptors to bind Grb2 directly uncovers differences in signaling ability between Grb2 and ShcA binding sites. <i>FEBS Letters</i> , 2012, 586, 3658-3664.	2.8	9
6	Interactions of the NPXY microdomains of the low density lipoprotein receptor-related protein 1. <i>Proteomics</i> , 2009, 9, 5016-5028.	2.2	59
7	Structural and Functional Consequences of Tyrosine Phosphorylation in the LRP1 Cytoplasmic Domain. <i>Journal of Biological Chemistry</i> , 2008, 283, 15656-15664.	3.4	37
8	MitoNEET is an iron-containing outer mitochondrial membrane protein that regulates oxidative capacity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 5318-5323.	7.1	251
9	The Outer Mitochondrial Membrane Protein mitoNEET Contains a Novel Redox-active 2Fe-2S Cluster*. <i>Journal of Biological Chemistry</i> , 2007, 282, 23745-23749.	3.4	145
10	Purification and Identification of Protein-Tyrosine Kinase-binding Proteins Using Synthetic Phosphopeptides as Affinity Reagents. <i>Molecular and Cellular Proteomics</i> , 2004, 3, 887-895.	3.8	5
11	v-Src induces Shc binding to tyrosine 63 in the cytoplasmic domain of the LDL receptor-related protein 1. <i>Oncogene</i> , 2003, 22, 3589-3597.	5.9	57
12	Multiple Regions of Internalin B Contribute to Its Ability to Turn on the Ras-Mitogen-activated Protein Kinase Pathway. <i>Journal of Biological Chemistry</i> , 2003, 278, 7783-7789.	3.4	43
13	Phosphorylation of LRP1 Regulation of Transport and Signal Transduction. <i>Trends in Cardiovascular Medicine</i> , 2002, 12, 160-165.	4.9	56
14	C-Cbl binds the CSF-1 receptor at tyrosine 973, a novel phosphorylation site in the receptor's carboxy-terminus. <i>Oncogene</i> , 2002, 21, 1079-1089.	5.9	73
15	Tyrosine-phosphorylated Low Density Lipoprotein Receptor-related Protein 1 (LRP1) Associates with the Adaptor Protein SHC in SRC-transformed Cells. <i>Journal of Biological Chemistry</i> , 2001, 276, 19119-19125.	3.4	108
16	RasGAP-Associated Endoribonuclease G3BP: Selective RNA Degradation and Phosphorylation-Dependent Localization. <i>Molecular and Cellular Biology</i> , 2001, 21, 7747-7760.	2.3	167
17	The Shc adaptor protein forms interdependent phosphotyrosine-mediated protein complexes in mast cells stimulated with interleukin 3. <i>Blood</i> , 2000, 96, 132-138.	1.4	20
18	The Shc adaptor protein forms interdependent phosphotyrosine-mediated protein complexes in mast cells stimulated with interleukin 3. <i>Blood</i> , 2000, 96, 132-138.	1.4	1

#	ARTICLE	IF	CITATIONS
19	Re-engineering the target specificity of the insulin receptor by modification of a PTB domain binding site. <i>Oncogene</i> , 1999, 18, 3071-3075.	5.9	13
20	Phosphorylation of Rat Serine 105 or Mouse Threonine 217 in C/EBP β Is Required for Hepatocyte Proliferation Induced by TGF β . <i>Molecular Cell</i> , 1999, 4, 1087-1092.	9.7	170
21	Phosphopeptide Mapping and Identification of Phosphorylation Sites. <i>Current Protocols in Molecular Biology</i> , 1999, 48, Unit 18.9.	2.9	4
22	Phosphopeptide Mapping and Identification of Phosphorylation Sites. <i>Current Protocols in Protein Science</i> , 1999, 18, Unit13.9.	2.8	7
23	The Shc adaptor protein is highly phosphorylated at conserved, twin tyrosine residues (Y239/240) that mediate protein-protein interactions. <i>Current Biology</i> , 1996, 6, 1435-1444.	3.9	207
24	Characterization of the Phosphotyrosine-binding Domain of the Drosophila Shc Protein. <i>Journal of Biological Chemistry</i> , 1996, 271, 31855-31862.	3.4	16
25	A conserved amino-terminal Shc domain binds to phosphotyrosine motifs in activated receptors and phosphopeptides. <i>Current Biology</i> , 1995, 5, 404-412.	3.9	173
26	The Receptor-like Protein-tyrosine Phosphatase, RPTP β , Is Phosphorylated by Protein Kinase C on Two Serines Close to the Inner Face of the Plasma Membrane. <i>Journal of Biological Chemistry</i> , 1995, 270, 10587-10594.	3.4	70
27	The PTB domain: a new protein module implicated in signal transduction. <i>Trends in Biochemical Sciences</i> , 1995, 20, 277-280.	7.5	228
28	Integrin-mediated signal transduction linked to Ras pathway by GRB2 binding to focal adhesion kinase. <i>Nature</i> , 1994, 372, 786-791.	27.8	1,528
29	Phosphopeptide mapping and phosphoamino acid analysis by electrophoresis and chromatography on thin-layer cellulose plates. <i>Electrophoresis</i> , 1994, 15, 544-554.	2.4	136
30	Receptor Protein-Tyrosine Kinases and Their Signal Transduction Pathways. <i>Annual Review of Cell Biology</i> , 1994, 10, 251-337.	26.1	1,194
31	Transactivation by NF-IL6/LAP is enhanced by phosphorylation of its activation domain. <i>Nature</i> , 1993, 364, 544-547.	27.8	343
32	[11] Phosphopeptide mapping and phosphoamino acid analysis by two-dimensional separation on thin-layer cellulose plates. <i>Methods in Enzymology</i> , 1991, 201, 110-149.	1.0	1,485