

Javier F Botto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10859491/publications.pdf>

Version: 2024-02-01

30
papers

2,165
citations

279798

23
h-index

454955

30
g-index

30
all docs

30
docs citations

30
times ranked

2336
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Ultra-High-Density QTL Marker Mapping for Seedling Photomorphogenesis Mediating Arabidopsis Establishment in Southern Patagonia. <i>Frontiers in Plant Science</i> , 2021, 12, 677728. | 3.6 | 3 |
| 2 | BBX21 reduces abscisic acid sensitivity, mesophyll conductance and chloroplast electron transport capacity to increase photosynthesis and water use efficiency in potato plants cultivated under moderated drought. <i>Plant Journal</i> , 2021, 108, 1131-1144. | 5.7 | 11 |
| 3 | Heterologous Expression of <i>AtBBX21</i> Enhances the Rate of Photosynthesis and Alleviates Photoinhibition in <i>Solanum tuberosum</i> . <i>Plant Physiology</i> , 2018, 177, 369-380. | 4.8 | 27 |
| 4 | SALT OVERLY SENSITIVE 2 (SOS2) and Interacting Partners SOS3 and ABSCISIC ACID-INSENSITIVE 2 (ABI2) Promote Red-Light-Dependent Germination and Seedling Deetiolation in <i>Arabidopsis</i> . <i>International Journal of Plant Sciences</i> , 2017, 178, 485-493. | 1.3 | 16 |
| 5 | A single haplotype hyposensitive to light and requiring strong vernalization dominates <i>Arabidopsis thaliana</i> populations in Patagonia, Argentina. <i>Molecular Ecology</i> , 2017, 26, 3389-3404. | 3.9 | 11 |
| 6 | Molecular mechanisms underlying the entrance in secondary dormancy of <i>Arabidopsis</i> seeds. <i>Plant, Cell and Environment</i> , 2016, 39, 213-221. | 5.7 | 34 |
| 7 | The Multifaceted Roles of HY5 in Plant Growth and Development. <i>Molecular Plant</i> , 2016, 9, 1353-1365. | 8.3 | 465 |
| 8 | The transcriptional regulator BBX24 impairs DELLA activity to promote shade avoidance in <i>Arabidopsis thaliana</i> . <i>Nature Communications</i> , 2015, 6, 6202. | 12.8 | 96 |
| 9 | Plasticity to simulated shade is associated with altitude in structured populations of <i>Arabidopsis thaliana</i> . <i>Plant, Cell and Environment</i> , 2015, 38, 1321-1332. | 5.7 | 19 |
| 10 | Physiological responses of spring rapeseed (<i>Brassica napus</i>) to red/far-red ratios and irradiance during pre- and post-flowering stages. <i>Physiologia Plantarum</i> , 2014, 152, 784-794. | 5.2 | 24 |
| 11 | The BBX family of plant transcription factors. <i>Trends in Plant Science</i> , 2014, 19, 460-470. | 8.8 | 370 |
| 12 | BBX proteins in green plants: Insights into their evolution, structure, feature and functional diversification. <i>Gene</i> , 2013, 531, 44-52. | 2.2 | 122 |
| 13 | The receptor-like kinase ERECTA contributes to the shade-avoidance syndrome in a background-dependent manner. <i>Annals of Botany</i> , 2013, 111, 811-819. | 2.9 | 38 |
| 14 | Transcriptional Programs Related to Phytochrome A Function in Arabidopsis Seed Germination. <i>Molecular Plant</i> , 2013, 6, 1261-1273. | 8.3 | 34 |
| 15 | Molecular interactions of BBX24 and BBX25 with HYH, HY5 HOMOLOG, to modulate <i>Arabidopsis</i> seedling development. <i>Plant Signaling and Behavior</i> , 2013, 8, e25208. | 2.4 | 52 |
| 16 | The <i>Arabidopsis</i> B-BOX Protein BBX25 Interacts with HY5, Negatively Regulating <i>BBX22</i> Expression to Suppress Seedling Photomorphogenesis. <i>Plant Cell</i> , 2013, 25, 1243-1257. | 6.6 | 189 |
| 17 | Function of B-BOX under shade. <i>Plant Signaling and Behavior</i> , 2011, 6, 101-104. | 2.4 | 41 |
| 18 | Genetic mapping of natural variation in a shade avoidance response: ELF3 is the candidate gene for a QTL in hypocotyl growth regulation. <i>Journal of Experimental Botany</i> , 2011, 62, 167-176. | 4.8 | 40 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | AtBBX21 and COP1 genetically interact in the regulation of shade avoidance. <i>Plant Journal</i> , 2010, 64, 551-562. | 5.7 | 92 |
| 20 | Germination variation in <i>Arabidopsis thaliana</i> accessions under moderate osmotic and salt stresses. <i>Annals of Botany</i> , 2010, 106, 833-842. | 2.9 | 71 |
| 21 | The Heterotrimeric Gâ€protein Complex Modulates Light Sensitivity in <i>Arabidopsis thaliana</i> Seed Germination. <i>Photochemistry and Photobiology</i> , 2009, 85, 949-954. | 2.5 | 23 |
| 22 | Manipulation of Light Environment to Produce High-quality Poinsettia Plants. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2009, 44, 702-706. | 1.0 | 26 |
| 23 | Light-related Loci Controlling Seed Germination in Ler Ã— Cvi and Bay-0 Ã— Sha Recombinant Inbred-line Populations of <i>Arabidopsis thaliana</i> . <i>Annals of Botany</i> , 2008, 102, 631-642. | 2.9 | 30 |
| 24 | Seasonal and plantâ€density dependency for quantitative trait loci affecting flowering time in multiple populations of <i>Arabidopsis thaliana</i> . <i>Plant, Cell and Environment</i> , 2007, 30, 1465-1479. | 5.7 | 31 |
| 25 | New <i>Arabidopsis</i> Recombinant Inbred Lines (<i>Landsberg erecta</i> Ã— Nossen) Reveal Natural Variation in Phytochrome-Mediated Responses. <i>Plant Physiology</i> , 2005, 138, 1126-1135. | 4.8 | 20 |
| 26 | Mapping Quantitative Trait Loci in Multiple Populations of <i>Arabidopsis thaliana</i> Identifies Natural Allelic Variation for Trichome Density. <i>Genetics</i> , 2005, 169, 1649-1658. | 2.9 | 85 |
| 27 | The Cape Verde Islands Allele of Cryptochrome 2 Enhances Cotyledon Unfolding in the Absence of Blue Light in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2003, 133, 1547-1556. | 4.8 | 46 |
| 28 | Burial conditions affect light responses of <i>Datura ferox</i> seeds. <i>Seed Science Research</i> , 1998, 8, 423-429. | 1.7 | 40 |
| 29 | The effect of light during and after soil cultivation with different tillage implements on weed seedling emergence. <i>Weed Science</i> , 1998, 46, 351-357. | 1.5 | 52 |
| 30 | Role of Phytochrome B in the Induction of Seed Germination by Light in <i>Arabidopsis thaliana</i> . <i>Journal of Plant Physiology</i> , 1995, 146, 307-312. | 3.5 | 57 |