

# Josefine Roswall

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1086185/publications.pdf>

Version: 2024-02-01

16  
papers

2,717  
citations

1040056

9  
h-index

996975

15  
g-index

16  
all docs

16  
docs citations

16  
times ranked

4717  
citing authors

#	ARTICLE	IF	CITATIONS
1	Developmental trajectory of the healthy human gut microbiota during the first 5 years of life. <i>Cell Host and Microbe</i> , 2021, 29, 765-776.e3.	11.0	208
2	Serum osteocalcin levels at four months of age were associated with neurodevelopment at four years of age in term-born children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2021, , .	1.5	2
3	Nutrition- and feeding practice-related risk factors for rapid weight gain during the first year of life: a population-based birth cohort study. <i>BMC Pediatrics</i> , 2020, 20, 507.	1.7	10
4	A prospective study of perinatal and metabolic risk factors for early childhood caries. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2020, 109, 2356-2361.	1.5	11
5	MRI estimates of brown adipose tissue in children – Associations to adiposity, osteocalcin, and thigh muscle volume. <i>Magnetic Resonance Imaging</i> , 2019, 58, 135-142.	1.8	8
6	Infant body mass index growth patterns predicted overweight at five years, waist-to-height ratio did not add to this predictivity. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2019, 108, 945-953.	1.5	0
7	Longitudinal birth cohort study found that a significant proportion of children had abnormal metabolic profiles and insulin resistance at 6 years of age. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2019, 108, 486-492.	1.5	10
8	Body mass index classification misses to identify children with an elevated waist-to-height ratio at 5 years of age. <i>Pediatric Research</i> , 2019, 85, 30-35.	2.3	6
9	Parents with overweight children two and five years of age did not perceive them as weighing too much. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2018, 107, 1060-1064.	1.5	10
10	Serum $\omega$ 6 and $\omega$ 9 Fatty Acids Correlate With Serum IGF1 and Growth Up to 4 Months of Age in Healthy Infants. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2018, 66, 141-146.	1.8	9
11	Influence of mode of delivery, family and nursing determinants on early childhood caries development: a prospective cohort study. <i>Acta Odontologica Scandinavica</i> , 2018, 76, 595-599.	1.6	20
12	Overweight at four years of age in a Swedish birth cohort: influence of neighbourhood-level purchasing power. <i>BMC Public Health</i> , 2016, 16, 546.	2.9	13
13	Salivary microflora and mode of delivery: a prospective case control study. <i>BMC Oral Health</i> , 2015, 15, 155.	2.3	12
14	Dynamics and Stabilization of the Human Gut Microbiome during the First Year of Life. <i>Cell Host and Microbe</i> , 2015, 17, 690-703.	11.0	2,276
15	Influence of neighbourhood purchasing power on breastfeeding at four months of age: a Swedish population-based cohort study. <i>BMC Public Health</i> , 2013, 13, 1077.	2.9	8
16	Factors associated with discontinuation of breastfeeding before 12 month of age. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2012, 101, 55-60.	1.5	114