Maarit Piirtola

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1086667/publications.pdf

Version: 2024-02-01

23 papers 1,401 citations

16 h-index 23 g-index

24 all docs

24 docs citations

times ranked

24

2086 citing authors

#	Article	IF	CITATIONS
1	Force Platform Measurements as Predictors of Falls among Older People – A Review. Gerontology, 2006, 52, 1-16.	1.4	538
2	Physical therapy approaches to reduce fall and fracture risk among older adults. Nature Reviews Endocrinology, 2010, 6, 396-407.	4.3	139
3	Factors contributing to psychological distress in the working population, with a special reference to gender difference. BMC Public Health, 2021, 21, 611.	1.2	123
4	Effectiveness of the Chaos Falls Clinic in preventing falls and injuries of home-dwelling older adults: A randomised controlled trial. Injury, 2014, 45, 265-271.	0.7	78
5	Fractures as predictors of excess mortality in the aged—A population-based study with a 12-year follow-up. European Journal of Epidemiology, 2008, 23, 747-755.	2.5	70
6	The Older Finnish Twin Cohort â€" 45 Years of Follow-up. Twin Research and Human Genetics, 2019, 22, 240-254.	0.3	68
7	A multifactorial fall prevention programme in the community-dwelling aged: predictors of adherence. European Journal of Public Health, 2007, 17, 464-470.	0.1	64
8	Association of current and former smoking with body mass index: A study of smoking discordant twin pairs from 21 twin cohorts. PLoS ONE, 2018, 13, e0200140.	1.1	57
9	Effects of risk-based multifactorial fall prevention on postural balance in the community-dwelling aged: A randomized controlled trial. Archives of Gerontology and Geriatrics, 2009, 48, 22-27.	1.4	32
10	Psychotropic drugs and the risk of fractures in old age: a prospective population-based study. BMC Public Health, 2010, 10, 396.	1.2	31
11	Correlates and genetics of selfâ€reported sleep and awake bruxism in a nationwide twin cohort. Journal of Oral Rehabilitation, 2020, 47, 1110-1119.	1.3	31
12	Opioids, antiepileptic and anticholinergic drugs and the risk of fractures in patients 65 years of age and older: a prospective population-based study. Age and Ageing, 2013, 42, 318-324.	0.7	26
13	Leisure-time physical inactivity and association with body mass index: a Finnish Twin Study with a 35-year follow-up. International Journal of Epidemiology, 2017, 46, 116-127.	0.9	26
14	Association between education and future leisure-time physical inactivity: a study of Finnish twins over a 35-year follow-up. BMC Public Health, 2016, 16, 720.	1.2	17
15	Incidence of fractures and changes over time among the aged in a Finnish municipality: a population-based 12-year follow-up. Aging Clinical and Experimental Research, 2007, 19, 269-276.	1.4	16
16	Predictors of fractures among the aged: a population-based study with 12-year follow-up in a Finnish municipality. Aging Clinical and Experimental Research, 2008, 20, 242-252.	1,4	16
17	A Study of Sedentary Behaviour in the Older Finnish Twin Cohort: A Cross Sectional Analysis. BioMed Research International, 2014, 2014, 1-9.	0.9	15
18	Effects of risk-based multifactorial fall prevention program on maximal isometric muscle strength in community-dwelling aged: a randomized controlled trial. Aging Clinical and Experimental Research, 2008, 20, 487-493.	1.4	13

#	Article	IF	CITATIONS
19	The associations of smoking dependence motives with depression among daily smokers. Addiction, 2021, 116, 2162-2174.	1.7	13
20	Fractures as an Independent Predictor of Functional Decline in Older People: A Population-Based Study with an 8-Year Follow-Up. Gerontology, 2012, 58, 296-304.	1.4	9
21	Association between long-term smoking and leisure-time physical inactivity: a cohort study among Finnish twins with a 35-year follow-up. International Journal of Public Health, 2017, 62, 819-829.	1.0	7
22	Associations of sitting time with leisureâ€time physical inactivity, education, and body mass index change. Scandinavian Journal of Medicine and Science in Sports, 2020, 30, 322-331.	1.3	7
23	Life course associations between smoking and depressive symptoms. A 30-year Finnish follow-up study. Nordic Journal of Psychiatry, 2022, , 1-1.	0.7	0