

Stefano Maini

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1087357/publications.pdf>

Version: 2024-02-01

22
papers

900
citations

687363

13
h-index

713466

21
g-index

22
all docs

22
docs citations

22
times ranked

932
citing authors

#	ARTICLE	IF	CITATIONS
1	Interactions between the Multicolored Asian Lady Beetle <i>Harmonia axyridis</i> and the Parasitoid <i>Dinocampus coccinellae</i> . <i>Insects</i> , 2016, 7, 67.	2.2	11
2	Pre-wintering conditions and post-winter performance in a solitary bee: does diapause impose an energetic cost on reproductive success?. <i>Ecological Entomology</i> , 2016, 41, 201-210.	2.2	44
3	<i>Macrocyclus albidus</i> (Copepoda: cyclopidae) for the Biocontrol of <i>Aedes albopictus</i> and <i>Culex pipiens</i> in Italy. <i>Journal of the American Mosquito Control Association</i> , 2015, 31, 32-43.	0.7	20
4	Effects of mechanical distribution on survival and reproduction of <i>Phytoseiulus persimilis</i> and <i>Amblyseius swirskii</i> . <i>Biosystems Engineering</i> , 2015, 129, 11-19.	4.3	7
5	Duration of prepupal summer dormancy regulates synchronization of adult diapause with winter temperatures in bees of the genus <i>Osmia</i> . <i>Journal of Insect Physiology</i> , 2012, 58, 924-933.	2.0	53
6	The long summer: Pre-wintering temperatures affect metabolic expenditure and winter survival in a solitary bee. <i>Journal of Insect Physiology</i> , 2011, 57, 1651-1659.	2.0	96
7	Dispersal and Survival of <i>Aedes albopictus</i> (Diptera: Culicidae) Males in Italian Urban Areas and Significance for Sterile Insect Technique Application. <i>Journal of Medical Entomology</i> , 2010, 47, 1082-1091.	1.8	66
8	<i>Apis mellifera</i> and <i>Osmia cornuta</i> as carriers for the secondary spread of <i>Bacillus subtilis</i> on apple flowers. <i>BioControl</i> , 2009, 54, 123-133.	2.0	24
9	Estimation of mortality by entomophages on exotic <i>Harmonia axyridis</i> versus native <i>Adalia bipunctata</i> in semi-field conditions in northern Italy. <i>BioControl</i> , 2008, 53, 277-287.	2.0	14
10	Occurrence of <i>Bacillus thuringiensis</i> harboring insecticidal cry1 genes in a corn field in Northern Italy. <i>Agronomy for Sustainable Development</i> , 2008, 28, 473-480.	5.3	3
11	Tolerance Thresholds for <i>Aedes albopictus</i> and <i>Aedes caspius</i> in Italian Urban Areas. <i>Journal of the American Mosquito Control Association</i> , 2008, 24, 377-386.	0.7	39
12	Estimation of mortality by entomophages on exotic <i>Harmonia axyridis</i> versus native <i>Adalia bipunctata</i> in semi-field conditions in northern Italy. , 2007, , 277-287.		1
13	Assessing delayed and acute toxicity of five formulated fungicides to <i>Osmia lignaria</i> Say and <i>Apis mellifera</i> . <i>Apidologie</i> , 2005, 36, 449-460.	2.0	105
14	Evaluation of a standard artificial flower design to feed individual bees known amounts of pesticides. <i>Apidologie</i> , 2005, 36, 379-387.	2.0	11
15	On the Competition Occurring Between <i>Aedes albopictus</i> and <i>Culex pipiens</i> (Diptera: Culicidae) in Italy. <i>Environmental Entomology</i> , 2003, 32, 1313-1321.	1.4	103
16	A method to feed individual bees (Hymenoptera: Apiformes) known amounts of pesticides. <i>Apidologie</i> , 2003, 34, 597-602.	2.0	28
17	On intra-guild predation and cannibalism in <i>Harmonia axyridis</i> (Pallas) and <i>Adalia bipunctata</i> L. (Coleoptera: Coccinellidae). <i>Biological Control</i> , 2002, 24, 110-116.	3.0	121
18	A new risk of occupational disease: allergic asthma and rhinoconjunctivitis in persons working with beneficial arthropods. <i>International Archives of Occupational and Environmental Health</i> , 1996, 68, 133-135.	2.3	10

#	ARTICLE	IF	CITATIONS
19	A new risk of occupational disease: allergic asthma and rhinoconjunctivitis in persons working with beneficial arthropods. <i>International Archives of Occupational and Environmental Health</i> , 1996, 68, 133-135.	2.3	1
20	A new risk of occupational disease: allergic asthma and rhinoconjunctivitis in persons working with beneficial arthropods. <i>International Archives of Occupational and Environmental Health</i> , 1994, 65, 291-294.	2.3	13
21	Attraction of male turnip moths <i>Agrotis segetum</i> (Lepidoptera: Noctuidae) to sex pheromone components and their mixtures at 11 sites in Europe, Asia, and Africa. <i>Journal of Chemical Ecology</i> , 1992, 18, 1337-1347.	1.8	77
22	Sex pheromone of European corn borer.. <i>Journal of Chemical Ecology</i> , 1988, 14, 1359-1366.	1.8	53