
Sandra Tenreiro

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1087578/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Guidelines for the use and interpretation of assays for monitoring autophagy (4th) Tj ETQq1 1 0.784314 rgBT /Ov	verlock 10 9.1	Tf 50 742 T 1,430
2	The YEASTRACT database: a tool for the analysis of transcription regulatory associations in Saccharomyces cerevisiae. Nucleic Acids Research, 2006, 34, D446-D451.	14.5	421
3	Protein phosphorylation in neurodegeneration: friend or foe?. Frontiers in Molecular Neuroscience, 2014, 7, 42.	2.9	203
4	YEASTRACT-DISCOVERER: new tools to improve the analysis of transcriptional regulatory associations in Saccharomyces cerevisiae. Nucleic Acids Research, 2007, 36, D132-D136.	14.5	140
5	DJ-1 interactions with α-synuclein attenuate aggregation and cellular toxicity in models of Parkinson's disease. Cell Death and Disease, 2014, 5, e1350-e1350.	6.3	130
6	Phosphorylation Modulates Clearance of Alpha-Synuclein Inclusions in a Yeast Model of Parkinson's Disease. PLoS Genetics, 2014, 10, e1004302.	3.5	114
7	SNCA (α-synuclein)-induced toxicity in yeast cells is dependent on Sir2-mediated mitophagy. Autophagy, 2012, 8, 1494-1509.	9.1	113
8	Expression of theAZR1 gene (ORF YGR224w), encoding a plasma membrane transporter of the major facilitator superfamily, is required for adaptation to acetic acid and resistance to azoles inSaccharomyces cerevisiae. Yeast, 2000, 16, 1469-1481.	1.7	91
9	Harnessing the power of yeast to unravel the molecular basis of neurodegeneration. Journal of Neurochemistry, 2013, 127, 438-452.	3.9	82
10	FLR1 gene (ORF YBR008c) is required for benomyl and methotrexate resistance inSaccharomyces cerevisiae and its benomyl-induced expression is dependent on Pdr3 transcriptional regulator. Yeast, 1999, 15, 1595-1608.	1.7	78
11	Simple is good: yeast models of neurodegeneration. FEMS Yeast Research, 2010, 10, 970-979.	2.3	77
12	Saccharomyces cerevisiae Aqr1 Is an Internal-Membrane Transporter Involved in Excretion of Amino Acids. Eukaryotic Cell, 2004, 3, 1492-1503.	3.4	76
13	Dtr1p, a Multidrug Resistance Transporter of the Major Facilitator Superfamily, Plays an Essential Role in Spore Wall Maturation in Saccharomyces cerevisiae. Eukaryotic Cell, 2002, 1, 799-810.	3.4	74
14	Visualization of cell-to-cell transmission of mutant huntingtin oligomers. PLOS Currents, 2011, 3, RRN1210.	1.4	74
15	AQR1 Gene (ORF YNL065w) Encodes a Plasma Membrane Transporter of the Major Facilitator Superfamily That Confers Resistance to Short-Chain Monocarboxylic Acids and Quinidine in Saccharomyces cerevisiae. Biochemical and Biophysical Research Communications, 2002, 292, 741-748.	2.1	73
16	(Poly)phenols protect from α-synuclein toxicity by reducing oxidative stress and promoting autophagy. Human Molecular Genetics, 2015, 24, 1717-1732.	2.9	66
17	The multidrug resistance transporters of the major facilitator superfamily, 6 years after disclosure of Saccharomyces cerevisiae genome sequence. Journal of Biotechnology, 2002, 98, 215-226.	3.8	65
18	From the baker to the bedside: yeast models of Parkinson's disease. Microbial Cell, 2015, 2, 262-279.	3.2	59

2

SANDRA TENREIRO

#	Article	IF	CITATIONS
19	Thermus silvanus sp. nov. and Thermus chliarophilus sp. nov., Two New Species Related to Thermus ruber but with Lower Growth Temperatures. International Journal of Systematic Bacteriology, 1995, 45, 633-639.	2.8	56
20	Parkinson Disease Mutant E46K Enhances α-Synuclein Phosphorylation in Mammalian Cell Lines, in Yeast, and in Vivo. Journal of Biological Chemistry, 2015, 290, 9412-9427.	3.4	52
21	Yeast models of Parkinson's disease-associated molecular pathologies. Current Opinion in Genetics and Development, 2017, 44, 74-83.	3.3	49
22	Saccharomyces cerevisiae Multidrug Resistance Transporter Qdr2 Is Implicated in Potassium Uptake, Providing a Physiological Advantage to Quinidine-Stressed Cells. Eukaryotic Cell, 2007, 6, 134-142.	3.4	48
23	Age-Related Macular Degeneration: Pathophysiology, Management, and Future Perspectives. Ophthalmologica, 2021, 244, 495-511.	1.9	48
24	Thermonema rossianum sp. nov., a New Thermophilic and Slightly Halophilic Species from Saline Hot Springs in Naples, Italy. International Journal of Systematic Bacteriology, 1997, 47, 122-126.	2.8	46
25	Transfer of extracellular vesicleâ€micro <scp>RNA</scp> controls germinal center reaction and antibody production. EMBO Reports, 2020, 21, e48925.	4.5	46
26	Saccharomyces cerevisiae Multidrug Transporter Qdr2p (Yil121wp): Localization and Function as a Quinidine Resistance Determinant. Antimicrobial Agents and Chemotherapy, 2004, 48, 2531-2537.	3.2	45
27	The yeast multidrug transporter Qdr3 (Ybr043c): localization and role as a determinant of resistance to quinidine, barban, cisplatin, and bleomycin. Biochemical and Biophysical Research Communications, 2005, 327, 952-959.	2.1	43
28	Inhibition of formation of α-synuclein inclusions by mannosylglycerate in a yeast model of Parkinson's disease. Biochimica Et Biophysica Acta - General Subjects, 2013, 1830, 4065-4072.	2.4	43
29	Transcriptional Activation of FLR1 Gene during Saccharomyces cerevisiae Adaptation to Growth with Benomyl: Role of Yap1p and Pdr3p. Biochemical and Biophysical Research Communications, 2001, 280, 216-222.	2.1	40
30	Resistance and Adaptation to Quinidine in Saccharomyces cerevisiae : Role of QDR1 (YIL120w), Encoding a Plasma Membrane Transporter of the Major Facilitator Superfamily Required for Multidrug Resistance. Antimicrobial Agents and Chemotherapy, 2001, 45, 1528-1534.	3.2	40
31	Adaptive response to the antimalarial drug artesunate in yeast involves Pdr1p/Pdr3p-mediated transcriptional activation of the resistance determinantsTPO1andPDR5. FEMS Yeast Research, 2006, 6, 1130-1139.	2.3	38
32	Fatty Composition of the Species of the Genera Thermus Acid Meiothermus. Systematic and Applied Microbiology, 1996, 19, 303-311.	2.8	37
33	PLK2 Modulates α-Synuclein Aggregation in Yeast and Mammalian Cells. Molecular Neurobiology, 2013, 48, 854-862.	4.0	37
34	Yeast response and tolerance to polyamine toxicity involving the drug : H+ antiporter Qdr3 and the transcription factors Yap1 and Gcn4. Microbiology (United Kingdom), 2011, 157, 945-956.	1.8	36
35	The effects of the novel A53E alpha-synuclein mutation on its oligomerization and aggregation. Acta Neuropathologica Communications, 2016, 4, 128.	5.2	35
36	Modulation of alpha-synuclein toxicity in yeast using a novel microfluidic-based gradient generator. Lab on A Chip, 2014, 14, 3949-3957.	6.0	33

SANDRA TENREIRO

#	Article	IF	CITATIONS
37	Yeast reveals similar molecular mechanisms underlying alpha- and beta-synuclein toxicity. Human Molecular Genetics, 2016, 25, 275-290.	2.9	29
38	Polar lipids and fatty acid composition ofThermus strains from New Zealand. Antonie Van Leeuwenhoek, 1994, 66, 357-363.	1.7	21
39	Transcriptomic Profiling of the <i>Saccharomyces cerevisiae</i> Response to Quinine Reveals a Glucose Limitation Response Attributable to Drug-Induced Inhibition of Glucose Uptake. Antimicrobial Agents and Chemotherapy, 2009, 53, 5213-5223.	3.2	21
40	(Poly)phenol-digested metabolites modulate alpha-synuclein toxicity by regulating proteostasis. Scientific Reports, 2018, 8, 6965.	3.3	20
41	DNA:DNA hybridization and chemotaxonomic studies of Thermus scotoductus. Research in Microbiology, 1995, 146, 315-324.	2.1	19
42	High-throughput study of alpha-synuclein expression in yeast using microfluidics for control of local cellular microenvironment. Biomicrofluidics, 2012, 6, 014109.	2.4	11
43	Macular Vascular Imaging and Connectivity Analysis Using High-Resolution Optical Coherence Tomography. Translational Vision Science and Technology, 2022, 11, 2.	2.2	10
44	Phycocyanin protects against Alpha-Synuclein toxicity in yeast. Journal of Functional Foods, 2017, 38, 553-560.	3.4	9
45	Retinal Progression Biomarkers of Early and Intermediate Age-Related Macular Degeneration. Life, 2022, 12, 36.	2.4	9
46	Neuroprotection or Neurotoxicity of Illicit Drugs on Parkinson's Disease. Life, 2020, 10, 86.	2.4	8
47	Formation of Lipofuscin-Like Autofluorescent Granules in the Retinal Pigment Epithelium Requires Lysosome Dysfunction. , 2021, 62, 39.		6
48	Impaired Proteostasis Contributes to Renal Tubular Dysgenesis. PLoS ONE, 2011, 6, e20854.	2.5	6
49	Identification of novel protein phosphatases as modifiers of alpha-synuclein aggregation in yeast. FEMS Yeast Research, 2018, 18, .	2.3	4
50	The synthetic cannabinoid JWH-018 modulates Saccharomyces cerevisiae energetic metabolism. FEMS Yeast Research, 2019, 19, .	2.3	2
51	CORRELATION STUDY BETWEEN DRUSEN MORPHOLOGY AND FUNDUS AUTOFLUORESCENCE. Retina, 2021, 41, 555-562.	1.7	2
52	Choroidal Vascular Impairment in Intermediate Age-Related Macular Degeneration. Diagnostics, 2022, 12, 1290.	2.6	2
53	Analysis of Protein Oligomeric Species by Sucrose Gradients. Methods in Molecular Biology, 2016, 1449, 331-339.	0.9	1
54	A biophysical perspective on the unexplored mechanisms driving Parkinson's disease by amphetamine-like stimulants. Neural Regeneration Research, 2021, 16, 2213.	3.0	1

#	Article	IF	CITATIONS
55	A levedura como modelo para estudar as bases moleculares da doença de Parkinson. Revista Brasileira De Ciências Do Envelhecimento Humano, 2015, 12, .	0.0	1
56	Identification of targets and mechanisms of resistance to imatinib and quinine using a molecular systems biology approach. , 2011, , .		0
57	Integration of Single Cell Traps, Chemical Gradient Generator and Photosensors in a Microfluidic Platform for the Study of Alpha-Synuclein Toxicity in Yeast. Procedia Engineering, 2014, 87, 92-95.	1.2	0
58	FLR1 gene (ORF YBR008c) is required for benomyl and methotrexate resistance in Saccharomyces cerevisiae and its benomylâ€induced expression is dependent on Pdr3 transcriptional regulator. Yeast, 1999, 15, 1595-1608.	1.7	0