

Brigitte Kasper

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10922217/publications.pdf>

Version: 2024-02-01

13
papers

1,454
citations

759233

12
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

1574
citing authors

#	ARTICLE	IF	CITATIONS
1	Syntaxin 11 is required for NK and CD8 ⁺ T cell cytotoxicity and neutrophil degranulation. <i>European Journal of Immunology</i> , 2013, 43, 194-208.	2.9	57
2	Molecular pathways of platelet factor 4/CXCL4 signaling. <i>European Journal of Cell Biology</i> , 2011, 90, 521-526.	3.6	60
3	CXCL4-induced monocyte survival, cytokine expression, and oxygen radical formation is regulated by sphingosine kinase 1. <i>European Journal of Immunology</i> , 2010, 40, 1162-1173.	2.9	17
4	Fc γ RIIA and Fc γ RIIIB Are Required for Autoantibody-Induced Tissue Damage in Experimental Human Models of Bullous Pemphigoid. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2841-2844.	0.7	48
5	Familial Hemophagocytic Lymphohistiocytosis Type 5 (FHL-5) Is Caused by Mutations in Munc18-2 and Impaired Binding to Syntaxin 11. <i>American Journal of Human Genetics</i> , 2009, 85, 482-492.	6.2	370
6	Platelet Factor 4 (CXC Chemokine Ligand 4) Differentially Regulates Respiratory Burst, Survival, and Cytokine Expression of Human Monocytes by Using Distinct Signaling Pathways. <i>Journal of Immunology</i> , 2007, 179, 2584-2591.	0.8	61
7	Neutrophil adhesion to endothelial cells induced by platelet factor 4 requires sequential activation of Ras, Syk, and JNK MAP kinases. <i>Blood</i> , 2006, 107, 1768-1775.	1.4	43
8	Linkage of familial hemophagocytic lymphohistiocytosis (FHL) type-4 to chromosome 6q24 and identification of mutations in syntaxin 11. <i>Human Molecular Genetics</i> , 2005, 14, 827-834.	2.9	502
9	Platelet Factor 4/CXCL4 Induces Phagocytosis and the Generation of Reactive Oxygen Metabolites in Mononuclear Phagocytes Independently of Gi Protein Activation or Intracellular Calcium Transients. <i>Journal of Immunology</i> , 2004, 173, 2060-2067.	0.8	92
10	Platelet factor 4 (PF-4)-induced neutrophil adhesion is controlled by src-kinases, whereas PF-4-mediated exocytosis requires the additional activation of p38 MAP kinase and phosphatidylinositol 3-kinase. <i>Blood</i> , 2004, 103, 1602-1610.	1.4	62
11	Platelet Factor 4 Inhibits Proliferation and Cytokine Release of Activated Human T Cells. <i>Journal of Immunology</i> , 2002, 169, 770-777.	0.8	100
12	Differential expression and regulation of GTPases (RhoA and Rac2) and GDIs (LyGDI and RhoGDI) in neutrophils from patients with severe congenital neutropenia. <i>Blood</i> , 2000, 95, 2947-2953.	1.4	32
13	Cytosolic proteins from neutrophilic granulocytes: A comparison between patients with severe chronic neutropenia and healthy donors. <i>Electrophoresis</i> , 1997, 18, 142-149.	2.4	10