
## Hugo Vanderstichele

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10949982/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The Alzheimer's Association international guidelines for handling of cerebrospinal fluid for routine clinical measurements of amyloid $\hat{l}^2$ and tau. Alzheimer's and Dementia, 2021, 17, 1575-1582.	0.8	51
2	Cerebrospinal fluid hemoglobin levels as markers of blood contamination: relevance for α-synuclein measurement. Clinical Chemistry and Laboratory Medicine, 2021, 59, 1653-1661.	2.3	2
3	First amyloid β1â€42 certified reference material for reâ€calibrating commercial immunoassays. Alzheimer's and Dementia, 2020, 16, 1493-1503.	0.8	42
4	Antibodyâ€based methods for the measurement of αâ€synuclein concentration in human cerebrospinal fluid – method comparison and round robin study. Journal of Neurochemistry, 2019, 149, 126-138.	3.9	44
5	Cerebrospinal Fluid Total and Phosphorylated α-Synuclein in Patients with Creutzfeldt–Jakob Disease and Synucleinopathy. Molecular Neurobiology, 2019, 56, 3476-3483.	4.0	26
6	Ultrasensitive Detection of Plasma Amyloid-β as a Biomarker for Cognitively Normal Elderly Individuals at Risk of Alzheimer's Disease. Journal of Alzheimer's Disease, 2019, 71, 775-783.	2.6	38
7	Cerebrospinal fluid levels of synaptic and neuronal integrity correlate with gray matter volume and amyloid load in the precuneus of cognitively intact older adults. Journal of Neurochemistry, 2019, 149, 139-157.	3.9	10
8	Synaptic biomarkers in CSF aid in diagnosis, correlate with cognition and predict progression in MCI and Alzheimer's disease. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2019, 5, 871-882.	3.7	79
9	APPâ€derived peptides reflect neurodegeneration in frontotemporal dementia. Annals of Clinical and Translational Neurology, 2019, 6, 2518-2530.	3.7	13
10	Automation on an Open-Access Platform of Alzheimer's Disease Biomarker Immunoassays. SLAS Technology, 2018, 23, 188-197.	1.9	5
11	Plasma Al̂² (Amyloid-l̂²) Levels and Severity and Progression of Small Vessel Disease. Stroke, 2018, 49, 884-890.	2.0	27
12	C-Reactive Protein, Plasma Amyloid-β Levels, and Their Interaction With Magnetic Resonance Imaging Markers. Stroke, 2018, 49, 2692-2698.	2.0	46
13	Commutability of the certified reference materials for the standardization of β-amyloid 1-42 assay in human cerebrospinal fluid: lessons for tau and β-amyloid 1-40 measurements. Clinical Chemistry and Laboratory Medicine, 2018, 56, 2058-2066.	2.3	27
14	The impact of preanalytical variables on measuring cerebrospinal fluid biomarkers for Alzheimer's disease diagnosis: A review. Alzheimer's and Dementia, 2018, 14, 1313-1333.	0.8	87
15	Plasma amyloid-β levels, cerebral atrophy and risk of dementia: a population-based study. Alzheimer's Research and Therapy, 2018, 10, 63.	6.2	39
16	Relevance of Aβ42/40 Ratio for Detection of Alzheimer Disease Pathology in Clinical Routine: The PLMR Scale. Frontiers in Aging Neuroscience, 2018, 10, 138.	3.4	59
17	CSF Aβ1–42 – an excellent but complicated Alzheimer's biomarker – a route to standardisation. Clinica Chimica Acta, 2017, 467, 27-33.	1.1	104
18	How to handle adsorption of cerebrospinal fluid amyloid $\hat{I}^2$ (1-42) in laboratory practice? Identifying		52

problematic handlings and resolving the issue by use of the  $\hat{Al^242}$  / $\hat{Al^240}$  ratio. , 2017, 13, 885-892.

#	Article	IF	CITATIONS
19	Accelerating drug development for Alzheimer's disease through the use of data standards. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2017, 3, 273-283.	3.7	10
20	Plasma Amyloid-β Levels, Cerebral Small Vessel Disease, and Cognition: The Rotterdam Study. Journal of Alzheimer's Disease, 2017, 60, 977-987.	2.6	43
21	A user's guide for αâ€synuclein biomarker studies in biological fluids: Perianalytical considerations. Movement Disorders, 2017, 32, 1117-1130.	3.9	54
22	Concordance Between Cerebrospinal Fluid Biomarkers with Alzheimer's Disease Pathology Between Three Independent Assay Platforms. Journal of Alzheimer's Disease, 2017, 61, 169-183.	2.6	21
23	Recommendations for cerebrospinal fluid collection for the analysis by ELISA of neurogranin trunc P75, α-synuclein, and total tau in combination with Al²(1–42)/Al²(1–40). Alzheimer's Research and Therapy, 2017, 9, 40.	6.2	17
24	Bloodâ€based biomarkers in Alzheimer disease: Current state of the science and a novel collaborative paradigm for advancing from discovery to clinic. Alzheimer's and Dementia, 2017, 13, 45-58.	0.8	227
25	Prevention of tau increase in cerebrospinal fluid of APP transgenic mice suggests downstream effect of BACE1 inhibition. Alzheimer's and Dementia, 2017, 13, 701-709.	0.8	35
26	Differential role of CSF fatty acid binding protein 3, α-synuclein, and Alzheimer's disease core biomarkers in Lewy body disorders and Alzheimer's dementia. Alzheimer's Research and Therapy, 2017, 9, 52.	6.2	101
27	Diagnostic Impact of Cerebrospinal Fluid Biomarker (Pre-)Analytical Variability in Alzheimer's Disease. Journal of Alzheimer's Disease, 2016, 51, 97-106.	2.6	20
28	Cerebrospinal Fluid Biomarkers for Alzheimer's Disease: A View of the Regulatory Science Qualification Landscape from the Coalition Against Major Diseases CSF Biomarker Team. Journal of Alzheimer's Disease, 2016, 55, 19-35.	2.6	35
29	Performance Evaluation of an Automated ELISA System for Alzheimer's Disease Detection in Clinical Routine. Journal of Alzheimer's Disease, 2016, 54, 55-67.	2.6	27
30	No diurnal variation of classical and candidate biomarkers of Alzheimer's disease in CSF. Molecular Neurodegeneration, 2016, 11, 65.	10.8	16
31	Prospective longitudinal course of cognition in older subjects with mild parkinsonian signs. Alzheimer's Research and Therapy, 2016, 8, 42.	6.2	14
32	<scp>CSF</scp> A <i>β</i> 42/A <i>β</i> 40 and A <i>β</i> 42/A <i>β</i> 38 ratios: better diagnostic markers of Alzheimer disease. Annals of Clinical and Translational Neurology, 2016, 3, 154-165.	3.7	329
33	Assessing the commutability of reference material formats for the harmonization of amyloid-Î <sup>2</sup> measurements. Clinical Chemistry and Laboratory Medicine, 2016, 54, 1177-1191.	2.3	49
34	A First Tetraplex Assay for the Simultaneous Quantification of Total α-Synuclein, Tau, β-Amyloid42 and DJ-1 in Human Cerebrospinal Fluid. PLoS ONE, 2016, 11, e0153564.	2.5	6
35	O4-11-06: The confidence level of established cut-off values for CSF Alzheimer's disease-specific biomarkers. , 2015, 11, P298-P298.		2
36	The Alzheimer's Disease Neuroimaging Initiative 2 Biomarker Core: A review of progress and plans. Alzheimer's and Dementia, 2015, 11, 772-791.	0.8	79

#	Article	IF	CITATIONS
37	Alzheimer's disease cerebrospinal fluid biomarker in cognitively normal subjects. Brain, 2015, 138, 2701-2715.	7.6	109
38	Cerebrospinal fluid biomarkers in trials for Alzheimer and Parkinson diseases. Nature Reviews Neurology, 2015, 11, 41-55.	10.1	144
39	Roadblocks for integration of novel biomarker concepts into clinical routine: the peptoid approach. Alzheimer's Research and Therapy, 2014, 6, 23.	6.2	3
40	Changes in plasma amyloid beta in a longitudinal study of aging and Alzheimer's disease. Alzheimer's and Dementia, 2014, 10, 53-61.	0.8	114
41	CSF biomarker variability in the Alzheimer's Association quality control program. Alzheimer's and Dementia, 2013, 9, 251-261.	0.8	344
42	Validation of Assays for Measurement of Amyloid-Î <sup>2</sup> Peptides in Cerebrospinal Fluid and Plasma Specimens from Patients with Alzheimer's Disease Treated with Solanezumab. Journal of Alzheimer's Disease, 2013, 34, 897-910.	2.6	17
43	Cerebrospinal Fluid Collection Tubes: A Critical Issue for Alzheimer Disease Diagnosis. Clinical Chemistry, 2012, 58, 787-789.	3.2	50
44	Validation of a Multiplex Assay for Simultaneous Quantification of Amyloid-β Peptide Species in Human Plasma with Utility for Measurements in Studies of Alzheimer's Disease Therapeutics. Journal of Alzheimer's Disease, 2012, 32, 905-918.	2.6	29
45	Comparison of Two Analytical Platforms for the Clinical Qualification of Alzheimer's Disease Biomarkers in Pathologically-Confirmed Dementia. Journal of Alzheimer's Disease, 2012, 33, 117-131.	2.6	40
46	Standardization of preanalytical aspects of cerebrospinal fluid biomarker testing for Alzheimer's disease diagnosis: A consensus paper from the Alzheimer's Biomarkers Standardization Initiative. Alzheimer's and Dementia, 2012, 8, 65-73.	0.8	271
47	Improved protocol for measurement of plasma β-amyloid in longitudinal evaluation of Alzheimer's Disease Neuroimaging Initiative study patients. , 2012, 8, 250-260.		56
48	Simultaneous analysis of cerebrospinal fluid biomarkers using microsphere-based xMAP multiplex technology for early detection of Alzheimer's disease. Methods, 2012, 56, 484-493.	3.8	85
49	Risk of Alzheimer's Disease Biological Misdiagnosis Linked to Cerebrospinal Collection Tubes. Journal of Alzheimer's Disease, 2012, 31, 13-20.	2.6	94
50	Potential sources of interference on Abeta immunoassays in biological samples. Alzheimer's Research and Therapy, 2012, 4, 39.	6.2	14
51	Reference measurement procedures for Alzheimer's disease cerebrospinal fluid biomarkers: definitions and approaches with focus on amyloid β42. Biomarkers in Medicine, 2012, 6, 409-417.	1.4	76
52	Analytical aspects of molecular Alzheimer's disease biomarkers. Biomarkers in Medicine, 2012, 6, 377-389.	1.4	26
53	The Alzheimer's Association external quality control program for cerebrospinal fluid biomarkers. Alzheimer's and Dementia, 2011, 7, 386.	0.8	354
54	Validation of ELISA Methods for Quantification of Total Tau and Phosphorylated-Tau181 in Human Cerebrospinal Fluid with Measurement in Specimens from Two Alzheimer's Disease Studies. Journal of Alzheimer's Disease, 2011, 26, 531-541.	2.6	23

#	Article	IF	CITATIONS
55	Qualification of the analytical and clinical performance of CSF biomarker analyses in ADNI. Acta Neuropathologica, 2011, 121, 597-609.	7.7	256
56	Factors affecting Aβ plasma levels and their utility as biomarkers in ADNI. Acta Neuropathologica, 2011, 122, 401-13.	7.7	151
57	Comparison of Analytical Platforms for Cerebrospinal Fluid Measures of β-Amyloid 1-42, Total tau, and P-tau <sub>181</sub> for Identifying Alzheimer Disease Amyloid Plaque Pathology. Archives of Neurology, 2011, 68, 1137.	4.5	161
58	Diagnosis-Independent Alzheimer Disease Biomarker Signature in Cognitively Normal Elderly People. Archives of Neurology, 2010, 67, 949.	4.5	407
59	Evaluation of plasma Aβ40 and Aβ42 as predictors of conversion to Alzheimer's disease in patients with mild cognitive impairment. Neurobiology of Aging, 2010, 31, 357-367.	3.1	242
60	Added diagnostic value of CSF biomarkers in differential dementia diagnosis. Neurobiology of Aging, 2010, 31, 1867-1876.	3.1	63
61	Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. Annals of Neurology, 2009, 65, 403-413.	5.3	1,803
62	Increased total-Tau levels in cerebrospinal fluid of pediatric hydrocephalus and brain tumor patients. European Journal of Paediatric Neurology, 2008, 12, 334-341.	1.6	20
63	Multiplexed quantification of dementia biomarkers in the CSF of patients with early dementias and MCI: A multicenter study. Neurobiology of Aging, 2008, 29, 812-818.	3.1	94
64	Diagnostic performance of a CSF-biomarker panel in autopsy-confirmed dementia. Neurobiology of Aging, 2008, 29, 1143-1159.	3.1	217
65	Intra-Individual Stability of CSF Biomarkers for Alzheimer's Disease over Two Years. Journal of Alzheimer's Disease, 2007, 12, 255-260.	2.6	117
66	Analytical performance and clinical utility of the INNOTEST® PHOSPHO-TAU(181P) assay for discrimination between Alzheimer's disease and dementia with Lewy bodies. Clinical Chemistry and Laboratory Medicine, 2006, 44, 1472-80.	2.3	145
67	Effects of Processing and Storage Conditions on Amyloid β (1–42) and Tau Concentrations in Cerebrospinal Fluid: Implications for Use in Clinical Practice. Clinical Chemistry, 2005, 51, 189-195.	3.2	151
68	Amino-Truncated β-Amyloid42 Peptides in Cerebrospinal Fluid and Prediction of Progression of Mild Cognitive Impairment. Clinical Chemistry, 2005, 51, 1650-1660.	3.2	82
69	Differences and Similarities between Two Frequently Used Assays for Amyloid β 42 in Cerebrospinal Fluid. Clinical Chemistry, 2005, 51, 1057-1060.	3.2	9
70	Simultaneous Measurement of β-Amyloid(1–42), Total Tau, and Phosphorylated Tau (Thr181) in Cerebrospinal Fluid by the xMAP Technology. Clinical Chemistry, 2005, 51, 336-345.	3.2	400
71	Measurement of Phosphorylated Tau Epitopes in the Differential Diagnosisof Alzheimer Disease. Archives of General Psychiatry, 2004, 61, 95.	12.3	390
72	Neurotoxicity Marker Profiles in the CSF are not Age-Dependent but Show Variation in Children Treated for Acute Lymphoblastic Leukemia. NeuroToxicology, 2004, 25, 471-480.	3.0	16

#	Article	IF	CITATIONS
73	Plasma Levels of β-Amyloid(1-40), β-Amyloid(1-42), and Total β-Amyloid Remain Unaffected in Adult Patients With Hypercholesterolemia After Treatment With Statins. Archives of Neurology, 2004, 61, 333.	4.5	109
74	Cerebrospinal fluid levels of total-tau, phospho-tau and Al²42 predicts development of Alzheimer's disease in patients with mild cognitive impairment. Acta Neurologica Scandinavica, 2003, 107, 47-51.	2.1	140
75	Unaltered Plasma Levels of β-Amyloid <sub>(1–40)</sub> and β-Amyloid <sub>(1–42)</sub> upon Stimulation of Human Platelets. Dementia and Geriatric Cognitive Disorders, 2003, 16, 93-97.	1.5	16
76	Transient increase in total tau but not phospho-tau in human cerebrospinal fluid after acute stroke. Neuroscience Letters, 2001, 297, 187-190.	2.1	401
77	Tau and Aβ42 in Cerebrospinal Fluid from Healthy Adults 21–93 Years of Age: Establishment of Reference Values. Clinical Chemistry, 2001, 47, 1776-1781.	3.2	420
78	The Cerebrospinal Fluid Levels of Tau, Growth-Associated Protein-43 and Soluble Amyloid Precursor Protein Correlate in Alzheimer's Disease, Reflecting a Common Pathophysiological Process. Dementia and Geriatric Cognitive Disorders, 2001, 12, 257-264.	1.5	102
79	Cerebrospinal fluid Ï,, and β-amyloid(1-42) in dementia disorders. Mechanisms of Ageing and Development, 2001, 122, 2005-2011.	4.6	50
80	Evaluation of CSF-tau and CSF-Aβ42 as Diagnostic Markers for Alzheimer Disease in Clinical Practice. Archives of Neurology, 2001, 58, 373-9.	4.5	487
81	The discrepancy between presenilin subcellular localization and <b>γ</b> -secretase processing of amyloid precursor protein. Journal of Cell Biology, 2001, 154, 731-740.	5.2	155
82	Cerebrospinal Fluid Markers for Alzheimer's Disease Evaluated after Acute Ischemic Stroke. Journal of Alzheimer's Disease, 2000, 2, 199-206.	2.6	183
83	Aging Increased Amyloid Peptide and Caused Amyloid Plaques in Brain of Old APP/V717I Transgenic Mice by a Different Mechanism than Mutant Presenilin1. Journal of Neuroscience, 2000, 20, 6452-6458.	3.6	107
84	Prominent Cerebral Amyloid Angiopathy in Transgenic Mice Overexpressing the London Mutant of Human APP in Neurons. American Journal of Pathology, 2000, 157, 1283-1298.	3.8	213
85	Standardization of measurement of β-amyloid <sub>(1-42)</sub> in cerebrospinal fluid and plasma. Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis, 2000, 7, 245-258.	3.0	286
86	Cerebrospinal Fluid β-Amyloid(1-42) in Alzheimer Disease. Archives of Neurology, 1999, 56, 673.	4.5	594
87	The Glu318Gly Substitution in Presenilin 1 Is Not Causally Related to Alzheimer Disease. American Journal of Human Genetics, 1999, 64, 290-292.	6.2	47
88	Evidence That Aβ42 Plasma Levels in Presenilin-1 Mutation Carriers Do not Allow for Prediction of Their Clinical Phenotype. Neurobiology of Disease, 1999, 6, 280-287.	4.4	48
89	Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. Nature, 1998, 391, 387-390.	27.8	1,765