

# Mathieu C Tamby

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10973347/publications.pdf>

Version: 2024-02-01

19  
papers

979  
citations

516710

16  
h-index

752698

20  
g-index

20  
all docs

20  
docs citations

20  
times ranked

1134  
citing authors

#	ARTICLE	IF	CITATIONS
1	New insights into the pathogenesis of systemic sclerosis. <i>Autoimmunity Reviews</i> , 2003, 2, 152-157.	5.8	141
2	Identification of Target Antigens of Antifibroblast Antibodies in Pulmonary Arterial Hypertension. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2008, 177, 1128-1134.	5.6	112
3	Pathogenesis of giant cell arteritis: More than just an inflammatory condition?. <i>Autoimmunity Reviews</i> , 2010, 9, 635-645.	5.8	110
4	Targets of anti-endothelial cell antibodies in pulmonary hypertension and scleroderma. <i>European Respiratory Journal</i> , 2012, 39, 1405-1414.	6.7	90
5	The Role of Inflammation and Autoimmunity in the Pathophysiology of Pulmonary Arterial Hypertension. <i>Clinical Reviews in Allergy and Immunology</i> , 2013, 44, 31-38.	6.5	85
6	IgG reactivity with a 100-kDa tissue and endothelial cell antigen identified as topoisomerase 1 distinguishes between limited and diffuse systemic sclerosis patients. <i>Clinical Immunology</i> , 2004, 111, 241-251.	3.2	49
7	Identification of target antigens of anti-endothelial cell and anti-vascular smooth muscle cell antibodies in patients with giant cell arteritis: a proteomic approach. <i>Arthritis Research and Therapy</i> , 2011, 13, R107.	3.5	44
8	Anti-endothelial cell antibodies from patients with limited cutaneous systemic sclerosis bind to centromeric protein B (CENP-B). <i>Clinical Immunology</i> , 2006, 120, 212-219.	3.2	42
9	IgG from patients with pulmonary arterial hypertension and/or systemic sclerosis binds to vascular smooth muscle cells and induces cell contraction. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 596-605.	0.9	41
10	Identification of target antigens of antiendothelial cell antibodies in healthy individuals: A proteomic approach. <i>Proteomics</i> , 2008, 8, 1000-1008.	2.2	39
11	Serum Eosinophil Cationic Protein: A Marker of Disease Activity in Churg-Strauss Syndrome. <i>Annals of the New York Academy of Sciences</i> , 2007, 1107, 392-399.	3.8	38
12	IgM and IgG autoantibodies from microscopic polyangiitis patients but not those with other small- and medium-sized vessel vasculitides recognize multiple endothelial cell antigens. <i>Clinical Immunology</i> , 2003, 109, 165-178.	3.2	34
13	Proteomes of umbilical vein and microvascular endothelial cells reflect distinct biological properties and influence immune recognition. <i>Proteomics</i> , 2012, 12, 2547-2555.	2.2	28
14	A Combined SDS-PAGE and Proteomics Approach to Identify Target Autoantigens in Healthy Individuals and Patients with Autoimmune Diseases. <i>Annals of the New York Academy of Sciences</i> , 2007, 1109, 538-549.	3.8	21
15	Identification of new autoantibody specificities directed at proteins involved in the transforming growth factor $\beta$ pathway in patients with systemic sclerosis. <i>Arthritis Research and Therapy</i> , 2011, 13, R74.	3.5	17
16	Analysis of autoantibody repertoires in small- and medium-sized vessels vasculitides. Evidence for specific perturbations in polyarteritis nodosa, microscopic polyangiitis, Churg's "Strauss syndrome and Wegener's granulomatosis. <i>Journal of Autoimmunity</i> , 2005, 24, 169-179.	6.5	15
17	Immunoblotting on HEp-2 cells increases the detection of antitopoisomerase 1 antibodies in patients with systemic sclerosis. <i>Clinical Immunology</i> , 2007, 123, 82-88.	3.2	9
18	Antitopoisomerase 1 Antibodies in Systemic Sclerosis: How to Improve the Detection?. <i>Annals of the New York Academy of Sciences</i> , 2007, 1109, 221-228.	3.8	7

#	ARTICLE	IF	CITATIONS
19	IgG from patients with systemic sclerosis bind to DNA antitopoisomerase 1 in normal human fibroblasts extracts. <i>Biologics: Targets and Therapy</i> , 2008, 2, 583.	3.2	6