

Linjie Tian

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10999539/publications.pdf>

Version: 2024-02-01

16
papers

484
citations

759233

12
h-index

940533

16
g-index

16
all docs

16
docs citations

16
times ranked

711
citing authors

#	ARTICLE	IF	CITATIONS
1	Liposome Preparation for the Analysis of Lipid- α Receptor Interaction and Efferocytosis. <i>Current Protocols in Immunology</i> , 2018, 120, 14.44.1-14.44.21.	3.6	8
2	Dendritic cells expressing immunoreceptor CD300f are critical for controlling chronic gut inflammation. <i>Journal of Clinical Investigation</i> , 2017, 127, 1905-1917.	8.2	17
3	Lipopolysaccharide-Induced CD300b Receptor Binding to Toll-like Receptor 4 Alters Signaling to Drive Cytokine Responses that Enhance Septic Shock. <i>Immunity</i> , 2016, 44, 1365-1378.	14.3	54
4	Emerging role of CD300 receptors in regulating myeloid cell efferocytosis. <i>Molecular and Cellular Oncology</i> , 2015, 2, e964625.	0.7	29
5	CD300f associates with IL-4 receptor β and amplifies IL-4-induced immune cell responses. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 8708-8713.	7.1	48
6	p85 β recruitment by the CD300f phosphatidyserine receptor mediates apoptotic cell clearance required for autoimmunity suppression. <i>Nature Communications</i> , 2014, 5, 3146.	12.8	77
7	C-terminal peptides of chemokine-like factor 1 signal through chemokine receptor CCR4 to cross-desensitize the CXCR4. <i>Biochemical and Biophysical Research Communications</i> , 2011, 409, 356-361.	2.1	13
8	Cutting Edge: Mouse CD300f (CMRF-35 α -Like Molecule-1) Recognizes Outer Membrane-Exposed Phosphatidyserine and Can Promote Phagocytosis. <i>Journal of Immunology</i> , 2011, 187, 3483-3487.	0.8	76
9	Rat CC chemokine receptor 4 is the functional homologue of human CC chemokine receptor 4 and can interact with human CCL17 and CCL22. <i>Science Bulletin</i> , 2010, 55, 1388-1395.	1.7	1
10	Potent antitumor activities of recombinant human PDCD5 protein in combination with chemotherapy drugs in K562 cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 396, 224-230.	2.1	32
11	Human CMTM2/CKLFSF2 enhances the ligand-induced transactivation of the androgen receptor. <i>Science Bulletin</i> , 2009, 54, 1050-1057.	9.0	1
12	Interactions of Peptides from Secreted Human CKLF1 and the N-Terminal Extracellular Tail of CCR4 Analyzed by CZE. <i>Chromatographia</i> , 2009, 70, 287-292.	1.3	2
13	Two C-terminal peptides of human CKLF1 interact with the chemokine receptor CCR4. <i>International Journal of Biochemistry and Cell Biology</i> , 2008, 40, 909-919.	2.8	37
14	An alternative splice form of CMTM8 induces apoptosis. <i>International Journal of Biochemistry and Cell Biology</i> , 2007, 39, 2107-2119.	2.8	19
15	CMTM8 induces caspase-dependent and -independent apoptosis through a mitochondria-mediated pathway. <i>Journal of Cellular Physiology</i> , 2007, 211, 112-120.	4.1	31
16	Cellular Uptake of Exogenous Human PDCD5 Protein. <i>Journal of Biological Chemistry</i> , 2006, 281, 24803-24817.	3.4	39