Abdelouahid Tajar

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11008522/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Low vitamin D and the risk of developing chronic widespread pain: results from the European male ageing study. BMC Musculoskeletal Disorders, 2016, 17, 32.	1.9	25
2	The ability of three different models of frailty to predict all-cause mortality: Results from the European Male Aging Study (EMAS). Archives of Gerontology and Geriatrics, 2013, 57, 360-368.	3.0	121
3	Frailty and Sexual Health in Older European Men. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2013, 68, 837-844.	3.6	32
4	The association of frailty with serum 25-hydroxyvitamin D and parathyroid hormone levels in older European men. Age and Ageing, 2013, 42, 352-359.	1.6	74
5	Cohort Profile: The European Male Ageing Study. International Journal of Epidemiology, 2013, 42, 391-401.	1.9	41
6	The Onset of Widespread Musculoskeletal Pain Is Associated with a Decrease in Healthy Ageing in Older People: A Population-Based Prospective Study. PLoS ONE, 2013, 8, e59858.	2.5	33
7	Characteristics of Androgen Deficiency in Late-Onset Hypogonadism: Results from the European Male Aging Study (EMAS). Journal of Clinical Endocrinology and Metabolism, 2012, 97, 1508-1516.	3.6	258
8	Comparison of serum testosterone and estradiol measurements in 3174 European men using platform immunoassay and mass spectrometry; relevance for the diagnostics in aging men. European Journal of Endocrinology, 2012, 166, 983-991.	3.7	169
9	Association of hypogonadism with vitamin D status: the European Male Ageing Study. European Journal of Endocrinology, 2012, 166, 77-85.	3.7	166
10	Reference Ranges for Testosterone in Men Generated Using Liquid Chromatography Tandem Mass Spectrometry in a Community-Based Sample of Healthy Nonobese Young Men in the Framingham Heart Study and Applied to Three Geographically Distinct Cohorts. Journal of Clinical Endocrinology and Metabolism, 2011, 96, 2430-2439.	3.6	332
11	Lower vitamin D levels are associated with depression among community-dwelling European men. Journal of Psychopharmacology, 2011, 25, 1320-1328.	4.0	99
12	Low testosterone in ageing men: a modifiable risk factor for frailty?. Trends in Endocrinology and Metabolism, 2011, 22, 491-498.	7.1	13
13	Frailty in Relation to Variations in Hormone Levels of the Hypothalamic-Pituitary-Testicular Axis in Older Men: Results From the European Male Aging Study. Journal of the American Geriatrics Society, 2011, 59, 814-821.	2.6	52
14	Elevated levels of gonadotrophins but not sex steroids are associated with musculoskeletal pain in middle-aged and older European men. Pain, 2011, 152, 1495-1501.	4.2	24
15	The Relationships between Sex Hormones and Sexual Function in Middle-Aged and Older European Men. Journal of Clinical Endocrinology and Metabolism, 2011, 96, E1577-E1587.	3.6	103
16	The Effect of Musculoskeletal Pain on Sexual Function in Middle-aged and Elderly European Men: Results from the European Male Ageing Study. Journal of Rheumatology, 2011, 38, 370-377.	2.0	16
17	Influence of bone remodelling rate on quantitative ultrasound parameters at the calcaneus and DXA BMDa of the hip and spine in middle-aged and elderly European men: the European Male Ageing Study (EMAS). European Journal of Endocrinology, 2011, 165, 977-986.	3.7	28
18	Obesity and weight management in the elderly. British Medical Bulletin, 2011, 97, 169-196.	6.9	249

Abdelouahid Tajar

#	Article	IF	CITATIONS
19	Impaired quality of life and sexual function in overweight and obese men: the European Male Ageing Study. European Journal of Endocrinology, 2011, 164, 1003-1011.	3.7	90
20	Do the Effects of Testosterone on Muscle Strength, Physical Function, Body Composition, And Quality of Life Persist Six Months after Treatment in Intermediate-Frail and Frail Elderly Men?. Journal of Clinical Endocrinology and Metabolism, 2011, 96, 454-458.	3.6	83
21	Genetic Determinants of Serum Testosterone Concentrations in Men. PLoS Genetics, 2011, 7, e1002313.	3.5	178
22	Characteristics of Secondary, Primary, and Compensated Hypogonadism in Aging Men: Evidence from the European Male Ageing Study. Journal of Clinical Endocrinology and Metabolism, 2010, 95, 1810-1818.	3.6	481
23	Endogenous hormones, androgen receptor CAG repeat length and fluid cognition in middle-aged and older men: results from the European Male Ageing Study. European Journal of Endocrinology, 2010, 162, 1155-1164.	3.7	25
24	Musculoskeletal pain is associated with very low levels of vitamin D in men: results from the European Male Ageing Study. Annals of the Rheumatic Diseases, 2010, 69, 1448-1452.	0.9	86
25	Identification of Late-Onset Hypogonadism in Middle-Aged and Elderly Men. New England Journal of Medicine, 2010, 363, 123-135.	27.0	1,274
26	On Concordance Measures for Discrete Data and Dependence Properties of Poisson Model. Journal of Probability and Statistics, 2009, 2009, 1-15.	0.7	3
27	Perturbed Insulin-like Growth Factor-1 (IGF-1) and IGF Binding Protein-3 Are Not Associated with Chronic Widespread Pain in Men: Results from the European Male Ageing Study. Journal of Rheumatology, 2009, 36, 2523-2530.	2.0	3
28	The association between different cognitive domains and age in a multiâ€centre study of middleâ€aged and older European men. International Journal of Geriatric Psychiatry, 2009, 24, 1257-1266.	2.7	10
29	The European Male Ageing Study (EMAS): design, methods and recruitment. Journal of Developmental and Physical Disabilities, 2009, 32, 11-24.	3.6	137
30	Association between 25-hydroxyvitamin D levels and cognitive performance in middle-aged and older European men. Journal of Neurology, Neurosurgery and Psychiatry, 2009, 80, 722-729.	1.9	130
31	Assessment of Sexual Health in Aging Men in Europe: Development and Validation of the European Male Ageing Study Sexual Function Questionnaire. Journal of Sexual Medicine, 2008, 5, 1374-1385.	0.6	80
32	Hypothalamic-Pituitary-Testicular Axis Disruptions in Older Men Are Differentially Linked to Age and Modifiable Risk Factors: The European Male Aging Study. Journal of Clinical Endocrinology and Metabolism, 2008, 93, 2737-2745.	3.6	790