

Daniella J Furman

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11049456/publications.pdf>

Version: 2024-02-01

17
papers

2,134
citations

623734

14
h-index

888059

17
g-index

18
all docs

18
docs citations

18
times ranked

3757
citing authors

#	ARTICLE	IF	CITATIONS
1	Functional Neuroimaging of Major Depressive Disorder: A Meta-Analysis and New Integration of Baseline Activation and Neural Response Data. <i>American Journal of Psychiatry</i> , 2012, 169, 693-703.	7.2	660
2	Default-Mode and Task-Positive Network Activity in Major Depressive Disorder: Implications for Adaptive and Maladaptive Rumination. <i>Biological Psychiatry</i> , 2011, 70, 327-333.	1.3	646
3	Frontostriatal functional connectivity in major depressive disorder. <i>Biology of Mood & Anxiety Disorders</i> , 2011, 1, 11.	4.7	166
4	Variant in oxytocin receptor gene is associated with amygdala volume. <i>Psychoneuroendocrinology</i> , 2011, 36, 891-897.	2.7	126
5	Interoceptive awareness, positive affect, and decision making in Major Depressive Disorder. <i>Journal of Affective Disorders</i> , 2013, 151, 780-785.	4.1	108
6	Variant in oxytocin receptor gene is associated with amygdala volume. <i>Psychoneuroendocrinology</i> , 2011, 36, 891-897.	2.7	97
7	Anticipatory and consummatory pleasure and displeasure in major depressive disorder: An experience sampling study.. <i>Journal of Abnormal Psychology</i> , 2017, 126, 149-159.	1.9	62
8	Dopamine Synthesis Capacity is Associated with D2/3 Receptor Binding but Not Dopamine Release. <i>Neuropsychopharmacology</i> , 2018, 43, 1201-1211.	5.4	43
9	Amygdala responses to salient social cues vary with oxytocin receptor genotype in youth. <i>Neuropsychologia</i> , 2015, 79, 1-9.	1.6	38
10	Habenula responses to potential and actual loss in major depression: preliminary evidence for lateralized dysfunction. <i>Social Cognitive and Affective Neuroscience</i> , 2016, 11, 843-851.	3.0	36
11	Dopaminergic Mechanisms Underlying Normal Variation in Trait Anxiety. <i>Journal of Neuroscience</i> , 2019, 39, 2735-2744.	3.6	36
12	Altered timing of amygdala activation during sad mood elaboration as a function of 5-HTTLPR. <i>Social Cognitive and Affective Neuroscience</i> , 2011, 6, 270-276.	3.0	35
13	Associations among locus coeruleus catecholamines, tau pathology, and memory in aging. <i>Neuropsychopharmacology</i> , 2022, 47, 1106-1113.	5.4	27
14	Effects of Dopaminergic Drugs on Cognitive Control Processes Vary by Genotype. <i>Journal of Cognitive Neuroscience</i> , 2020, 32, 804-821.	2.3	18
15	Activation of the medial prefrontal and posterior cingulate cortex during encoding of negative material predicts symptom worsening in major depression. <i>NeuroReport</i> , 2014, 25, 324-329.	1.2	17
16	Augmenting Frontal Dopamine Tone Enhances Maintenance over Gating Processes in Working Memory. <i>Journal of Cognitive Neuroscience</i> , 2021, 33, 1753-1765.	2.3	13
17	Enhancing dopamine tone modulates global and local cortical perfusion as a function of COMT val158met genotype. <i>NeuroImage</i> , 2021, 242, 118472.	4.2	5