

Rh Belmaker

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11055107/publications.pdf>

Version: 2024-02-01

97
papers

6,196
citations

109321

35
h-index

69250

77
g-index

99
all docs

99
docs citations

99
times ranked

6578
citing authors

#	ARTICLE	IF	CITATIONS
1	A multi-national, multi-disciplinary Delphi consensus study on using omega-3 polyunsaturated fatty acids (n-3 PUFAs) for the treatment of major depressive disorder. <i>Journal of Affective Disorders</i> , 2020, 265, 233-238.	4.1	12
2	International Society for Nutritional Psychiatry Research Practice Guidelines for Omega-3 Fatty Acids in the Treatment of Major Depressive Disorder. <i>Psychotherapy and Psychosomatics</i> , 2019, 88, 263-273.	8.8	114
3	Influence of birth cohort on age of onset cluster analysis in bipolar I disorder. <i>European Psychiatry</i> , 2015, 30, 99-105.	0.2	28
4	Molecular effects of lithium are partially mimicked by inositol-monophosphatase (IMPA)1 knockout mice in a brain region-dependent manner. <i>European Neuropsychopharmacology</i> , 2015, 25, 425-434.	0.7	23
5	Inositol-Deficient Food Augments a Behavioral Effect of Long-Term Lithium Treatment Mediated by Inositol Monophosphatase Inhibition. <i>Journal of Clinical Psychopharmacology</i> , 2015, 35, 175-177.	1.4	3
6	Lurasidone and Bipolar Disorder. <i>American Journal of Psychiatry</i> , 2014, 171, 131-133.	7.2	14
7	Inhibition of inositol monophosphatase (IMPase) at the calbindin-D28k binding site: Molecular and behavioral aspects. <i>European Neuropsychopharmacology</i> , 2013, 23, 1806-1815.	0.7	9
8	Behavioral addictions in euthymic patients with bipolar I disorder: a comparison to controls. <i>International Journal of Bipolar Disorders</i> , 2013, 1, 27.	2.2	5
9	The inositol monophosphatase inhibitor L-690,330 affects pilocarpine-behavior and the forced swim test. <i>Psychopharmacology</i> , 2013, 227, 503-508.	3.1	5
10	Acute Intracerebroventricular Inositol Does Not Reverse the Effect of Chronic Lithium Treatment in the Forced Swim Test. <i>Neuropsychobiology</i> , 2013, 68, 189-192.	1.9	8
11	Effects of lithium on lipopolysaccharide-induced inflammation in rat primary glia cells. <i>Innate Immunity</i> , 2012, 18, 447-458.	2.4	62
12	Individual differences and evidence-based psychopharmacology. <i>BMC Medicine</i> , 2012, 10, 110.	5.5	15
13	Gene-expression studies in understanding the mechanism of action of lithium. <i>Expert Review of Neurotherapeutics</i> , 2012, 12, 93-97.	2.8	14
14	Hyperhomocysteinemia does not affect global DNA methylation and nicotinamide N-methyltransferase expression in mice. <i>Journal of Psychopharmacology</i> , 2011, 25, 976-981.	4.0	9
15	Valnoctamide as a valproate substitute with low teratogenic potential in mania: a double-blind, controlled, add-on clinical trial. <i>Bipolar Disorders</i> , 2010, 12, 376-382.	1.9	43
16	The New Lithium Clinic. <i>Neuropsychobiology</i> , 2010, 62, 17-26.	1.9	16
17	No gross abnormality of plasma homocysteine after acute methionine loading in clinically stabilized patients with schizophrenia. <i>Asian Journal of Psychiatry</i> , 2010, 3, 64-66.	2.0	3
18	Identification of eukaryotic elongation factor-2 as a novel cellular target of lithium and glycogen synthase kinase-3. <i>Molecular and Cellular Neurosciences</i> , 2010, 45, 449-455.	2.2	18

#	ARTICLE	IF	CITATIONS
19	Subtyping Major Depressive Disorder. <i>Psychotherapy and Psychosomatics</i> , 2010, 79, 131-135.	8.8	103
20	Omega 3 Fatty Acid Treatment in Autism. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2009, 19, 449-451.	1.3	58
21	Knockout mice in understanding the mechanism of action of lithium. <i>Biochemical Society Transactions</i> , 2009, 37, 1121-1125.	3.4	48
22	Neuropsychological correlates of homocysteine levels in euthymic bipolar patients. <i>Journal of Affective Disorders</i> , 2008, 105, 229-233.	4.1	27
23	No association between global leukocyte DNA methylation and homocysteine levels in schizophrenia patients. <i>Schizophrenia Research</i> , 2008, 101, 50-57.	2.0	55
24	Major Depressive Disorder. <i>New England Journal of Medicine</i> , 2008, 358, 55-68.	27.0	1,600
25	Antidepressive-like effects of rapamycin in animal models: Implications for mTOR inhibition as a new target for treatment of affective disorders. <i>Brain Research Bulletin</i> , 2008, 76, 469-473.	3.0	121
26	Glycogen synthase kinase-3 β heterozygote knockout mice as a model of findings in postmortem schizophrenia brain or as a model of behaviors mimicking lithium action: negative results. <i>Behavioural Pharmacology</i> , 2008, 19, 217-224.	1.7	38
27	Treatment of Bipolar Depression. <i>New England Journal of Medicine</i> , 2007, 356, 1771-1773.	27.0	23
28	Lithium's effect in forced-swim test is blood level dependent but not dependent on weight loss. <i>Behavioural Pharmacology</i> , 2007, 18, 77-80.	1.7	52
29	Mitochondrial DNA HV lineage increases the susceptibility to schizophrenia among Israeli Arabs. <i>Schizophrenia Research</i> , 2007, 94, 354-358.	2.0	39
30	Lithium's pilocarpine seizures as a model for lithium action in mania. <i>Neuroscience and Biobehavioral Reviews</i> , 2007, 31, 843-849.	6.1	20
31	P.8.a.005 Familial heritability of increased homocysteine in schizophrenia. <i>European Neuropsychopharmacology</i> , 2006, 16, S537.	0.7	0
32	Homocysteine-Reducing Strategies Improve Symptoms in Chronic Schizophrenic Patients with Hyperhomocysteinemia. <i>Biological Psychiatry</i> , 2006, 60, 265-269.	1.3	117
33	SMIT1 haploinsufficiency causes brain inositol deficiency without affecting lithium-sensitive behavior. <i>Molecular Genetics and Metabolism</i> , 2006, 88, 384-388.	1.1	32
34	Transmission disequilibrium and haplotype analyses of the G72/G30 locus: Suggestive linkage to schizophrenia in Palestinian Arabs living in the North of Israel. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2006, 141B, 91-95.	1.7	49
35	Omega-3 Treatment of Childhood Depression: A Controlled, Double-Blind Pilot Study. <i>American Journal of Psychiatry</i> , 2006, 163, 1098-1100.	7.2	303
36	Nutritional and life style determinants of plasma homocysteine in schizophrenia patients. <i>European Neuropsychopharmacology</i> , 2005, 15, 291-295.	0.7	23

#	ARTICLE	IF	CITATIONS
37	High homocysteine serum levels in young male schizophrenia and bipolar patients and in an animal model. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2005, 29, 1181-1191.	4.8	52
38	Epi-inositol is ineffective in Porsolt Forced Swim Test model of depression. <i>Neuropsychiatric Disease and Treatment</i> , 2005, 1, 189-190.	2.2	2
39	Homocysteine levels in newly admitted schizophrenic patients. <i>Journal of Psychiatric Research</i> , 2004, 38, 413-416.	3.1	98
40	Bipolar Disorder. <i>New England Journal of Medicine</i> , 2004, 351, 476-486.	27.0	468
41	Lithium inhibitable enzymes in postmortem brain of bipolar patients. <i>Journal of Psychiatric Research</i> , 2003, 37, 433-442.	3.1	14
42	Right prefrontal TMS versus sham treatment of mania: a controlled study. <i>Bipolar Disorders</i> , 2003, 5, 36-39.	1.9	80
43	The effect of lithium on expression of genes for inositol biosynthetic enzymes in mouse hippocampus; a comparison with the yeast model. <i>Molecular Brain Research</i> , 2003, 115, 104-110.	2.3	33
44	GSK-3 and the neurodevelopmental hypothesis of schizophrenia. <i>European Neuropsychopharmacology</i> , 2002, 12, 13-25.	0.7	122
45	Reduced inositol content in lymphocyte-derived cell lines from bipolar patients. <i>Bipolar Disorders</i> , 2002, 4, 67-69.	1.9	26
46	Low GSK-3 activity in frontal cortex of schizophrenic patients. <i>Schizophrenia Research</i> , 2001, 52, 101-105.	2.0	130
47	The antidepressant activity of inositol in the forced swim test involves 5-HT ₂ receptors. <i>Behavioural Brain Research</i> , 2001, 118, 77-83.	2.2	58
48	No evidence for linkage by transmission disequilibrium test analysis of microsatellite marker D22S278 and schizophrenia in a Palestinian Arab and in a German population. <i>American Journal of Medical Genetics Part A</i> , 2001, 105, 328-331.	2.4	3
49	The effects of inositol treatment in animal models of psychiatric disorders. <i>Journal of Affective Disorders</i> , 2001, 62, 113-121.	4.1	56
50	Nordidemnin potently inhibits inositol uptake in cultured astrocytes and dose-dependently augments lithium's proconvulsant effect in vivo. , 2000, 60, 116-121.		11
51	No association between the dopamine D ₃ receptor 3'UTR polymorphism and schizophrenia in a family-based study of a Palestinian Arab population. <i>American Journal of Medical Genetics Part A</i> , 2000, 96, 778-780.	2.4	18
52	Epi-inositol: A potential antidepressant. <i>Drug Development Research</i> , 2000, 50, 309-315.	2.9	4
53	Rorschach markers in offspring of manic-depressive patients. <i>Journal of Affective Disorders</i> , 2000, 59, 231-236.	4.1	11
54	Chronic treatment of human astrocytoma cells with lithium, carbamazepine or valproic acid decreases inositol uptake at high inositol concentrations but increases it at low inositol concentrations. <i>Brain Research</i> , 2000, 855, 158-161.	2.2	40

#	ARTICLE	IF	CITATIONS
55	Psychological responses in family members after the Hebron massacre. <i>Depression and Anxiety</i> , 1999, 9, 27-31.	4.1	29
56	Rat brain monoamines after acute and chronic myo-inositol treatment. <i>European Neuropsychopharmacology</i> , 1999, 10, 27-30.	0.7	7
57	Chronic treatment with lithium and pretreatment with excess inositol reduce inositol pool size in astrocytes by different mechanisms. <i>Brain Research</i> , 1998, 787, 34-40.	2.2	24
58	Transcranial Magnetic Stimulation in Mania: A Controlled Study. <i>American Journal of Psychiatry</i> , 1998, 155, 1608-1610.	7.2	185
59	S.07.05 The effects of transcranial magnetic stimulation on $\hat{1}^2$ -adrenergic receptors and brain monoamines. <i>European Neuropsychopharmacology</i> , 1997, 7, S93-S94.	0.7	0
60	The Effects of TMS on Animal Models of Depression, $\hat{1}^2$ -Adrenergic Receptors, and Brain Monoamines. <i>CNS Spectrums</i> , 1997, 2, 26-30.	1.2	24
61	Phorbol ester intracerebroventricularly induces a behavioral hypoactivity that is not affected by chronic or acute lithium. <i>European Neuropsychopharmacology</i> , 1996, 6, 39-41.	0.7	4
62	Differential uptake of myo-inositol in vivo into rat brain areas. <i>European Neuropsychopharmacology</i> , 1996, 6, 73-75.	0.7	38
63	Lack of effect of ECS on rat brain inositol monophosphatase activity and inositol levels and of i.c.v. inositol on ictal and post-ictal length. <i>Journal of Psychiatric Research</i> , 1996, 30, 39-43.	3.1	0
64	TPQ in euthymic manic-depressive patients. <i>Journal of Psychiatric Research</i> , 1996, 30, 353-357.	3.1	78
65	Lack of effect of 6 g inositol treatment on post-ECT cognitive function in humans. <i>Journal of Psychiatric Research</i> , 1995, 29, 487-489.	3.1	11
66	The effect of transcranial magnetic stimulation of rat brain on behavioral models of depression. <i>Brain Research</i> , 1995, 699, 130-132.	2.2	109
67	High-dose peripheral inositol raises brain inositol levels and reverses behavioral effects of inositol depletion by lithium. <i>Pharmacology Biochemistry and Behavior</i> , 1994, 49, 341-343.	2.9	32
68	Genetic markers, temperament, and psychopathology. <i>Biological Psychiatry</i> , 1994, 36, 71-72.	1.3	16
69	Transcranial magnetic stimulation in depression and schizophrenia. <i>European Neuropsychopharmacology</i> , 1994, 4, 287-288.	0.7	115
70	CSF inositol in schizophrenia and high-dose inositol treatment of schizophrenia. <i>European Neuropsychopharmacology</i> , 1994, 4, 487-490.	0.7	18
71	Behavioral evidence for the existence of two pools of cellular inositol. <i>European Neuropsychopharmacology</i> , 1994, 4, 463-467.	0.7	28
72	The 4-dedimethylamino derivative of tetracycline loses both ability to block cyclic AMP accumulation and ability to inhibit rat motor activity. <i>European Neuropsychopharmacology</i> , 1994, 4, 419-420.	0.7	3

#	ARTICLE	IF	CITATIONS
73	The effect of inositol on cognitive processes and mood states in normal volunteers. <i>European Neuropsychopharmacology</i> , 1994, 4, 417.	0.7	6
74	Inositol. <i>European Neuropsychopharmacology</i> , 1994, 4, 165-166.	0.7	0
75	The effect of transcranial magnetic stimulation compared with electroconvulsive shock on rat apomorphine-induced stereotypy. <i>European Neuropsychopharmacology</i> , 1994, 4, 449.	0.7	28
76	Effects of inositol on lithium-induced EEG abnormalities. <i>European Neuropsychopharmacology</i> , 1994, 4, 419.	0.7	0
77	Species differences in susceptibility to Li-pilocarpine seizures. <i>European Neuropsychopharmacology</i> , 1994, 4, 428-429.	0.7	4
78	Mechanism of lithium lethality in rats. <i>Journal of Psychiatric Research</i> , 1993, 27, 415-422.	3.1	1
79	Inositol treatment raises CSF inositol levels. <i>Brain Research</i> , 1993, 627, 168-170.	2.2	68
80	Dose-response and time curve of inositol prevention of Li-pilocarpine seizures. <i>European Neuropsychopharmacology</i> , 1993, 3, 428-429.	0.7	12
81	Biochemical, behavioral, and clinical studies of the role of inositol in lithium treatment and depression. <i>Biological Psychiatry</i> , 1993, 34, 839-852.	1.3	105
82	Lack of effect of inositol treatment in chronic schizophrenia. <i>Biological Psychiatry</i> , 1993, 33, 673-675.	1.3	16
83	Linkage of a normal personality trait to the color-blindness gene: preliminary evidence. <i>Biological Psychiatry</i> , 1993, 34, 581-583.	1.3	9
84	Raised monophosphatase activity in schizophrenic patients. <i>Clinica Chimica Acta</i> , 1992, 209, 89-93.	1.1	20
85	FAILURE OF ADDITION OF LITHIUM TO IMIPRAMINE TO ENHANCE ACTIVITY IN RATS OR MOOD IN NORMAL VOLUNTEERS. <i>Basic and Clinical Pharmacology and Toxicology</i> , 1992, 71, 18-25.	0.0	0
86	Inhibition by antibiotic tetracyclines of rat cortical noradrenergic adenylate cyclase and amphetamine-induced hyperactivity. <i>Pharmacology Biochemistry and Behavior</i> , 1990, 37, 417-424.	2.9	22
87	Inositol-1-phosphatase in red blood cells of manic-depressive patients before and during treatment with lithium. <i>Biological Psychiatry</i> , 1990, 27, 552-555.	1.3	29
88	Lithium research: State of the art. <i>Biological Psychiatry</i> , 1990, 27, 1279-1281.	1.3	15
89	Intracerebroventricular myo-inositol antagonizes lithium-induced suppression of rearing behaviour in rats. <i>Brain Research</i> , 1990, 534, 345-347.	2.2	38
90	Lithium inhibits adrenergic and cholinergic increases in GTP binding in rat cortex. <i>Nature</i> , 1988, 331, 440-442.	27.8	489

#	ARTICLE	IF	CITATIONS
91	Effects of lithium in vitro and ex vivo on components of the adenylate cyclase system in membranes from the cerebral cortex of the rat. <i>Neuropharmacology</i> , 1987, 26, 211-217.	4.1	142
92	Interstrain correlation between behavioural effects of lithium and effects on cortical cyclic AMP. <i>Pharmacology Biochemistry and Behavior</i> , 1986, 24, 9-13.	2.9	36
93	Effects of lithium in vitro on noradrenaline-induced cyclic AMP accumulation in rat cortical slices after reserpine-induced supersensitivity. <i>Neuropharmacology</i> , 1985, 24, 353-355.	4.1	6
94	Effect of Treatment and Withdrawal from Chronic Lithium in Rats on Stimulant-Induced Responses. <i>Neuropsychobiology</i> , 1984, 11, 28-32.	1.9	31
95	Rorschach Markers in Euthymic Manic-Depressive Illness. <i>Neuropsychobiology</i> , 1984, 12, 96-100.	1.9	12
96	The effects of chronic lithium and ECT on A1 and A2 adenosine receptor systems in rat brain. <i>Brain Research</i> , 1984, 291, 188-192.	2.2	46
97	Lack of Benefit from Magnesium in Lithium Toxicity. <i>Neuropsychobiology</i> , 1982, 8, 10-11.	1.9	5