

# Luiz Felipe De Campos-Lobato

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11187139/publications.pdf>

Version: 2024-02-01

14  
papers

833  
citations

1163117

8  
h-index

1125743

13  
g-index

14  
all docs

14  
docs citations

14  
times ranked

1101  
citing authors

#	ARTICLE	IF	CITATIONS
1	Is It Safe to Recommend Cholecystectomy Whenever Gallstones Develop After Bariatric Surgery?. Obesity Surgery, 2021, 31, 179-184.	2.1	2
2	Rectal prolapse surgery in males and females: An ACS NSQIP-based comparative analysis of over 12,000 patients. American Journal of Surgery, 2020, 220, 697-705.	1.8	6
3	Surgical Management of Complex Enteric Fistulas in Crohn's Disease. Clinics in Colon and Rectal Surgery, 2019, 32, 268-272.	1.1	2
4	Current Views on the Interval Between Neoadjuvant Chemoradiation and Surgery for Rectal Cancer. Current Colorectal Cancer Reports, 2017, 13, 234-239.	0.5	0
5	Extralevator Abdominal Perineal Excision Versus Standard Abdominal Perineal Excision: Impact on Quality of the Resected Specimen and Postoperative Morbidity. World Journal of Surgery, 2017, 41, 2160-2167.	1.6	25
6	Less Than 12 Nodes in the Surgical Specimen After Total Mesorectal Excision Following Neoadjuvant Chemoradiation: It means more than you think!. Annals of Surgical Oncology, 2013, 20, 3398-3406.	1.5	59
7	Prone or Lithotomy Positioning During an Abdominoperineal Resection for Rectal Cancer Results in Comparable Oncologic Outcomes. Diseases of the Colon and Rectum, 2011, 54, 939-946.	1.3	61
8	Pathologic Complete Response After Neoadjuvant Treatment for Rectal Cancer Decreases Distant Recurrence and Could Eradicate Local Recurrence. Annals of Surgical Oncology, 2011, 18, 1590-1598.	1.5	165
9	Neoadjuvant Therapy for Rectal Cancer: The Impact of Longer Interval Between Chemoradiation and Surgery. Journal of Gastrointestinal Surgery, 2011, 15, 444-450.	1.7	130
10	Response to Letter to the Editor: Neoadjuvant Therapy for Rectal Cancer: The Impact of Longer Interval Between Chemoradiation and Surgery. Journal of Gastrointestinal Surgery, 2011, 15, 1293.	1.7	1
11	Downstaging Without Complete Pathologic Response After Neoadjuvant Treatment Improves Cancer Outcomes for cIII but not cII Rectal Cancers. Annals of Surgical Oncology, 2010, 17, 1758-1766.	1.5	45
12	Enterocutaneous Fistula Associated with Malignancy and Prior Radiation Therapy. Clinics in Colon and Rectal Surgery, 2010, 23, 176-181.	1.1	17
13	How to evaluate risk and identify stage II patients requiring referral to a medical oncologist: a surgeon's perspective. Oncology, 2010, 24, 14-6.	0.5	4
14	Predictive Factors of Pathologic Complete Response After Neoadjuvant Chemoradiation for Rectal Cancer. Annals of Surgery, 2009, 250, 582-589.	4.2	316