

# Per Sundberg

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11236252/publications.pdf>

Version: 2024-02-01

87

papers

2,517

citations

186265

28

h-index

233421

45

g-index

87

all docs

87

docs citations

87

times ranked

1507

citing authors

#	ARTICLE	IF	CITATIONS
1	Phylum Nemertea., 2020,, 121-123.	0	
2	Phylum Nemertea., 2019,, 145-147.	0	
3	Nemertean taxonomyâ€”Implementing changes in the higher ranks, dismissing Anopla and Enopla. <i>Zoologica Scripta</i> , 2019, 48, 118-119.	1.7	26
4	Dispersal and speciation: The cross Atlantic relationship of two parasitic cnidarians. <i>Molecular Phylogenetics and Evolution</i> , 2018, 126, 346-355.	2.7	6
5	The future of nemertean taxonomy (phylum Nemertea) â€” a proposal. <i>Zoologica Scripta</i> , 2016, 45, 579-582.	1.7	22
6	Phylum Nemertea., 2016,, 111-113.	0	
7	Evaluating the Utility of Single-Locus DNA Barcoding for the Identification of Ribbon Worms (Phylum) Tj ETQq1 1 0.784314 rgBT /Overl...	2.5	37
8	Thirty-Five Years of Nemertean (Nemertea) Researchâ€”Past, Present, and Future. <i>Zoological Science</i> , 2015, 32, 501-506.	0.7	8
9	Species Diversity of <i>Ramphogordius sanguineus</i> / <i>Lineus ruber</i> -Like Nemerteans (Nemertea) Tj ETQq1 1 0.784314 rgBT /Overlock 1 579-589.	0.7	25
10	DNA barcoding supports identification of Malacobdella species (Nemertea: Hoplonemertea). <i>Zoological Studies</i> , 2015, 54, e10.	0.3	7
11	Phylum Nemertea., 2015,, 205-209.	3	
12	Mutation and Selection Cause Codon Usage and Bias in Mitochondrial Genomes of Ribbon Worms (Nemertea). <i>PLoS ONE</i> , 2014, 9, e85631.	2.5	72
13	A Transcriptomic Approach to Ribbon Worm Systematics (Nemertea): Resolving the Pilidiophora Problem. <i>Molecular Biology and Evolution</i> , 2014, 31, 3206-3215.	8.9	68
14	Complete mitochondrial genome sequences of two parasitic/commensal nemerteans, <i>Gononemertes parasita</i> and <i>Nemertopsis tetriclitophila</i> (Nemertea: Hoplonemertea). <i>Parasites and Vectors</i> , 2014, 7, 273.	2.5	6
15	Taxonomic Identity of a Tetrodotoxin-Accumulating Ribbon-worm <i>Cephalothrix simula</i> (Nemertea: Palaeonemertea): A Species Artificially Introduced from the Pacific to Europe. <i>Zoological Science</i> , 2013, 30, 985-997.	0.7	20
16	A comparative study of nemertean complete mitochondrial genomes, including two new ones for <i>Nectonemertes cf. mirabilis</i> and <i>Zygeupolia rubens</i> , may elucidate the fundamental pattern for the phylum Nemertea. <i>BMC Genomics</i> , 2012, 13, 139.	2.8	16
17	Disentangling ribbon worm relationships: multiâ€—locus analysis supports traditional classification of the phylum Nemertea. <i>Cladistics</i> , 2012, 28, 141-159.	3.3	107
18	A DNA-based description of a new nemertean (phylum Nemertea) species. <i>Marine Biology Research</i> , 2011, 7, 63-70.	0.7	35

#	ARTICLE	IF	CITATIONS
19	Systematics and phylogeny of the hoplonemertean genus <i>Diplomma</i> (Nemertea) based on molecular and morphological evidence. <i>Zoological Journal of the Linnean Society</i> , 2011, 161, 695-722.	2.3	16
20	The mitochondrial genomes of two nemerteans, <i>Cephalothrix</i> sp. (Nemertea: Palaeonemertea) and <i>Paranemertes</i> cf. <i>peregrina</i> (Nemertea: Hoplonemertea). <i>Molecular Biology Reports</i> , 2011, 38, 4509-4525.	2.3	16
21	Parasitic anemone infects the invasive ctenophore <i>Mnemiopsis leidyi</i> in the North East Atlantic. <i>Biological Invasions</i> , 2010, 12, 1003-1009.	2.4	15
22	The <i>Lanius excubitor</i> (Aves, Passeriformes) conundrum—Taxonomic dilemma when molecular and non-molecular data tell different stories. <i>Molecular Phylogenetics and Evolution</i> , 2010, 55, 347-357.	2.7	41
23	The first internal molecular phylogeny of the animal phylum Entoprocta (Kamptozoa). <i>Molecular Phylogenetics and Evolution</i> , 2010, 56, 370-379.	2.7	28
24	Nemertean taxonomy - time to change lane?. <i>Journal of Zoological Systematics and Evolutionary Research</i> , 2010, 48, 283.	1.4	19
25	Statistical Parsimony Networks and Species Assemblages in Cephalotrichid Nemerteans (Nemertea). <i>PLoS ONE</i> , 2010, 5, e12885.	2.5	70
26	Character-matrix based descriptions of two new nemertean (Nemertea) species. <i>Zoological Journal of the Linnean Society</i> , 2009, 157, 264-294.	2.3	36
27	The first comprehensive molecular phylogeny of Bryozoa (Ectoprocta) based on combined analyses of nuclear and mitochondrial genes. <i>Molecular Phylogenetics and Evolution</i> , 2009, 52, 225-233.	2.7	70
28	The complete mitochondrial genome of <i>Cephalothrix simula</i> (Iwata) (Nemertea: Palaeonemertea). <i>Gene</i> , 2009, 442, 8-17.	2.2	15
29	Intraspecific variation in <i>Tetrastremma laminariae</i> (Nemertini): an examination of one of its possible causes. <i>Journal of Zoology</i> , 2009, 192, 137-141.	1.7	6
30	Global diversity of nemerteans (Nemertea) in freshwater. <i>Hydrobiologia</i> , 2008, 595, 61-66.	2.0	19
31	Phylogeny and classification of the Old World Emberizini (Aves, Passeriformes). <i>Molecular Phylogenetics and Evolution</i> , 2008, 47, 960-973.	2.7	53
32	What is proper vouchering in phylogenetic studies of birds?—A reply to Peterson et al. (2007). <i>Molecular Phylogenetics and Evolution</i> , 2008, 48, 383-385.	2.7	1
33	Species delimitation based on multiple criteria: the Spotted Bush Warbler <i>&lt; i&gt;Bradypterus thoracicus&lt;/i&gt;</i> complex (Aves: Megaluridae). <i>Zoological Journal of the Linnean Society</i> , 2008, 154, 291-307.	2.3	53
34	The phylogenetic relationships between Amphinomidae, Archinomidae and Euphosinidae (Amphinomida: Aciculata: Polychaeta), inferred from molecular data. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2008, 88, 509-513.	0.8	21
35	Checklist of Nemertean Genera and Species Published between 1995 and 2007. <i>Species Diversity</i> , 2008, 13, 245-274.	0.4	98
36	Genetics do not reflect habitat differences in <i>Riseriellus occultus</i> (Heteronemertea, Nemertea) from Spain and Wales. <i>Marine Biology Research</i> , 2007, 3, 117-122.	0.7	10

#	ARTICLE	IF	CITATIONS
37	Swedish nemerteans (phylum Nemertea), with description of a new hoplonemertean genus and species. Journal of Natural History, 2007, 41, 2287-2299.	0.5	12
38	< i>Annulonemertes</i> (phylum Nemertea): when segments do not count. Biology Letters, 2007, 3, 570-573.	2.3	11
39	Phylogeny of benthic Phyllodocidae (Polychaeta) based on morphological and molecular data. Molecular Phylogenetics and Evolution, 2007, 45, 261-271.	2.7	22
40	Morphological, vocal and genetic divergence in the< i>Cettia acanthizoides</i> complex (Aves: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 622	2.3	19
41	BUILD-UP OF THE HIMALAYAN AVIFAUNA THROUGH IMMIGRATION: A BIOGEOGRAPHICAL ANALYSIS OF THEPHYLLOSCOPUSANDSEICERCUSWARBLERS. Evolution; International Journal of Organic Evolution, 2007, 61, 324-333.	2.3	100
42	Global diversity of nemerteans (Nemertea) in freshwater. , 2007, , 61-66.		0
43	Phylogeny and classification of the avian superfamily Sylvioidea. Molecular Phylogenetics and Evolution, 2006, 38, 381-397.	2.7	143
44	Phylogeography of Indonesian and Sino-Himalayan region bush warblers (Cettia, Aves). Molecular Phylogenetics and Evolution, 2006, 41, 556-565.	2.7	20
45	Non-monophyletic taxa and cryptic speciesâ€”Evidence from a molecular phylogeny of leaf-warblers (Phylloscopus, Aves). Molecular Phylogenetics and Evolution, 2005, 36, 261-276.	2.7	98
46	Genus Tetrastemma Ehrenberg, 1831 (Phylum Nemertea)â€”A natural group? Phylogenetic relationships inferred from partial 18S rRNA sequences. Molecular Phylogenetics and Evolution, 2005, 37, 144-152.	2.7	33
47	Phylogeny of Aphroditiformia (Polychaeta) based on molecular and morphological data. Molecular Phylogenetics and Evolution, 2005, 37, 494-502.	2.7	48
48	Phylogenetic position of Nerillidae and Aberranta (Polychaeta, Annelida), analysed by direct optimization of combined molecular and morphological data. Zoologica Scripta, 2005, 34, 313-328.	1.7	36
49	Delimiting species in the hoplonemertean genus Tetrastemma (phylum Nemertea): morphology is not concordant with phylogeny as evidenced from mtDNA sequences. Biological Journal of the Linnean Society, 2005, 86, 201-212.	1.6	47
50	GenusBaseodiscus(Nemertea: Heteronemertea): Molecular identification of a new species in a phylogenetic context. Journal of Natural History, 2005, 39, 3785-3793.	0.5	14
51	Non-monophly of the avian genus Seicercus (Aves: Sylviidae) revealed by mitochondrial DNA. Zoologica Scripta, 2004, 33, 501-510.	1.7	29
52	Phylogeny and evolution of reproductive modes in Autolytinae (Syllidae, Annelida). Molecular Phylogenetics and Evolution, 2003, 29, 235-249.	2.7	91
53	Phylogenetic analysis of a group of palaeonemerteans (Nemertea) including two new species from Queensland and the Great Barrier Reef, Australia. Zoologica Scripta, 2003, 32, 279-296.	1.7	18
54	Test of the monophly of Odostomiinae and Turbonilliinae (Gastropoda, Heterobranchia,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 62 Td (P	1.7	18

#	ARTICLE	IF	CITATIONS
55	Some heteronemerteans (Nemertea) from the Solomon Islands. <i>Journal of Natural History</i> , 2002, 36, 1785-1804.	0.5	9
56	Phylogenetic Relationships among Higher Nemertean (Nemertea) Taxa Inferred from 18S rDNA Sequences. <i>Molecular Phylogenetics and Evolution</i> , 2001, 20, 327-334.	2.7	69
57	Molecular Phylogeny of the Model Annelid <i>Ophryotrocha</i> . <i>Biological Bulletin</i> , 2001, 201, 193-203.	1.8	63
58	Useful Characters in Gastropod Phylogeny: Soft Information or Hard Facts. <i>Systematic Biology</i> , 2001, 50, 136-141.	5.6	13
59	Some Nemerteans (Nemertea) from Queensland and the Great Barrier Reef, Australia. <i>Zoological Science</i> , 2001, 18, 1259-1273.	0.7	8
60	Six new species of palaeonemerteans (Nemertea) from Hong Kong. <i>Zoological Journal of the Linnean Society</i> , 1999, 125, 151-196.	2.3	17
61	Taxonomy and Philosophy of Names. <i>Biology and Philosophy</i> , 1998, 13, 233-244.	1.4	27
62	Phylogenetic relationships and genetic distances between some monostiliferous interstitial nemerteans (Otostyphlonemertes, Hoplonemertea, Nemertea) indicated from the 16S rRNA gene. <i>Zoological Journal of the Linnean Society</i> , 1998, 123, 105-115.	2.3	17
63	Molecular Phylogeny of Some European Heteronemertean (Nemertea) Species and the Monophyletic Status of <i>Riseriellus</i> , <i>Lineus</i> , and <i>Micrura</i> . <i>Molecular Phylogenetics and Evolution</i> , 1998, 10, 271-280.	2.7	29
64	There is no support for Jensen's hypothesis of nemerteans as ancestors to the vertebrates. <i>Hydrobiologia</i> , 1997, 365, 47-54.	2.0	7
65	The nemerteans (Nemertea) of Rottnest Island, Western Australia. <i>Zoologica Scripta</i> , 1995, 24, 101-141.	1.7	19
66	Random Amplified Polymorphic Dna (Rapd) and Intraspecific Variation in <i>Oerstedia Dorsalis</i> (Hoplonemertea, Nemertea). <i>Journal of the Marine Biological Association of the United Kingdom</i> , 1995, 75, 483-490.	0.8	15
67	PHYLOGENY OF THE NEMERTEAN SUBCLASS PALAEONEMERTEA (ANOPLA, NEMERTEA). <i>Cladistics</i> , 1994, 10, 347-402.	3.3	38
68	Phylogenetic classification and the definition of taxon names. <i>Zoologica Scripta</i> , 1994, 23, 19-25.	1.7	59
69	Phylogeny of the Nemertean Subclass Palaeonemertea (Anopla, Nemertea). <i>Cladistics</i> , 1994, 10, 347-402.	3.3	2
70	Phylogeny, natural groups and nemertean classification. <i>Hydrobiologia</i> , 1993, 266, 103-113.	2.0	16
71	Phylogeny, natural groups and nemertean classification. , 1993, , 103-113.		6
72	A proposal for renaming the higher taxonomic categories in the phylum Nemertea. <i>Journal of Natural History</i> , 1991, 25, 45-48.	0.5	21

#	ARTICLE	IF	CITATIONS
73	Gibson's reclassification of the enoplans nemerteans (Enopla, Nemertea): a critique and cladistic analysis. <i>Zoologica Scripta</i> , 1990, 19, 133-140.	1.7	33
74	Phylogeny and Cladistic Classification of the Paramonostiliferous Family Plectonemertidae (Phylum) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 536	3.3	
75	Classification of the family Plectonemertidae (Nemertea): a phenetic comparison. <i>Zoological Journal of the Linnean Society</i> , 1989, 97, 57-68.	2.3	10
76	Phylogeny and cladistic classification of terrestrial nemerteans: the genera Pantinonemertes Moore & Gibson and Geonemertes Semper. <i>Zoological Journal of the Linnean Society</i> , 1989, 95, 363-372.	2.3	25
77	A new monostiliferous hoplonemertean (Nemertea), <i>Oerstedia striata</i> sp.n., from the west coast of Sweden. <i>Zoologica Scripta</i> , 1988, 17, 135-139.	1.7	11
78	Polymorphism in <i>Oerstedia dorsalis</i> (Abildgaard, 1806) revisited. <i>Hydrobiologia</i> , 1988, 156, 93-98.	2.0	20
79	Polymorphism in <i>Oerstedia dorsalis</i> (Abildgaard, 1806) revisited. , 1988, , 93-98.		3
80	A Possible Mechanism for the Evolution of Aposematic Colouration in Solitary Nemerteans (Phylum) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 527	2.7	27
81	A previously unrecognized report of a nemertean in the literature. <i>Archives of Natural History</i> , 1986, 13, 7-8.	0.3	9
82	Nemertean systematics and phenetic classification: an example from a group of hoplonemerteans. <i>Zoological Journal of the Linnean Society</i> , 1985, 85, 247-266.	2.3	5
83	Multivariate analysis of polymorphism in the hoplonemertean <i>Oerstedia dorsalis</i> (Abildgaard, 1806). <i>Journal of Experimental Marine Biology and Ecology</i> , 1984, 78, 1-22.	1.5	25
84	Tubulanus annulatus, an aposematic nemertean?. <i>Biological Journal of the Linnean Society</i> , 1979, 12, 177-179.	1.6	10
85	Statistical analysis of variation in characters in <i>Tetrastemma laminariae</i> (Nemertini), with a redescription of the species. <i>Journal of Zoology</i> , 1979, 189, 39-56.	1.7	40
86	Littoral nemerteans from the Faroe Islands. <i>Sarsia</i> , 1977, 63, 185-190.	0.5	2
87	Polymorphism hides cryptic species in <i>Oerstedia dorsalis</i> (Nemertea, Hoplonemertea). <i>Biological Journal of the Linnean Society</i> , 0, 98, 556-567.	1.6	22