Jingling Jin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11381708/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	NKT Cells Coexpressing a GD2-Specific Chimeric Antigen Receptor and IL15 Show Enhanced <i>In Vivo</i> Persistence and Antitumor Activity against Neuroblastoma. Clinical Cancer Research, 2019, 25, 7126-7138.	7.0	112
2	Small molecule inhibitor agerafenib effectively suppresses neuroblastoma tumor growth in mouse models via inhibiting ERK MAPK signaling. Cancer Letters, 2019, 457, 129-141.	7.2	16
3	Targeting LRH‑1 in hepatoblastoma cell lines causes decreased proliferation. Oncology Reports, 2018, 41, 143-153.	2.6	14
4	Methylation of PCDH19 predicts poor prognosis of hepatocellular carcinoma. Asia-Pacific Journal of Clinical Oncology, 2018, 14, e352-e358.	1.1	19
5	IL-21 Selectively Protects CD62L+ NKT Cells and Enhances Their Effector Functions for Adoptive Immunotherapy. Journal of Immunology, 2018, 201, 2141-2153.	0.8	40
6	The second-generation ALK inhibitor alectinib effectively induces apoptosis in human neuroblastoma cells and inhibits tumor growth in a TH-MYCN transgenic neuroblastoma mouse model. Cancer Letters, 2017, 400, 61-68.	7.2	37
7	A Novel Cell Line Based Orthotopic Xenograft Mouse Model That Recapitulates Human Hepatoblastoma. Scientific Reports, 2017, 7, 17751.	3.3	29
8	EWS-FLI1 and RNA helicase A interaction inhibitor YK-4-279 inhibits growth of neuroblastoma. Oncotarget, 2017, 8, 94780-94792.	1.8	5
9	Activation of CDK4 Triggers Development of Non-alcoholic Fatty Liver Disease. Cell Reports, 2016, 16, 744-756.	6.4	41
10	Cooperation of C/EBP family proteins and chromatin remodeling proteins is essential for termination of liver regeneration. Hepatology, 2015, 61, 315-325.	7.3	56
11	Age-associated Change of C/EBP Family Proteins Causes Severe Liver Injury and Acceleration of Liver Proliferation after CCl4 Treatments. Journal of Biological Chemistry, 2014, 289, 1106-1118.	3.4	30
12	Farnesoid X receptor inhibits gankyrin in mouse livers and prevents development of liver cancer. Hepatology, 2013, 57, 1098-1106.	7.3	61
13	Increased Expression of Enzymes of Triglyceride Synthesis Is Essential for the Development of Hepatic Steatosis. Cell Reports, 2013, 3, 831-843.	6.4	63
14	Farnesoid X receptor directly regulates xenobiotic detoxification genes in the long-lived Little mice. Mechanisms of Ageing and Development, 2013, 134, 407-415.	4.6	17
15	Transcriptional and Translational Regulation of C/EBPÎ ² -HDAC1 Protein Complexes Controls Different Levels of p53, SIRT1, and PGC1α Proteins at the Early and Late Stages of Liver Cancer. Journal of Biological Chemistry, 2013, 288, 14451-14462.	3.4	38
16	The reduction of SIRT1 in livers of old mice leads to impaired body homeostasis and to inhibition of liver proliferation. Hepatology, 2011, 54, 989-998.	7.3	87
17	Epigenetic changes play critical role in ageâ€associated dysfunctions of the liver. Aging Cell, 2010, 9, 895-910	6.7	30
18	Elimination of C/EBPα through the ubiquitin-proteasome system promotes the development of liver cancer in mice. Journal of Clinical Investigation, 2010, 120, 2549-2562.	8.2	50

JINGLING JIN

#	Article	IF	CITATIONS
19	The Age-Associated Decline of Glycogen Synthase Kinase 3β Plays a Critical Role in the Inhibition of Liver Regeneration. Molecular and Cellular Biology, 2009, 29, 3867-3880.	2.3	60
20	GSK3β-cyclin D3-CUGBP1-elF2 pathway in aging and in Myotonic Dystrophy. Cell Cycle, 2009, 8, 2356-2359.	2.6	22
21	GSK3β and aging liver. Aging, 2009, 1, 582-585.	3.1	27
22	Cloning and expression profiling of testis-expressed piRNA-like RNAs. Rna, 2007, 13, 1693-1702.	3.5	70
23	A PCR-based method for detection and quantification of small RNAs. Biochemical and Biophysical Research Communications, 2006, 351, 756-763.	2.1	153