

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11382118/publications.pdf>

Version: 2024-02-01

73
papers

7,663
citations

53794
45
h-index

91884
69
g-index

73
all docs

73
docs citations

73
times ranked

3393
citing authors

#	ARTICLE	IF	CITATIONS
1	Basic and clinical aspects of visceral hyperalgesia. Gastroenterology, 1994, 107, 271-293.	1.3	875
2	Colorectal distension as a noxious visceral stimulus: physiologic and pharmacologic characterization of pseudoeffective reflexes in the rat. Brain Research, 1988, 450, 153-169.	2.2	636
3	Descending modulation of pain. Neuroscience and Biobehavioral Reviews, 2004, 27, 729-737.	6.1	503
4	Nitric oxide mediates the thermal hyperalgesia produced in a model of neuropathic pain in the rat. Neuroscience, 1992, 50, 7-10.	2.3	358
5	The possible role of glia in nociceptive processing and hyperalgesia in the spinal cord of the rat. Neuropharmacology, 1994, 33, 1471-1478.	4.1	319
6	Vagal afferent modulation of nociception. Brain Research Reviews, 1992, 17, 77-99.	9.0	268
7	Production of endogenous nitric oxide and activation of soluble guanylate cyclase are required for N-methyl-D-aspartate-produced facilitation of the nociceptive tail-flick reflex. European Journal of Pharmacology, 1992, 214, 93-96.	3.5	254
8	Relative contributions of the nucleus raphe magnus and adjacent medullary reticular formation to the inhibition by stimulation in the periaqueductal gray of a spinal nociceptive reflex in the pentobarbital-anesthetized rat. Brain Research, 1984, 305, 77-87.	2.2	248
9	Evaluation of the periaqueductal central gray (PAG) as a morphine-specific locus of action and examination of morphine-induced and stimulation-produced analgesia at coincident PAG loci. Brain Research, 1977, 124, 283-303.	2.2	232
10	The role of nitric oxide in the development and maintenance of the hyperalgesia produced by intraplantar injection of carrageenan in the rat. Neuroscience, 1994, 60, 367-374.	2.3	225
11	Characterization of coeruleospinal inhibition of the nociceptive tail-flick reflex in the rat: Mediation by spinal $\text{I}\pm 2$ -adrenoceptors. Brain Research, 1986, 364, 315-330.	2.2	193
12	Characterization of inhibition of a spinal nociceptive reflex by stimulation medially and laterally in the midbrain and medulla in the pentobarbital-anesthetized rat. Brain Research, 1984, 305, 67-76.	2.2	191
13	Neural Upregulation in Interstitial Cystitis. Urology, 2007, 69, S24-S33.	1.0	171
14	Intracolonic zymosan produces visceral hyperalgesia in the rat that is mediated by spinal NMDA and non-NMDA receptors. Brain Research, 1996, 736, 7-15.	2.2	159
15	Differential c-Fos expression in the nucleus of the solitary tract and spinal cord following noxious gastric distention in the rat. Neuroscience, 1996, 74, 873-884.	2.3	148
16	Spinal serotonin receptors mediate descending facilitation of a nociceptive reflex from the nuclei reticularis gigantocellularis and gigantocellularis pars alpha in the rat. Brain Research, 1991, 550, 35-48.	2.2	147
17	Stimulation-produced spinal inhibition from the midbrain in the rat is mediated by an excitatory amino acid neurotransmitter in the medial medulla. Journal of Neuroscience, 1986, 6, 1803-1813.	3.6	134
18	Characterization of inhibition of the spinal nociceptive tail-flick reflex in the rat from the medullary lateral reticular nucleus. Journal of Neuroscience, 1986, 6, 701-713.	3.6	124

#	ARTICLE	IF	CITATIONS
19	Evidence that spinal 5-HT1, 5-HT2 and 5-HT3 receptor subtypes modulate responses to noxious colorectal distension in the rat. Brain Research, 1991, 538, 64-75.	2.2	108
20	Further behavioral evidence that colorectal distension is a "noxious" visceral stimulus in rats. Neuroscience Letters, 1991, 131, 113-116.	2.1	103
21	Intraplantar zymosan as a reliable, quantifiable model of thermal and mechanical hyperalgesia in the rat. European Journal of Pain, 1997, 1, 43-52.	2.8	100
22	Vagal afferent modulation of a nociceptive reflex in rats: involvement of spinal opioid and monoamine receptors. Brain Research, 1988, 446, 285-294.	2.2	97
23	Acute thermal hyperalgesia in the rat is produced by activation of N-methyl-d-aspartate receptors and protein kinase c and production of nitric oxide. Neuroscience, 1996, 71, 327-335.	2.3	82
24	Noxious distention of viscera results in differential c-Fos expression in second order sensory neurons receiving "sympathetic" or "parasympathetic" input. Neuroscience Letters, 1994, 180, 71-75.	2.1	80
25	Spinal cholinergic and monoaminergic receptors mediate descending inhibition from the nuclei reticularis gigantocellularis and gigantocellularis pars alpha in the rat. Brain Research, 1990, 535, 67-78.	2.2	78
26	Brain-stem relays mediating stimulation-produced antinociception from the lateral hypothalamus in the rat. Journal of Neuroscience, 1988, 8, 2652-2663.	3.6	73
27	Role of neurokinin 3 receptors on responses to colorectal distention in the rat: Electrophysiological and behavioral studies. Gastroenterology, 1999, 116, 1124-1131.	1.3	73
28	Effects of intracolonic acetic acid on responses to colorectal distension in the rat. Brain Research, 1995, 672, 77-82.	2.2	70
29	An evaluation of stimulation-produced analgesia in the cat. Experimental Neurology, 1978, 62, 570-579.	4.1	69
30	Antinociception and cardiovascular responses produced by intravenous morphine: the role of vagal afferents. Brain Research, 1991, 543, 256-270.	2.2	69
31	Effect of spinal norepinephrine depletion on descending inhibition of the tail flick reflex from the locus coeruleus and lateral reticular nucleus in the rat. Brain Research, 1987, 400, 40-52.	2.2	64
32	Characterization of the role of spinal n-methyl-d-aspartate receptors in thermal nociception in the rat. Neuroscience, 1993, 57, 385-395.	2.3	61
33	Peripheral and Central P2X3 Receptor Contributions to Colon Mechanosensitivity and Hypersensitivity in the Mouse. Gastroenterology, 2009, 137, 2096-2104.	1.3	61
34	A distension control device useful for quantitative studies of hollow organ sensation. Physiology and Behavior, 1987, 41, 635-638.	2.1	57
35	Spinal cord NADPH-diaphorase histochemical staining but not nitric oxide synthase immunoreactivity increases following carrageenan-produced hindpaw inflammation in the rat. Brain Research, 1994, 668, 204-210.	2.2	57
36	Spinal Mediators of Hyperalgesia. Drugs, 1994, 47, 10-20.	10.9	57

#	ARTICLE	IF	CITATIONS
37	NMDA and quisqualate modulation of visceral nociception in the rat. Brain Research, 1994, 651, 215-226.	2.2	55
38	N-methyl-d-aspartate receptor-mediated changes in thermal nociception: Allosteric modulation at glycine and polyamine recognition sites. Neuroscience, 1994, 63, 925-936.	2.3	53
39	Characterization of superficial T13-L2 dorsal horn neurons encoding for colorectal distension in the rat: comparison with neurons in deep laminae. Brain Research, 1989, 486, 301-309.	2.2	52
40	Brainstem and spinal pathways mediating descending inhibition from the medullary lateral reticular nucleus in the rat. Brain Research, 1988, 440, 109-122.	2.2	51
41	Spinal monoaminergic receptors mediate the antinociception produced by glutamate in the medullary lateral reticular nucleus. Journal of Neuroscience, 1987, 7, 2862-2873.	3.6	50
42	Spinal monoamine mediation of stimulation-produced antinociception from the lateral hypothalamus. Brain Research, 1987, 403, 290-300.	2.2	49
43	Characterization of antinociception produced by glutamate microinjection in the nucleus tractus solitarius and the nucleus reticularis ventralis. Journal of Neuroscience, 1988, 8, 4675-4684.	3.6	49
44	Opioid, cholinergic and $\hat{1}$ -adrenergic influences on the modulation of nociception from the lateral reticular nucleus of the rat. Brain Research, 1986, 384, 282-293.	2.2	48
45	The peripheral nociceptive actions of intravenously administered 5-HT in the rat requires dual activation of both 5-HT ₂ and 5-HT ₃ receptor subtypes. Brain Research, 1991, 561, 61-68.	2.2	48
46	Vagal afferent-mediated inhibition of a nociceptive reflex by intravenous serotonin i the rat. I. Characterization. Brain Research, 1990, 524, 90-100.	2.2	46
47	NADPH-diaphorase histochemistry provides evidence for a bilateral, somatotopically inappropriate response to unilateral hindpaw inflammation in the rat. Brain Research, 1994, 647, 113-123.	2.2	44
48	Activation of lamina I spinal cord neurons that express the substance P receptor in visceral nociception and hyperalgesia. Journal of Pain, 2002, 3, 3-11.	1.4	44
49	Lesion in nucleus reticularis gigantocellularis: effect on the antinociception produced by micro-injection of morphine and focal electrical stimulation in the periaqueductal gray matter. Brain Research, 1982, 231, 143-152.	2.2	42
50	Is there a role for an endothelium-derived relaxing factor in nociception?. Brain Research, 1990, 531, 342-345.	2.2	42
51	Locus coeruleus lesions in the rat enhance the antinociceptive potency of centrally administered clonidine but not morphine. Brain Research, 1985, 341, 320-330.	2.2	40
52	Attenuation of c-Fos expression in the rat lumbosacral spinal cord by morphine or tramadol following noxious colorectal distention. Brain Research, 1995, 701, 175-182.	2.2	40
53	Hind Paw Incision in the Rat Produces Long-Lasting Colon Hypersensitivity. Journal of Pain, 2008, 9, 246-253.	1.4	33
54	Responses of primary afferents and spinal dorsal horn neurons to thermal and mechanical stimuli before and during zymosan-induced inflammation of the rat hindpaw. Brain Research, 1997, 772, 135-148.	2.2	30

#	ARTICLE	IF	CITATIONS
55	Intrathecal coadministration of clonidine with serotonin receptor agonists produces supra-additive visceral antinociception in the rat. Brain Research, 1991, 555, 35-42.	2.2	28
56	The role of CNS NMDA receptors and nitric oxide in visceral hyperalgesia. European Journal of Pharmacology, 2001, 429, 319-325.	3.5	28
57	Dissociation of antinociceptive from cardiovascular effects of stimulation in the lateral reticular nucleus in the rat. Brain Research, 1987, 405, 140-149.	2.2	26
58	Chapter 14 Acute mechanical hyperalgesia in the rat can be produced by coactivation of spinal ionotropic AMPA and metabotropic glutamate receptors, activation of phospholipase A2 and generation of cyclooxygenase products. Progress in Brain Research, 1996, 110, 177-192.	1.4	26
59	Chapter 6. Visceral polymodal receptors. Progress in Brain Research, 1996, 113, 101-112.	1.4	26
60	Chapter 27 The glutamate synapse: A target in the pharmacological management of hyperalgesic pain states. Progress in Brain Research, 1998, 116, 407-420.	1.4	25
61	Expression of nitric oxide synthase type II in the spinal cord under conditions producing thermal hyperalgesia. Journal of Chemical Neuroanatomy, 1996, 10, 221-229.	2.1	24
62	Role of sensitized pelvic nerve afferents from the inflamed rat colon in the maintenance of visceral hyperalgesia. Progress in Brain Research, 2000, 129, 375-387.	1.4	21
63	Neonatal capsaicin treatment abolishes the nociceptive responses to intravenous 5-HT in the rat. Brain Research, 1991, 542, 212-218.	2.2	20
64	Medullary substrates of descending spinal inhibition activated by intravenous administration of [d-Ala2]methionine enkephalinamide in the rat. Brain Research, 1987, 411, 236-247.	2.2	18
65	ATTENUATION OF PETHIDINE-INDUCED ANTINOCICEPTION BY ZIMELIDINE, AN INHIBITOR OF 5-HYDROXYTRYPTAMINE REUPTAKE. British Journal of Pharmacology, 1980, 70, 411-414.	5.4	13
66	It's Chickens and Eggs All Over Again: Is Central Reorganization the Result or Cause of Persistent Visceral Pain?. Gastroenterology, 2007, 132, 1618-1620.	1.3	13
67	Ethanol dose-dependently attenuates NMDA-mediated thermal hyperalgesia in the rat. Neuroscience Letters, 1993, 154, 137-140.	2.1	11
68	Visceral pain: basic mechanisms. , 2006, , 721-736.		9
69	Models of Visceral Pain: Colorectal Distension (CRD). Current Protocols in Pharmacology, 2004, 25, Unit 5.36.	4.0	6
70	Production of Reversible Local Blockage of Neuronal Function. Methods in Neurosciences, 1991, , 122-138.	0.5	5
71	Visceral Pain. , 2008, , 543-569.		3
72	Dolor visceral. , 2007, , 741-758.		1

#	ARTICLE	IF	CITATIONS
73	Visceral Pain. , 2009, , 189-194.		0