

Aya Kadota

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1138431/publications.pdf>

Version: 2024-02-01

145
papers

2,986
citations

236925

25
h-index

206112

48
g-index

150
all docs

150
docs citations

150
times ranked

5098
citing authors

#	ARTICLE	IF	CITATIONS
1	Cardiovascular Disease and Risk Factors in Asia. <i>Circulation</i> , 2008, 118, 2702-2709.	1.6	604
2	Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. <i>Nature Genetics</i> , 2019, 51, 379-386.	21.4	164
3	Brachialâ€“ankle pulse wave velocity predicts all-cause mortality in the general population: findings from the Takashima study, Japan. <i>Hypertension Research</i> , 2010, 33, 922-925.	2.7	125
4	Relationship Between Metabolic Risk Factor Clustering and Cardiovascular Mortality Stratified by High Blood Glucose and Obesity. <i>Diabetes Care</i> , 2007, 30, 1533-1538.	8.6	120
5	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019, 24, 1.	3.4	84
6	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population. <i>Journal of Hypertension</i> , 2017, 35, 401-408.	0.5	74
7	Impact of Metabolic Syndrome on the Risk of Cardiovascular Disease Mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014, 113, 84-89.	1.6	69
8	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014, 112, 916-924.	2.3	59
9	Increased Aortic Calcification Is Associated With Arterial Stiffness Progression in Multiethnic Middle-Aged Men. <i>Hypertension</i> , 2017, 69, 102-108.	2.7	51
10	Epidemiology of hypertension in Japan: beyond the new 2019 Japanese guidelines. <i>Hypertension Research</i> , 2020, 43, 1344-1351.	2.7	49
11	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016, 23, 800-809.	2.0	48
12	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016, 246, 141-147.	0.8	48
13	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, 1703-1708.	2.4	44
14	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 755-766.	2.0	43
15	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980â€“2010. <i>Hypertension Research</i> , 2015, 38, 790-795.	2.7	39
16	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016, 5, .	3.7	39
17	Effects of Stroke Education of Junior High School Students on Stroke Knowledge of Their Parents. <i>Stroke</i> , 2015, 46, 572-574.	2.0	37
18	Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. <i>Medicine and Science in Sports and Exercise</i> , 2018, 50, 2433-2441.	0.4	36

#	ARTICLE	IF	CITATIONS
19	Does High-Sensitivity C-Reactive Protein or Low-Density Lipoprotein Cholesterol Show a Stronger Relationship with the Cardio-Ankle Vascular Index in Healthy Community Dwellers?: the KOBE Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 1027-1034.	2.0	35
20	Genome-wide association meta-analysis identifies GP2 gene risk variants for pancreatic cancer. <i>Nature Communications</i> , 2020, 11, 3175.	12.8	34
21	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , 2017, 117, 260-266.	2.3	31
22	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016, 4, 225-232.	1.8	30
23	Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. <i>International Journal of Cardiology</i> , 2018, 267, 177-182.	1.7	30
24	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016, 220, 262-267.	1.7	29
25	The National Integrated Project for Prospective Observation of Non-communicable Disease and its Trends in the Aged 2010 (NIPPON DATA2010): Objectives, Design, and Population Characteristics. <i>Journal of Epidemiology</i> , 2018, 28, S2-S9.	2.4	29
26	Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. <i>Atherosclerosis</i> , 2018, 273, 145-152.	0.8	27
27	Relationship Between Socioeconomic Status and the Prevalence of Underweight, Overweight or Obesity in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S10-S16.	2.4	26
28	Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. <i>International Journal of Cardiology</i> , 2015, 189, 67-72.	1.7	24
29	Associations of Nutrient Patterns with the Prevalence of Metabolic Syndrome: Results from the Baseline Data of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2019, 11, 990.	4.1	24
30	Association of Dietary Acid Load with the Prevalence of Metabolic Syndrome among Participants in Baseline Survey of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2020, 12, 1605.	4.1	23
31	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014, 236, 237-243.	0.8	22
32	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 783-792.	2.9	22
33	Re-evaluation of the associations of egg intake with serum total cholesterol and cause-specific and total mortality in Japanese women. <i>European Journal of Clinical Nutrition</i> , 2018, 72, 841-847.	2.9	21
34	Twelve-year trends of increasing overweight and obesity in patients with diabetes: the Shiga Diabetes Clinical Survey. <i>Endocrine Journal</i> , 2018, 65, 527-536.	1.6	21
35	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. <i>Hypertension Research</i> , 2019, 42, 411-418.	2.7	21
36	Red blood cell fatty acid patterns from 7 countries: Focus on the Omega-3 index. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2022, 179, 102418.	2.2	21

#	ARTICLE	IF	CITATIONS
37	Underweight Young Women Without Later Weight Gain Are at High Risk for Osteopenia After Midlife: The KOBE Study. <i>Journal of Epidemiology</i> , 2016, 26, 572-578.	2.4	20
38	Macronutrient Intake and Socioeconomic Status: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S17-S22.	2.4	19
39	Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. <i>Stroke</i> , 2020, 51, 3584-3591.	2.0	19
40	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015, 239, 444-450.	0.8	18
41	Dietary Inflammatory Index Positively Associated With High-Sensitivity C-Reactive Protein Level in Japanese From NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2020, 30, 98-107.	2.4	18
42	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 606-620.	2.0	17
43	Socioeconomic Status and Knowledge of Cardiovascular Risk Factors: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S46-S52.	2.4	17
44	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 198-206.	2.0	17
45	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 67-76.	2.9	17
46	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , 2016, 213, 460-467.	7.5	16
47	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S29-S34.	2.4	15
48	Association between plasma levels of homocysteine, folate, and vitamin B12, and dietary folate intake and hypertension in a cross-sectional study. <i>Scientific Reports</i> , 2020, 10, 18499.	3.3	14
49	Identification of a novel uterine leiomyoma GWAS locus in a Japanese population. <i>Scientific Reports</i> , 2020, 10, 1197.	3.3	14
50	Sedentary Time is Associated with Cardiometabolic Diseases in A Large Japanese Population: A Cross-Sectional Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1097-1107.	2.0	14
51	Differences Between Coronary Artery Calcification and Aortic Artery Calcification in Relation to Cardiovascular Disease Risk Factors in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 452-464.	2.0	13
52	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1195-1202.e1.	1.5	12
53	Relationships among Socioeconomic Factors and Self-rated Health in Japanese Adults: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S66-S72.	2.4	12
54	The relationship between serum levels of LOX-1 ligand containing ApoAI as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. <i>Atherosclerosis</i> , 2020, 313, 20-25.	0.8	12

#	ARTICLE	IF	CITATIONS
55	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 995-1009.	2.0	12
56	Relationships between Family Histories of Stroke and of Hypertension and Stroke Mortality: NIPPON DATA80, 1980-1999. <i>Hypertension Research</i> , 2008, 31, 1525-1531.	2.7	11
57	Relationship of moderate metabolic risk factor clustering to cardiovascular disease mortality in non-lean Japanese: A 15-year follow-up of NIPPON DATA90. <i>Atherosclerosis</i> , 2011, 215, 209-213.	0.8	11
58	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: A 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014, 64, 218-224.	1.9	11
59	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia. <i>Journal of Clinical Lipidology</i> , 2016, 10, 172-180.e1.	1.5	11
60	Genetic Variants of <i>RAMP2</i> and <i>CLR</i> are Associated with Stroke. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 1267-1281.	2.0	11
61	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S59-S65.	2.4	11
62	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018, 82, 2542-2548.	1.6	11
63	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan. NIPPON DATA80. <i>Circulation Journal</i> , 2019, 83, 1254-1260.	1.6	11
64	Socioeconomic and lifestyle factors associated with depressive tendencies in general Japanese men and women: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019, 24, 37.	3.4	11
65	The interaction between ABCA1 polymorphism and physical activity on the HDL-cholesterol levels in a Japanese population. <i>Journal of Lipid Research</i> , 2020, 61, 86-94.	4.2	11
66	Relationship of household salt intake level with long-term all-cause and cardiovascular disease mortality in Japan: NIPPON DATA80. <i>Hypertension Research</i> , 2020, 43, 132-139.	2.7	11
67	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018, 37, 182-188.	5.0	10
68	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S23-S28.	2.4	10
69	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020, 30, 244-252.	2.4	10
70	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , 2016, 252, 116-121.	0.8	9
71	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , 2018, 16, 166-173.	1.3	9
72	Comparison of carotid plaque burden among healthy middle-aged men living in the US, Japan, and South Korea. <i>International Journal of Cardiology</i> , 2018, 266, 245-249.	1.7	9

#	ARTICLE	IF	CITATIONS
73	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019, 29, 133-138.	2.4	9
74	Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. <i>Journal of Epidemiology</i> , 2020, 30, 121-127.	2.4	9
75	Estimated 24 h Urinary Sodium-to-Potassium Ratio Is Related to Renal Function Decline: A 6-Year Cohort Study of Japanese Urban Residents. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 5811.	2.6	9
76	Impact of <i>PSCA</i> Polymorphisms on the Risk of Duodenal Ulcer. <i>Journal of Epidemiology</i> , 2021, 31, 12-20.	2.4	9
77	Relationship Between Non-fasting Triglycerides and Cardiovascular Disease Mortality in a 20-year Follow-up Study of a Japanese General Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2022, 32, 303-313.	2.4	9
78	Effect of Underlying Cardiometabolic Diseases on the Association Between Sedentary Time and All-Cause Mortality in a Large Japanese Population: A Cohort Analysis Based on the J-MICC Study. <i>Journal of the American Heart Association</i> , 2021, 10, e018293.	3.7	9
79	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , 2019, 29, 205-212.	2.4	9
80	Relationship between Kidney Function and Subclinical Atherosclerosis Progression Evaluated by Coronary Artery Calcification. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 1359-1371.	2.0	9
81	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , 2016, 39, 670-679.	2.7	8
82	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , 2017, 24, 505-513.	1.8	8
83	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018, 18, 799-805.	1.5	8
84	Factors Related to Participation in Health Examinations for Japanese National Health Insurance: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S53-S58.	2.4	8
85	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 417-423.	2.3	8
86	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013, 99, 1024-1029.	2.9	7
87	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men: A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , 2019, 83, 1506-1513.	1.6	7
88	One-year weight loss maintenance outcomes following a worksite-based weight reduction program among Japanese men with cardiovascular risk factors. <i>Journal of Occupational Health</i> , 2019, 61, 189-196.	2.1	7
89	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019, 37, 1676-1681.	0.5	7
90	Alcohol consumption and cognitive function in elderly Japanese men. <i>Alcohol</i> , 2020, 85, 145-152.	1.7	7

#	ARTICLE	IF	CITATIONS
91	Smoking habits and progression of coronary and aortic artery calcification: A 5-year follow-up of community-dwelling Japanese men. <i>International Journal of Cardiology</i> , 2020, 314, 89-94.	1.7	7
92	Relationship of Higher-level Functional Capacity With Long-term Mortality in Japanese Older People: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2023, 33, 136-141.	2.4	7
93	A Comparison of Segment-Specific and Composite Measures of Carotid Intima-Media Thickness and their Relationships with Coronary Calcium. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 282-295.	2.0	7
94	Trends in medical performance in diabetic patients in primary care clinics compared with those in hospitals: Shiga Diabetes Clinical Survey, Japan, 2000â€“2012. <i>Diabetology International</i> , 2017, 8, 59-68.	1.4	6
95	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , 2021, 44, 80-87.	2.7	6
96	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , 2021, 90, 57-65.	1.7	6
97	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. <i>British Journal of Nutrition</i> , 2021, 126, 1843-1851.	2.3	6
98	Self-reported Sleep Duration and Subclinical Atherosclerosis in a General Population of Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 186-198.	2.0	5
99	Risk Factors That Most Accurately Predict Coronary Artery Disease Based on the Duration of Follow-upâ€“â€• NIPPON DATA80 â€•. <i>Circulation Journal</i> , 2021, 85, 908-913.	1.6	5
100	Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. <i>PLoS ONE</i> , 2021, 16, e0254706.	2.5	5
101	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 252, 253-262.	1.2	5
102	Association of Coronary Artery Calcification with Estimated Coronary Heart Disease Risk from Prediction Models in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 477-489.	2.0	4
103	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S40-S45.	2.4	4
104	Time-Related Changes in Relationships Between the Keys Score, Dietary Lipids, and Serum Total Cholesterol in Japanâ€“â€• NIPPON DATA80/90/2010 â€•. <i>Circulation Journal</i> , 2018, 83, 147-155.	1.6	4
105	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical) <i>TJ ETQq1 1 0.784314 rgB2/@verlock 10 Tf 50 1</i>		
106	Association of Red Meat Intake with the Risk of Cardiovascular Mortality in General Japanese Stratified by Kidney Function: NIPPON DATA80. <i>Nutrients</i> , 2020, 12, 3707.	4.1	4
107	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , 2016, 11, e0157563.	2.5	4
108	Eating Slowly Is Associated with Undernutrition among Community-Dwelling Adult Men and Older Adult Women. <i>Nutrients</i> , 2022, 14, 54.	4.1	4

#	ARTICLE	IF	CITATIONS
109	Serum long-chain n-3 polyunsaturated fatty acids and aortic calcification in middle-aged men: The population-based cross-sectional ERA-JUMP study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019, 29, 837-846.	2.6	3
110	Anthropometric Obesity Indices were Stronger than CT-Based Indices in Associations with Carotid Intima-Media Thickness in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 1102-1114.	2.0	3
111	Cross-sectional association of bone mineral density with coronary artery calcification in an international multi-ethnic population-based cohort of men aged 40â€“49: ERA JUMP study. <i>IJC Heart and Vasculature</i> , 2020, 30, 100618.	1.1	3
112	Relationship Between Calcium Intake and Impaired Activities of Daily Living in a Japanese Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2021, 31, 119-124.	2.4	3
113	Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 57.	3.4	3
114	Association between Stress-Coping Strategy and Functional Disability in the General Older Adult Population: The Takashima Study. <i>Gerontology</i> , 2022, 68, 699-706.	2.8	3
115	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. <i>Journal of Nutritional Science</i> , 2021, 10, e61.	1.9	3
116	Physique at Birth and Cardiovascular Disease Risk Factors in Japanese Urban Residents: the KOBE Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, , .	2.0	3
117	Ventricular Premature Complexes and Their Associated Factors in a General Population of Japanese Men. <i>American Journal of Cardiology</i> , 2022, 169, 51-56.	1.6	3
118	Association of equol producing status with aortic calcification in middle-aged Japanese men: The ERA JUMP study. <i>International Journal of Cardiology</i> , 2022, 352, 158-164.	1.7	3
119	Relationship of leisure-time and household physical activity level and type with cardiovascular disease: secondary analysis of the Takashima Study data. <i>BMC Cardiovascular Disorders</i> , 2022, 22, 132.	1.7	3
120	Trends in Prevalence, Treatment, and Control of Hypertension According to 40-Year-Old Life Expectancy at Prefectures in Japan from the National Health and Nutrition Surveys. <i>Nutrients</i> , 2022, 14, 1219.	4.1	3
121	Influence of cigarette smoking and inflammatory gene polymorphisms on glycated hemoglobin in the Japanese general population. <i>Preventive Medicine Reports</i> , 2016, 3, 288-295.	1.8	2
122	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S35-S39.	2.4	2
123	Electrocardiographic Left Atrial Abnormality and B-Type Natriuretic Peptide in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 34-43.	2.0	2
124	Relationship of Four Blood Pressure Indexes to Subclinical Cerebrovascular Diseases Assessed by Brain MRI in General Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 174-187.	2.0	2
125	Lipoprotein Particle Profiles Compared With Standard Lipids in the Association With Subclinical Aortic Valve Calcification in Apparently Healthy Japanese Men. <i>Circulation Journal</i> , 2021, 85, 1076-1082.	1.6	2
126	Relationship of Ambient Temperature Parameters to Stroke Incidence in a Japanese Populationâ€“ Takashima Stroke Registry, Japan, 1988â€“2010 â€“. <i>Circulation Journal</i> , 2021, 85, 2215-2221.	1.6	2

#	ARTICLE	IF	CITATIONS
127	Sex-specific Relationship Between Stress Coping Strategies and All-cause Mortality: Japan Multi-Institutional Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2023, 33, 236-245.	2.4	2
128	The association of reproductive history with hypertension and obesity according to menopausal status: the J-MICC Study. <i>Hypertension Research</i> , 2022, 45, 708-714.	2.7	2
129	Predictors of lower limb fractures in general Japanese: NIPPON DATA90. <i>PLoS ONE</i> , 2022, 17, e0261716.	2.5	2
130	Premature Atrial Contractions and Their Determinants in a General Population of Japanese Men. <i>Circulation Journal</i> , 2022, 86, 1298-1306.	1.6	2
131	Effects of blood pressure and the renin-angiotensin system on platelet activation in type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2010, 1, 196-201.	2.4	1
132	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 746-746.	2.0	1
133	Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. <i>Open Heart</i> , 2019, 6, e001119.	2.3	1
134	Genome-wide association study of serum prostate-specific antigen levels based on 1000 Genomes imputed data in Japanese: the Japan Multi-Institutional Collaborative Cohort Study. <i>Nagoya Journal of Medical Science</i> , 2021, 83, 183-194.	0.3	1
135	Association of perceived stress and coping strategies with the renal function in middle-aged and older Japanese men and women. <i>Scientific Reports</i> , 2022, 12, 291.	3.3	1
136	Effect of the interaction between physical activity and estimated macronutrient intake on HbA1c: population-based cross-sectional and longitudinal studies. <i>BMJ Open Diabetes Research and Care</i> , 2022, 10, e002479.	2.8	1
137	A genome-wide association study on adherence to low-carbohydrate diets in Japanese. <i>European Journal of Clinical Nutrition</i> , 2022, , .	2.9	1
138	Impact of Flushing Response on the Relationship between Alcohol Consumption and Gamma-glutamyl Transpeptidase: the KOBE study. <i>Japanese Journal of Alcohol Studies and Drug Dependence</i> , 2016, 51, 173-183.	0.4	1
139	Association between C-Reactive Protein Levels and Functional Disability in the General Older-Population: The Takashima Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2023, 30, 56-65.	2.0	1
140	Eighteen-year trends in the management of patients with diabetes in the Shiga Diabetes Clinical Survey: overall trends and differences by age group. <i>Diabetology International</i> , 2022, 13, 566-574.	1.4	1
141	ICAD12: OBJECTIVELY MEASURED PHYSICAL ACTIVITY AND BRAIN VOLUME IN JAPANESE ADULT MEN: FINDINGS FROM THE SHIGA EPIDEMIOLOGICAL STUDY OF SUBCLINICAL ATHEROSCLEROSIS. <i>Alzheimer's and Dementia</i> , 2019, 15, P96.	0.8	0
142	The Estimated Absolute Risk of Coronary Artery Disease and Subclinical Atherosclerosis. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 1260-1262.	2.0	0
143	913 Factors of premature atrial contractions among general Japanese men. <i>International Journal of Epidemiology</i> , 2021, 50, .	1.9	0
144	Impact of resting heart rate on cardiovascular mortality according to serum albumin levels in a 24-year follow-up study on a general Japanese population: NIPPON DATA80. <i>Journal of Epidemiology</i> , 2021, , .	2.4	0

#	ARTICLE	IF	CITATIONS
145	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men. <i>Cerebrovascular Diseases</i> , 2022, 51, 774-780.	1.7	0