## Aya Kadota

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1138431/publications.pdf

Version: 2024-02-01

236925 206112 2,986 145 25 48 citations h-index g-index papers 150 150 150 5098 times ranked docs citations citing authors all docs

#	Article	IF	CITATIONS
1	Relationship of Higher-level Functional Capacity With Long-term Mortality in Japanese Older People: NIPPON DATA90. Journal of Epidemiology, 2023, 33, 136-141.	2.4	7
2	Sex-specific Relationship Between Stress Coping Strategies and All-cause Mortality: Japan Multi-Institutional Collaborative Cohort Study. Journal of Epidemiology, 2023, 33, 236-245.	2.4	2
3	Association between C-Reactive Protein Levels and Functional Disability in the General Older-Population: The Takashima Study. Journal of Atherosclerosis and Thrombosis, 2023, 30, 56-65.	2.0	1
4	Relationship of Four Blood Pressure Indexes to Subclinical Cerebrovascular Diseases Assessed by Brain MRI in General Japanese Men. Journal of Atherosclerosis and Thrombosis, 2022, 29, 174-187.	2.0	2
5	Relationship Between Non-fasting Triglycerides and Cardiovascular Disease Mortality in a 20-year Follow-up Study of a Japanese General Population: NIPPON DATA90. Journal of Epidemiology, 2022, 32, 303-313.	2.4	9
6	Association between Stress-Coping Strategy and Functional Disability in the General Older Adult Population: The Takashima Study. Gerontology, 2022, 68, 699-706.	2.8	3
7	A Comparison of Segment-Specific and Composite Measures of Carotid Intima-Media Thickness and their Relationships with Coronary Calcium. Journal of Atherosclerosis and Thrombosis, 2022, 29, 282-295.	2.0	7
8	Relationship between Kidney Function and Subclinical Atherosclerosis Progression Evaluated by Coronary Artery Calcification. Journal of Atherosclerosis and Thrombosis, 2022, 29, 1359-1371.	2.0	9
9	Association of perceived stress and coping strategies with the renal function in middle-aged and older Japanese men and women. Scientific Reports, 2022, 12, 291.	3.3	1
10	Ventricular Premature Complexes and Their Associated Factors in a General Population of Japanese Men. American Journal of Cardiology, 2022, 169, 51-56.	1.6	3
11	The association of reproductive history with hypertension and obesity according to menopausal status: the J-MICC Study. Hypertension Research, 2022, 45, 708-714.	2.7	2
12	Effect of the interaction between physical activity and estimated macronutrient intake on HbA1c: population-based cross-sectional and longitudinal studies. BMJ Open Diabetes Research and Care, 2022, 10, e002479.	2.8	1
13	Predictors of lower limb fractures in general Japanese: NIPPON DATA90. PLoS ONE, 2022, 17, e0261716.	2.5	2
14	Association of equol producing status with aortic calcification in middle-aged Japanese men: The ERA JUMP study. International Journal of Cardiology, 2022, 352, 158-164.	1.7	3
15	A genome-wide association study on adherence to low-carbohydrate diets in Japanese. European Journal of Clinical Nutrition, 2022, , .	2.9	1
16	Premature Atrial Contractions and Their Determinants in a General Population of Japanese Men. Circulation Journal, 2022, 86, 1298-1306.	1.6	2
17	Eighteen-year trends in the management of patients with diabetes in the Shiga Diabetes Clinical Survey: overall trends and differences by age group. Diabetology International, 2022, 13, 566-574.	1.4	1
18	Relationship of leisure-time and household physical activity level and type with cardiovascular disease: secondary analysis of the Takashima Study data. BMC Cardiovascular Disorders, 2022, 22, 132.	1.7	3

#	Article	IF	Citations
19	Trends in Prevalence, Treatment, and Control of Hypertension According to 40-Year-Old Life Expectancy at Prefectures in Japan from the National Health and Nutrition Surveys. Nutrients, 2022, 14, 1219.	4.1	3
20	Red blood cell fatty acid patterns from 7 countries: Focus on the Omega-3 index. Prostaglandins Leukotrienes and Essential Fatty Acids, 2022, 179, 102418.	2.2	21
21	Eating Slowly Is Associated with Undernutrition among Community-Dwelling Adult Men and Older Adult Women. Nutrients, 2022, 14, 54.	4.1	4
22	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men. Cerebrovascular Diseases, 2022, 51, 774-780.	1.7	0
23	Electrocardiographic Left Atrial Abnormality and B-Type Natriuretic Peptide in a General Japanese Population: NIPPON DATA2010. Journal of Atherosclerosis and Thrombosis, 2021, 28, 34-43.	2.0	2
24	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. Hypertension Research, 2021, 44, 80-87.	2.7	6
25	Impact of <i>PSCA</i> Polymorphisms on the Risk of Duodenal Ulcer. Journal of Epidemiology, 2021, 31, 12-20.	2.4	9
26	The Estimated Absolute Risk of Coronary Artery Disease and Subclinical Atherosclerosis. Journal of Atherosclerosis and Thrombosis, 2021, 28, 1260-1262.	2.0	0
27	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. Alcohol, 2021, 90, 57-65.	1.7	6
28	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. British Journal of Nutrition, 2021, 126, 1843-1851.	2.3	6
29	Relationship Between Calcium Intake and Impaired Activities of Daily Living in a Japanese Population: NIPPON DATA90. Journal of Epidemiology, 2021, 31, 119-124.	2.4	3
30	Risk Factors That Most Accurately Predict Coronary Artery Disease Based on the Duration of Follow-up ― NIPPON DATA80 ―. Circulation Journal, 2021, 85, 908-913.	1.6	5
31	Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. Environmental Health and Preventive Medicine, 2021, 26, 57.	3.4	3
32	Lipoprotein Particle Profiles Compared With Standard Lipids in the Association With Subclinical Aortic Valve Calcification in Apparently Healthy Japanese Men. Circulation Journal, 2021, 85, 1076-1082.	1.6	2
33	Effect of Underlying Cardiometabolic Diseases on the Association Between Sedentary Time and Allâ€Cause Mortality in a Large Japanese Population: A Cohort Analysis Based on the Jâ€MICC Study. Journal of the American Heart Association, 2021, 10, e018293.	3.7	9
34	Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. PLoS ONE, 2021, 16, e0254706.	2.5	5
35	Relationship of Ambient Temperature Parameters to Stroke Incidence in a Japanese Population ― Takashima Stroke Registry, Japan, 1988–2010 ―. Circulation Journal, 2021, 85, 2215-2221.	1.6	2
36	913Factors of premature atrial contractions among general Japanese men. International Journal of Epidemiology, 2021, 50, .	1.9	0

#	Article	IF	CITATIONS
37	Impact of resting heart rate on cardiovascular mortality according to serum albumin levels in a 24-year follow-up study on a general Japanese population: NIPPON DATA80. Journal of Epidemiology, 2021, , .	2.4	0
38	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. Journal of Nutritional Science, 2021, 10, e61.	1.9	3
39	Genome-wide association study of serum prostate-specific antigen levels based on 1000 Genomes imputed data in Japanese: the Japan Multi-Institutional Collaborative Cohort Study. Nagoya Journal of Medical Science, 2021, 83, 183-194.	0.3	1
40	Dietary Inflammatory Index Positively Associated With High-Sensitivity C-Reactive Protein Level in Japanese From NIPPON DATA2010. Journal of Epidemiology, 2020, 30, 98-107.	2.4	18
41	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. Journal of Epidemiology, 2020, 30, 244-252.	2.4	10
42	Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. Journal of Epidemiology, 2020, 30, 121-127.	2.4	9
43	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. European Journal of Clinical Nutrition, 2020, 74, 67-76.	2.9	17
44	The interaction between ABCA1 polymorphism and physical activity on the HDL-cholesterol levels in a Japanese population. Journal of Lipid Research, 2020, 61, 86-94.	4.2	11
45	Alcohol consumption and cognitive function in elderly Japanese men. Alcohol, 2020, 85, 145-152.	1.7	7
46	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical) Tj ETQq0 0 0 rgBT /Ove	rlo <b>e</b> 1810 T	f 5 <b>0</b> 377 Td (
47	Relationship of household salt intake level with long-term all-cause and cardiovascular disease mortality in Japan: NIPPON DATA80. Hypertension Research, 2020, 43, 132-139.	2.7	11
48	The relationship between serum levels of LOX-1 ligand containing ApoAl as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. Atherosclerosis, 2020, 313, 20-25.	0.8	12
49	Cross-sectional association of bone mineral density with coronary artery calcification in an international multi-ethnic population-based cohort of men aged 40–49: ERA JUMP study. IJC Heart and Vasculature, 2020, 30, 100618.	1.1	3
50	Association between plasma levels of homocysteine, folate, and vitamin B12, and dietary folate intake and hypertension in a cross-sectional study. Scientific Reports, 2020, 10, 18499.	3.3	14
51	Estimated 24 h Urinary Sodium-to-Potassium Ratio Is Related to Renal Function Decline: A 6-Year Cohort Study of Japanese Urban Residents. International Journal of Environmental Research and Public Health, 2020, 17, 5811.	2.6	9
52	Association of Red Meat Intake with the Risk of Cardiovascular Mortality in General Japanese Stratified by Kidney Function: NIPPON DATA80. Nutrients, 2020, 12, 3707.	4.1	4
53	Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. Stroke, 2020, 51, 3584-3591.	2.0	19
54	Smoking habits and progression of coronary and aortic artery calcification: A 5-year follow-up of community-dwelling Japanese men. International Journal of Cardiology, 2020, 314, 89-94.	1.7	7

#	Article	IF	CITATIONS
55	Association of Dietary Acid Load with the Prevalence of Metabolic Syndrome among Participants in Baseline Survey of the Japan Multi-Institutional Collaborative Cohort Study. Nutrients, 2020, 12, 1605.	4.1	23
56	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 995-1009.	2.0	12
57	Epidemiology of hypertension in Japan: beyond the new 2019 Japanese guidelines. Hypertension Research, 2020, 43, 1344-1351.	2.7	49
58	Genome-wide association meta-analysis identifies GP2 gene risk variants for pancreatic cancer. Nature Communications, 2020, 11, 3175.	12.8	34
59	Identification of a novel uterine leiomyoma GWAS locus in a Japanese population. Scientific Reports, 2020, 10, 1197.	3.3	14
60	Sedentary Time is Associated with Cardiometabolic Diseases in A Large Japanese Population: A Cross-Sectional Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 1097-1107.	2.0	14
61	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. Tohoku Journal of Experimental Medicine, 2020, 252, 253-262.	1.2	5
62	Physique at Birth and Cardiovascular Disease Risk Factors in Japanese Urban Residents: the KOBE Study. Journal of Atherosclerosis and Thrombosis, 2020, , .	2.0	3
63	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. Journal of Epidemiology, 2019, 29, 133-138.	2.4	9
64	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. European Journal of Clinical Nutrition, 2019, 73, 783-792.	2.9	22
65	Serum long-chain n-3 polyunsaturated fatty acids and aortic calcification in middle-aged men: The population-based cross-sectional ERA-JUMP study. Nutrition, Metabolism and Cardiovascular Diseases, 2019, 29, 837-846.	2.6	3
66	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Journal of Atherosclerosis and Thrombosis, 2019, 26, 746-746.	2.0	1
67	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan ― NIPPON DATA80 ―. Circulation Journal, 2019, 83, 1254-1260.	1.6	11
68	Socioeconomic and lifestyle factors associated with depressive tendencies in general Japanese men and women: NIPPON DATA2010. Environmental Health and Preventive Medicine, 2019, 24, 37.	3.4	11
69	Associations of Nutrient Patterns with the Prevalence of Metabolic Syndrome: Results from the Baseline Data of the Japan Multi-Institutional Collaborative Cohort Study. Nutrients, 2019, 11, 990.	4.1	24
70	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men ― A 20-Year Follow-up of NIPPON DATA90 ―. Circulation Journal, 2019, 83, 1506-1513.	1.6	7
71	Anthropometric Obesity Indices were Stronger than CT-Based Indices in Associations with Carotid Intima-Media Thickness in Japanese Men. Journal of Atherosclerosis and Thrombosis, 2019, 26, 1102-1114.	2.0	3
72	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. Journal of Diabetes and Its Complications, 2019, 33, 417-423.	2.3	8

#	Article	IF	CITATIONS
73	Differences Between Coronary Artery Calcification and Aortic Artery Calcification in Relation to Cardiovascular Disease Risk Factors in Japanese Men. Journal of Atherosclerosis and Thrombosis, 2019, 26, 452-464.	2.0	13
74	Oneâ€year weight loss maintenance outcomes following a worksiteâ€based weight reduction program among Japanese men with cardiovascular risk factors. Journal of Occupational Health, 2019, 61, 189-196.	2.1	7
75	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. Hypertension Research, 2019, 42, 411-418.	2.7	21
76	Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. Open Heart, 2019, 6, e001119.	2.3	1
77	ICâ€Pâ€112: OBJECTIVELY MEASURED PHYSICAL ACTIVITY AND BRAIN VOLUME IN JAPANESE ADULT MEN: FINDIN FROM THE SHIGA EPIDEMIOLOGICAL STUDY OF SUBCLINICAL ATHEROSCLEROSIS. Alzheimer's and Dementia, 2019, 15, P96.	IGS 0.8	0
78	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. Journal of Hypertension, 2019, 37, 1676-1681.	0.5	7
79	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. Environmental Health and Preventive Medicine, 2019, 24, 1.	3.4	84
80	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality inÂa 15-Year Follow-Up Study ofÂthe General Japanese Population. Journal of Atherosclerosis and Thrombosis, 2019, 26, 198-206.	2.0	17
81	Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. Nature Genetics, 2019, 51, 379-386.	21.4	164
82	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Journal of Epidemiology, 2019, 29, 205-212.	2.4	9
83	Re-evaluation of the associations of egg intake with serum total cholesterol and cause-specific and total mortality in Japanese women. European Journal of Clinical Nutrition, 2018, 72, 841-847.	2.9	21
84	Association of Coronary Artery Calcification with Estimated Coronary Heart Disease Risk from Prediction Models in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Journal of Atherosclerosis and Thrombosis, 2018, 25, 477-489.	2.0	4
85	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. Metabolic Syndrome and Related Disorders, 2018, 16, 166-173.	1.3	9
86	Overweight or underweight and the risk of decline in activities of daily living in a 22â€year cohort study of a Japanese sample. Geriatrics and Gerontology International, 2018, 18, 799-805.	1.5	8
87	Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. Atherosclerosis, 2018, 273, 145-152.	0.8	27
88	Dietary tofu intake and long-term risk of death from stroke in a general population. Clinical Nutrition, 2018, 37, 182-188.	5.0	10
89	Relationships among Socioeconomic Factors and Self-rated Health in Japanese Adults: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S66-S72.	2.4	12
90	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S59-S65.	2.4	11

#	Article	IF	CITATIONS
91	Self-reported Sleep Duration and Subclinical Atherosclerosis in a General Population of Japanese Men. Journal of Atherosclerosis and Thrombosis, 2018, 25, 186-198.	2.0	5
92	The National Integrated Project for Prospective Observation of Non-communicable Disease and its Trends in the Aged 2010 (NIPPON DATA2010): Objectives, Design, and Population Characteristics. Journal of Epidemiology, 2018, 28, S2-S9.	2.4	29
93	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S23-S28.	2.4	10
94	Relationship Between Socioeconomic Status and the Prevalence of Underweight, Overweight or Obesity in a General Japanese Population: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S10-S16.	2.4	26
95	Macronutrient Intake and Socioeconomic Status: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S17-S22.	2.4	19
96	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S29-S34.	2.4	15
97	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S35-S39.	2.4	2
98	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. Journal of Atherosclerosis and Thrombosis, 2018, 25, 606-620.	2.0	17
99	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S40-S45.	2.4	4
100	Time-Related Changes in Relationships Between the Keys Score, Dietary Lipids, and Serum Total Cholesterol in Japan ― NIPPON DATA80/90/2010 ―. Circulation Journal, 2018, 83, 147-155.	1.6	4
101	Socioeconomic Status and Knowledge of Cardiovascular Risk Factors: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S46-S52.	2.4	17
102	Factors Related to Participation in Health Examinations for Japanese National Health Insurance: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S53-S58.	2.4	8
103	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. Circulation Journal, 2018, 82, 2542-2548.	1.6	11
104	Comparison of carotid plaque burden among healthy middle-aged men living in the US, Japan, and South Korea. International Journal of Cardiology, 2018, 266, 245-249.	1.7	9
105	Twelve-year trends of increasing overweight and obesity in patients with diabetes: the Shiga Diabetes Clinical Survey. Endocrine Journal, 2018, 65, 527-536.	1.6	21
106	Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. International Journal of Cardiology, 2018, 267, 177-182.	1.7	30
107	Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. Medicine and Science in Sports and Exercise, 2018, 50, 2433-2441.	0.4	36
108	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. British Journal of Nutrition, 2017, 117, 260-266.	2.3	31

#	Article	IF	CITATIONS
109	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population. Journal of Hypertension, 2017, 35, 401-408.	0.5	74
110	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. European Journal of Preventive Cardiology, 2017, 24, 505-513.	1.8	8
111	Trends in medical performance in diabetic patients in primary care clinics compared with those in hospitals: Shiga Diabetes Clinical Survey, Japan, 2000–2012. Diabetology International, 2017, 8, 59-68.	1.4	6
112	Increased Aortic Calcification Is Associated With Arterial Stiffness Progression in Multiethnic Middle-Aged Men. Hypertension, 2017, 69, 102-108.	2.7	51
113	Genetic Variants of <i>RAMP2</i> and <i>CLR</i> are Associated with Stroke. Journal of Atherosclerosis and Thrombosis, 2017, 24, 1267-1281.	2.0	11
114	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. Journal of Atherosclerosis and Thrombosis, 2016, 23, 800-809.	2.0	48
115	Underweight Young Women Without Later Weight Gain Are at High Risk for Osteopenia After Midlife: The KOBE Study. Journal of Epidemiology, 2016, 26, 572-578.	2.4	20
116	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. Environmental Pollution, 2016, 213, 460-467.	7.5	16
117	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. Hypertension Research, 2016, 39, 670-679.	2.7	8
118	Influence of cigarette smoking and inflammatory gene polymorphisms on glycated hemoglobin in the Japanese general population. Preventive Medicine Reports, 2016, 3, 288-295.	1.8	2
119	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. Journal of the American Heart Association, 2016, 5, .	3.7	39
120	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. Atherosclerosis, 2016, 252, 116-121.	0.8	9
121	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Preventive Medicine Reports, 2016, 4, 225-232.	1.8	30
122	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. International Journal of Cardiology, 2016, 220, 262-267.	1.7	29
123	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components. Arteriosclerosis, Thrombosis, and Vascular Biology, 2016, 36, 1703-1708.	2.4	44
124	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. Atherosclerosis, 2016, 246, 141-147.	0.8	48
125	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemiaLOX-1 ligand and IMT in Japanese. Journal of Clinical Lipidology, 2016, 10, 172-180.e1.	1.5	11
126	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. Journal of Clinical Lipidology, 2016, 10, 1195-1202.e1.	1.5	12

#	Article	IF	CITATIONS
127	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. PLoS ONE, 2016, 11, e0157563.	2.5	4
128	Impact of Flushing Response on the Relationship between Alcohol Consumption and Gamma-glutamyl Transpeptidase: the KOBE study. Japanese Journal of Alcohol Studies and Drug Dependence, 2016, 51, 173-183.	0.4	1
129	Effects of Stroke Education of Junior High School Students on Stroke Knowledge of Their Parents. Stroke, 2015, 46, 572-574.	2.0	37
130	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. Atherosclerosis, 2015, 239, 444-450.	0.8	18
131	Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. International Journal of Cardiology, 2015, 189, 67-72.	1.7	24
132	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980–2010. Hypertension Research, 2015, 38, 790-795.	2.7	39
133	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. British Journal of Nutrition, 2014, 112, 916-924.	2.3	59
134	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: A 24-year follow-up of Japanese general population. Journal of Cardiology, 2014, 64, 218-224.	1.9	11
135	Impact of Metabolic Syndrome on the Risk of Cardiovascular Disease Mortality in the United States and in Japan. American Journal of Cardiology, 2014, 113, 84-89.	1.6	69
136	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. Atherosclerosis, 2014, 236, 237-243.	0.8	22
137	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. Heart, 2013, 99, 1024-1029.	2.9	7
138	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Journal of Atherosclerosis and Thrombosis, 2013, 20, 755-766.	2.0	43
139	Does High-Sensitivity C-Reactive Protein or Low-Density Lipoprotein Cholesterol Show a Stronger Relationship with the Cardio-Ankle Vascular Index in Healthy Community Dwellers?: the KOBE Study. Journal of Atherosclerosis and Thrombosis, 2012, 19, 1027-1034.	2.0	35
140	Relationship of moderate metabolic risk factor clustering to cardiovascular disease mortality in non-lean Japanese: A 15-year follow-up of NIPPON DATA90. Atherosclerosis, 2011, 215, 209-213.	0.8	11
141	Brachial–ankle pulse wave velocity predicts all-cause mortality in the general population: findings from the Takashima study, Japan. Hypertension Research, 2010, 33, 922-925.	2.7	125
142	Effects of blood pressure and the reninâ $\in$ angiotensin system on platelet activation in typeâ $\in$ f2 diabetes. Journal of Diabetes Investigation, 2010, 1, 196-201.	2.4	1
143	Relationships between Family Histories of Stroke and of Hypertension and Stroke Mortality: NIPPON DATA80, 1980-1999. Hypertension Research, 2008, 31, 1525-1531.	2.7	11
144	Cardiovascular Disease and Risk Factors in Asia. Circulation, 2008, 118, 2702-2709.	1.6	604

#	Article	IF	CITATIONS
145	Relationship Between Metabolic Risk Factor Clustering and Cardiovascular Mortality Stratified by High Blood Glucose and Obesity. Diabetes Care, 2007, 30, 1533-1538.	8.6	120