Kenji Wakai

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1143539/publications.pdf

Version: 2024-02-01

411 papers

16,975 citations

65 h-index 27406 106 g-index

414 all docs

414 docs citations

times ranked

414

19213 citing authors

#	Article	IF	CITATIONS
1	Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. Clinical and Experimental Nephrology, 2007, 11, 41-50.	1.6	488
2	Prevalence and Clinicoepidemiological Features of Moyamoya Disease in Japan. Stroke, 2008, 39, 42-47.	2.0	404
3	The Relationship between Green Tea and Total Caffeine Intake and Risk for Self-Reported Type 2 Diabetes among Japanese Adults. Annals of Internal Medicine, 2006, 144, 554.	3.9	389
4	Prospective Cohort Study of the Risk of Prostate Cancer among Rotating-Shift Workers: Findings from the Japan Collaborative Cohort Study. American Journal of Epidemiology, 2006, 164, 549-555.	3.4	348
5	Dietary Risk Factors for Inflammatory Bowel Disease. Inflammatory Bowel Diseases, 2005, 11, 154-163.	1.9	338
6	Common Defects of ABCG2, a High-Capacity Urate Exporter, Cause Gout: A Function-Based Genetic Analysis in a Japanese Population. Science Translational Medicine, 2009, 1, 5ra11.	12.4	334
7	Epidemiological features of Moyamoya disease in Japan: findings from a nationwide survey. Clinical Neurology and Neurosurgery, 1997, 99, S1-S5.	1.4	330
8	An Overview of Regular Dialysis Treatment in Japan (As of 31 December 2013). Therapeutic Apheresis and Dialysis, 2015, 19, 540-574.	0.9	275
9	Prevalence of chronic kidney disease (CKD) in the Japanese general population predicted by the MDRD equation modified by a Japanese coefficient. Clinical and Experimental Nephrology, 2007, 11, 156-163.	1.6	267
10	Fish, ω-3 Polyunsaturated Fatty Acids, and Mortality From Cardiovascular Diseases in a Nationwide Community-Based Cohort of Japanese Men and Women. Journal of the American College of Cardiology, 2008, 52, 988-996.	2.8	251
11	Dietary Intake and Sources of Isoflavones Among Japanese. Nutrition and Cancer, 1999, 33, 139-145.	2.0	205
12	A scoring system to predict renal outcome in IgA nephropathy: a nationwide 10-year prospective cohort study. Nephrology Dialysis Transplantation, 2009, 24, 3068-3074.	0.7	203
13	Smoking Increases the Treatment Failure for Helicobacter pylori Eradication. American Journal of Medicine, 2006, 119, 217-224.	1.5	164
14	Overview of Regular Dialysis Treatment in Japan (as of 31 December 2008). Therapeutic Apheresis and Dialysis, 2010, 14, 505-540.	0.9	151
15	A scoring system to predict renal outcome in IgA nephropathy: from a nationwide prospective study. Nephrology Dialysis Transplantation, 2006, 21, 2800-2808.	0.7	147
16	Serum phytoestrogens and prostate cancer risk in a nested caseâ€control study among Japanese men. Cancer Science, 2004, 95, 65-71.	3.9	143
17	Associations of medical status and physical fitness with periodontal disease. Journal of Clinical Periodontology, 1999, 26, 664-672.	4.9	136
18	Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. Nature Genetics, 2017, 49, 953-958.	21.4	136

#	Article	IF	Citations
19	Reproducibility and Validity of a Self-administered Food Frequency Questionnaire Used in the JACC Study. Journal of Epidemiology, 2005, 15, S9-S23.	2.4	135
20	Cohort Profile of the Japan Collaborative Cohort Study at Final Follow-up. Journal of Epidemiology, 2013, 23, 227-232.	2.4	134
21	Estimated Prevalence and Incidence of Adult Still's Disease: Findings by a Nationwide Epidemiological Survey in Japan. Journal of Epidemiology, 1997, 7, 221-225.	2.4	132
22	Overview of Regular Dialysis Treatment in <scp>J</scp> apan (as of 31 <scp>D</scp> ecember 2011). Therapeutic Apheresis and Dialysis, 2013, 17, 567-611.	0.9	132
23	A Review of Food Frequency Questionnaires Developed and Validated in Japan. Journal of Epidemiology, 2009, 19, 1-11.	2.4	125
24	ABCG2 dysfunction causes hyperuricemia due to both renal urate underexcretion and renal urate overload. Scientific Reports, 2014, 4, 3755.	3.3	125
25	Glomerular hyperfiltration in prediabetes and prehypertension. Nephrology Dialysis Transplantation, 2012, 27, 1821-1825.	0.7	124
26	Dietary Fiber Intake Is Associated with Reduced Risk of Mortality from Cardiovascular Disease among Japanese Men and Women , ,. Journal of Nutrition, 2010, 140, 1445-1453.	2.9	119
27	Alcohol Dehydrogenase 2 His47Arg Polymorphism Influences Drinking Habit Independently of Aldehyde Dehydrogenase 2 Glu487Lys Polymorphism: Analysis of 2,299 Japanese Subjects. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 1009-1013.	2.5	116
28	An Overview of Regular Dialysis Treatment in <scp>J</scp> apan (as of 31 <scp>D</scp> ecember 2012). Therapeutic Apheresis and Dialysis, 2014, 18, 535-602.	0.9	115
29	GWAS of clinically defined gout and subtypes identifies multiple susceptibility loci that include urate transporter genes. Annals of the Rheumatic Diseases, 2017, 76, 869-877.	0.9	114
30	Tooth loss and intakes of nutrients and foods: a nationwide survey of Japanese dentists. Community Dentistry and Oral Epidemiology, 2010, 38, 43-49.	1.9	111
31	An Overview of Regular Dialysis Treatment in Japan (As of 31 December 2010). Therapeutic Apheresis and Dialysis, 2012, 16, 483-521.	0.9	111
32	Nationwide epidemiological survey of chronic pancreatitis in Japan. Journal of Gastroenterology, 2000, 35, 136-141.	5.1	110
33	Alcohol Drinking and Colorectal Cancer in Japanese: A Pooled Analysis of Results from Five Cohort Studies. American Journal of Epidemiology, 2008, 167, 1397-1406.	3.4	107
34	Common dysfunctional variants in ABCG2 are a major cause of early-onset gout. Scientific Reports, 2013, 3, 2014.	3.3	105
35	Body Mass Index and Mortality From All Causes and Major Causes in Japanese: Results of a Pooled Analysis of 7 Large-Scale Cohort Studies. Journal of Epidemiology, 2011, 21, 417-430.	2.4	100
36	Dietary intakes of fat and fatty acids and risk of breast cancer: A prospective study in Japan. Cancer Science, 2005, 96, 590-599.	3.9	97

#	Article	IF	CITATIONS
37	Tobacco Smoking and Gastric Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 800-807.	1.3	95
38	The Global DNA Methylation Surrogate LINE-1 Methylation Is Correlated with MGMT Promoter Methylation and Is a Better Prognostic Factor for Glioma. PLoS ONE, 2011, 6, e23332.	2.5	95
39	Leptin Is Associated with an Increased Female Colorectal Cancer Risk: A Nested Case-Control Study in Japan. Oncology, 2005, 68, 454-461.	1.9	94
40	Tobacco Smoking and Lung Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiological Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 309-324.	1.3	94
41	Diabetes mellitus and cancer risk: Pooled analysis of eight cohort studies in Japan. Cancer Science, 2013, 104, 1499-1507.	3.9	94
42	Profile of Participants and Genotype Distributions of 108 Polymorphisms in a Cross-Sectional Study of Associations of Genotypes With Lifestyle and Clinical Factors: A Project in the Japan Multi-Institutional Collaborative Cohort (J-MICC) Study. Journal of Epidemiology, 2011, 21, 223-235.	2.4	92
43	An Overview of Regular Dialysis Treatment in Japan (As of 31 December 2007). Therapeutic Apheresis and Dialysis, 2009, 13, 457-504.	0.9	90
44	Alcohol Consumption and Mortality among Middle-aged and Elderly Japanese Men and Women. Annals of Epidemiology, 2005, 15, 590-597.	1.9	87
45	Risk factors differ for non-small-cell lung cancers with and without EGFR mutation: assessment of smoking and sex by a case-control study in Japanese. Cancer Science, 2007, 98, 96-101.	3.9	86
46	Benefits of interferon $\hat{\mathbf{e}}^2$ and temozolomide combination therapy for newly diagnosed primary glioblastoma with the unmethylated MGMT promoter. Cancer, 2011, 117, 1721-1730.	4.1	85
47	Tobacco Smoking and Colorectal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 25-39.	1.3	83
48	Overview of Regular Dialysis Treatment in Japan (as of 31 December 2009). Therapeutic Apheresis and Dialysis, 2012, 16, 11-53.	0.9	83
49	Dietary Risk Factors for Colon and Rectal Cancers: A Comparative Case-Control Study. Journal of Epidemiology, 2006, 16, 125-135.	2.4	81
50	One-carbon metabolism related gene polymorphisms interact with alcohol drinking to influence the risk of colorectal cancer in Japan. Carcinogenesis, 2005, 26, 2164-2171.	2.8	80
51	A Comparison of Ovarian Metastasis between Squamous Cell Carcinoma and Adenocarcinoma of the Uterine Cervix. Gynecologic Oncology, 2001, 82, 504-509.	1.4	79
52	Diet and Colorectal Cancer Mortality: Results From the Japan Collaborative Cohort Study. Nutrition and Cancer, 2004, 50, 23-32.	2.0	79
53	Soy Intake and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2014, 44, 282-295.	1.3	79
54	Trends in incidence of end-stage renal disease in Japan, 1983-2000: age-adjusted and age-specific rates by gender and cause. Nephrology Dialysis Transplantation, 2004, 19, 2044-2052.	0.7	78

#	Article	IF	CITATIONS
55	Acute exacerbation of IPF following diagnostic bronchoalveolar lavage procedures. Respiratory Medicine, 2012, 106, 436-442.	2.9	77
56	Breast cancer risk and erythrocyte compositions of nâ€3 highly unsaturated fatty acids in Japanese. International Journal of Cancer, 2007, 121, 377-385.	5.1	76
57	<i>Tollâ€Like Receptor 4 +</i> 3725 G/C Polymorphism, <i>Helicobacter pylori</i> Seropositivity, and the Risk of Gastric Atrophy and Gastric Cancer in Japanese. Helicobacter, 2009, 14, 47-53.	3.5	76
58	Dietary Fiber and Risk of Colorectal Cancer in the Japan Collaborative Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2007, 16, 668-675.	2.5	74
59	Genome-wide association study revealed novel loci which aggravate asymptomatic hyperuricaemia into gout. Annals of the Rheumatic Diseases, 2019, 78, 1430-1437.	0.9	73
60	Foods and Nutrients in Relation to Bladder Cancer Risk: A Case-Control Study in Aichi Prefecture, Central Japan. Nutrition and Cancer, 2000, 38, 13-22.	2.0	72
61	Serum Levels of Polyunsaturated Fatty Acids and Risk of Colorectal Cancer: A Prospective Study. American Journal of Epidemiology, 2005, 161, 462-471.	3.4	71
62	Gene?gene and gene?environment interactions between alcohol drinking habit and polymorphisms in alcohol-metabolizing enzyme genes and the risk of head and neck cancer in Japan. Cancer Science, 2007, 98, 1087-1091.	3.9	70
63	Common dysfunctional variants of ABCG2 have stronger impact on hyperuricemia progression than typical environmental risk factors. Scientific Reports, 2014, 4, 5227.	3.3	70
64	Secondhand smoke exposure and risk of lung cancer in Japan: a systematic review and meta-analysis of epidemiologic studies. Japanese Journal of Clinical Oncology, 2016, 46, 942-951.	1.3	70
65	Alcohol Drinking and Esophageal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2011, 41, 677-692.	1.3	69
66	Relationship between obesity and serum markers of oxidative stress and inflammation in Japanese. Asian Pacific Journal of Cancer Prevention, 2003, 4, 259-66.	1.2	69
67	Tea Consumption and Lung Cancer Risk: A Case-Control Study in Okinawa, Japan. Japanese Journal of Cancer Research, 1995, 86, 1027-1034.	1.7	67
68	Risk of Colorectal Cancer Is Linked to Erythrocyte Compositions of Fatty Acids as Biomarkers for Dietary Intakes of Fish, Fat, and Fatty Acids. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 1791-1798.	2.5	67
69	Attributable and absolute risk of lung cancer death by smoking status: Findings from the Japan collaborative cohort study. International Journal of Cancer, 2003, 105, 249-254.	5.1	66
70	Genome-wide meta-analysis identifies multiple novel loci associated with serum uric acid levels in Japanese individuals. Communications Biology, 2019, 2, 115.	4.4	66
71	Genetic Predisposition to Ischemic Stroke. Stroke, 2017, 48, 253-258.	2.0	64
72	The current status of the treatment for hemorrhagic type Moyamoya disease based on a 1995 nationwide survey in Japan. Clinical Neurology and Neurosurgery, 1997, 99, S183-S186.	1.4	63

#	Article	IF	CITATIONS
73	A Simple Food Frequency Questionnaire for Japanese Diet-Part I. Development of the Questionnaire, and Reproducibility and Validity for Food Groups. Journal of Epidemiology, 1999, 9, 216-226.	2.4	63
74	Overview of Regular Dialysis Treatment in Japan as of 31â€∫December 2006. Therapeutic Apheresis and Dialysis, 2008, 12, 428-456.	0.9	63
75	A Simple Food Frequency Questionnaire for Japanese Diet-Part II. Reproducibility and Validity for Nutrient Intakes Journal of Epidemiology, 1999, 9, 227-234.	2.4	62
76	Validity and Reliability of Single-item Questions about Physical Activity Journal of Epidemiology, 2001, 11, 211-218.	2.4	62
77	Foods and beverages in relation to urothelial cancer: Caseâ€control study in Japan. International Journal of Urology, 2004, 11, 11-19.	1.0	62
78	Gene–environment interactions between alcohol drinking and the MTHFR C677T polymorphism impact on esophageal cancer risk: results of a case–control study in Japan. Carcinogenesis, 2005, 26, 1285-1290.	2.8	62
79	Tobacco Smoking and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiological Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 387-394.	1.3	62
80	Combined Effect of miR-146a rs2910164 G/C Polymorphism and Toll-like Receptor 4 +3725 G/C Polymorphism on the Risk of Severe Gastric Atrophy in Japanese. Digestive Diseases and Sciences, 2011, 56, 1131-1137.	2.3	62
81	Risk modification byCYP1A1 andGSTM1 polymorphisms in the association of environmental tobacco smoke and lung cancer: A case-control study in Japanese nonsmoking women. International Journal of Cancer, 2003, 107, 139-144.	5.1	61
82	Meat Consumption and Colorectal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2014, 44, 641-650.	1.3	60
83	Dietary Habits and Risk of Lung Cancer Death in a Large-scale Cohort Study (JACC Study) in Japan by Sex and Smoking Habit. Japanese Journal of Cancer Research, 2001, 92, 1259-1269.	1.7	59
84	Consumption of soy foods and the risk of breast cancer: findings from the Japan Collaborative Cohort (JACC) Study. Cancer Causes and Control, 2007, 18, 801-808.	1.8	59
85	Impact of alcohol intake on total mortality and mortality from major causes in Japan: a pooled analysis of six large-scale cohort studies. Journal of Epidemiology and Community Health, 2012, 66, 448-456.	3.7	59
86	An intervention study of smoking cessation with feedback on genetic cancer susceptibility in Japan. Preventive Medicine, 2006, 42, 102-108.	3.4	58
87	Alcohol Drinking and Liver Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2008, 38, 816-838.	1.3	58
88	Esophageal cancer risk by ALDH2 and ADH2 polymorphisms and alcohol consumption: exploration of gene-environment and gene-gene interactions. Asian Pacific Journal of Cancer Prevention, 2005, 6, 256-62.	1.2	58
89	Genetic polymorphism in the phenobarbital-responsive enhancer module of the UDP-glucuronosyltransferase 1A1 gene and irinotecan toxicity. Pharmacogenetics and Genomics, 2005, 15, 35-41.	1.5	57
90	Overview of Regular Dialysis Treatment in Japan (as of 31â€fDecember 2005). Therapeutic Apheresis and Dialysis, 2007, 11, 411-441.	0.9	56

#	Article	IF	Citations
91	Coffee consumption and the risk of endometrial cancer: Evidence from a case-control study of female hormone-related cancers in Japan. Cancer Science, 2007, 98, 411-415.	3.9	54
92	One-carbon metabolism-related gene polymorphisms and risk of head and neck squamous cell carcinoma: Case-control study. Cancer Science, 2007, 98, 1439-1446.	3.9	54
93	Comparison of lifestyle risk factors by family history for gastric, breast, lung and colorectal cancer. Asian Pacific Journal of Cancer Prevention, 2004, 5, 419-27.	1.2	54
94	An Overview of Regular Dialysis Treatment in Japan (as of 31 December 2004). Therapeutic Apheresis and Dialysis, 2006, 10, 476-497.	0.9	53
95	Cigarette Smoking and Esophageal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 63-73.	1.3	53
96	A gene–gene interaction between ALDH2 Glu487Lys and ADH2 His47Arg polymorphisms regarding the risk of colorectal cancer in Japan. Carcinogenesis, 2006, 27, 1018-1023.	2.8	52
97	Long-term effects of a diet loosely restricting carbohydrates on HbA1c levels, BMI and tapering of sulfonylureas in type 2 diabetes: A 2-year follow-up study. Diabetes Research and Clinical Practice, 2008, 79, 350-356.	2.8	52
98	GWAS identifies two novel colorectal cancer loci at 16q24.1 and 20q13.12. Carcinogenesis, 2018, 39, 652-660.	2.8	52
99	Serum carotenoids and mortality from lung cancer: a case-control study nested in the Japan Collaborative Cohort (JACC) Study. Cancer Science, 2003, 94, 57-63.	3.9	51
100	Colorectal Cancer and Serum C-reactive Protein Levels: a Case-control Study Nested in the JACC Study. Journal of Epidemiology, 2005, 15, S185-S189.	2.4	50
101	Possible antenatal and perinatal related factors in development of cystic periventricular leukomalacia. Brain and Development, 2005, 27, 17-21.	1.1	50
102	Alcohol Drinking and Colorectal Cancer Risk: an Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2006, 36, 582-597.	1.3	50
103	Effect of dietary antioxidants and risk of oral, pharyngeal and laryngeal squamous cell carcinoma according to smoking and drinking habits. Cancer Science, 2006, 97, 760-767.	3.9	49
104	Active Smoking, Passive Smoking, and Breast Cancer Risk: Findings from the Japan Collaborative Cohort Study for Evaluation of Cancer Risk. Journal of Epidemiology, 2008, 18, 77-83.	2.4	49
105	12 new susceptibility loci for prostate cancer identified by genome-wide association study in Japanese population. Nature Communications, 2019, 10, 4422.	12.8	49
106	Risk stratification for progression of IgA nephropathy using a decision tree induction algorithm. Nephrology Dialysis Transplantation, 2008, 24, 1242-1247.	0.7	48
107	A common missense variant of monocarboxylate transporter 9 (MCT9/SLC16A9) gene is associated with renal overload gout, but not with all gout susceptibility. Human Cell, 2013, 26, 133-136.	2.7	48
108	Epidemiology of Breast Cancer in Japan. International Journal of Epidemiology, 1995, 24, 285-291.	1.9	47

#	Article	IF	Citations
109	Association of vegetable and fruit intake with gastric cancer risk among Japanese: a pooled analysis of four cohort studies. Annals of Oncology, 2014, 25, 1228-1233.	1.2	47
110	Fat intake and breast cancer risk in an area where fat intake is low: a case-control study in Indonesia. International Journal of Epidemiology, 2000, 29, 20-28.	1.9	46
111	Serum Heat Shock Protein 70 Levels and Lung Cancer Risk: A Case-Control Study Nested in a Large Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 1733-1737.	2.5	46
112	Cigarette Smoking and Liver Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among Japanese. Japanese Journal of Clinical Oncology, 2006, 36, 445-456.	1.3	46
113	Diabetes mellitus and kidney cancer risk: The results of Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC Study). International Journal of Urology, 2007, 14, 393-397.	1.0	46
114	Cigarette Smoking and Pancreas Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence in the Japanese Population. Japanese Journal of Clinical Oncology, 2011, 41, 1292-1302.	1.3	46
115	Obesity and Liver Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 212-221.	1.3	46
116	Multiple common and rare variants of <i>ABCG2</i> cause gout. RMD Open, 2017, 3, e000464.	3.8	46
117	Impact of menstrual and reproductive factors on breast cancer risk in Japan: Results of the JACC study. Cancer Science, 2005, 96, 57-62.	3.9	45
118	Green Tea Consumption and Gastric Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 335-346.	1.3	45
119	Obesity/Weight Gain and Breast Cancer Risk: Findings From the Japan Collaborative Cohort Study for the Evaluation of Cancer Risk. Journal of Epidemiology, 2013, 23, 139-145.	2.4	45
120	Oxidized human serum albumin as a possible correlation factor for atherosclerosis in a rural Japanese population: the results of the Yakumo Study. Environmental Health and Preventive Medicine, 2018, 23, 1.	3.4	45
121	Risk factors for sudden deafness: a case-control study. Auris Nasus Larynx, 1997, 24, 265-270.	1.2	44
122	Evaluation Based on Systematic Review of Epidemiological Evidence Among Japanese Populations: Tobacco Smoking and Total Cancer Risk. Japanese Journal of Clinical Oncology, 2005, 35, 404-411.	1.3	44
123	Risk modification in lung cancer by a dietary intake of preserved foods and soyfoods: findings from a case-control study in Okinawa, Japan. Lung Cancer, 1999, 25, 147-159.	2.0	43
124	The Japanese food score and risk of all-cause, CVD and cancer mortality: the Japan Collaborative Cohort Study. British Journal of Nutrition, 2018, 120, 464-471.	2.3	43
125	Lung Cancer Mortality and Serum Levels of Carotenoids, Retinol, Tocopherols, and Folic Acid in Men and Women: a Case-Control Study Nested in the JACC Study. Journal of Epidemiology, 2005, 15, S140-S149.	2.4	42
126	Hyperuricemia in acute gastroenteritis is caused by decreased urate excretion via ABCG2. Scientific Reports, 2016, 6, 31003.	3.3	42

#	Article	IF	Citations
127	The Relationship between Smoking Habits and Serum Levels of 8-OHdG, Oxidized LDL Antibodies, Mn-SOD and Carotenoids in Rural Japanese Residents Journal of Epidemiology, 2003, 13, 29-37.	2.4	41
128	Correlation of pathological grade and tumor stage of urothelial carcinomas with CD109 expression. Pathology International, 2010, 60, 735-743.	1.3	41
129	Annual Dialysis Data Report 2014, JSDT Renal Data Registry (JRDR). Renal Replacement Therapy, 2017, 3, .	0.7	41
130	Study Profile of the Japan Multi-institutional Collaborative Cohort (J-MICC) Study. Journal of Epidemiology, 2021, 31, 660-668.	2.4	41
131	Serum Insulin-like Growth Factors, Insulin-like Growth Factor-binding Protein-3, and Risk of Lung Cancer Death: A Case-control Study Nested in the Japan Collaborative Cohort (JACC) Study. Japanese Journal of Cancer Research, 2002, 93, 1279-1286.	1.7	40
132	Risk Factors for Kidney Cancer in a Japanese Population: Findings from the JACC Study. Journal of Epidemiology, 2005, 15, S203-S211.	2.4	40
133	Physical Activity and Colorectal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 2-13.	1.3	40
134	Decrease in risk of lung cancer death in Japanese men after smoking cessation by age at quitting: Pooled analysis of three large-scale cohort studies. Cancer Science, 2007, 98, 584-589.	3.9	39
135	Alcohol drinking and primary liver cancer: A pooled analysis of four Japanese cohort studies. International Journal of Cancer, 2012, 130, 2645-2653.	5.1	39
136	Genomeâ€wide association study identifies gastric cancer susceptibility loci at 12q24.11â€12 and 20q11.21. Cancer Science, 2018, 109, 4015-4024.	3.9	39
137	Baseline data of Shizuoka area in the Japan Multi-Institutional Collaborative Cohort Study (J-MICC) Tj ETQq1 1 0.	784314 rg	:BT ₃ Overlock
138	Maternal Active and Passive Smoking and Fetal Growth: A Prospective Study in Nagoya, Japan Journal of Epidemiology, 2000, 10, 335-343.	2.4	38
139	Effect of Physical Activity on Breast Cancer Risk: Findings of the Japan Collaborative Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 3396-3401.	2.5	38
140	Smoking and serum CA19â€9 levels according to <i>Lewis</i> and <i>secretor</i> genotypes. International Journal of Cancer, 2008, 123, 2880-2884.	5.1	37
141	Alcohol Drinking and Gastric Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2008, 38, 8-25.	1.3	37
142	Intake Frequency of Fish and Serum Levels of Long-chain n-3 Fatty Acids: A Cross-sectional Study within the Japan Collaborative Cohort Study. Journal of Epidemiology, 2005, 15, 211-218.	2.4	36
143	Effects of a low-carbohydrate diet on glycemic control in outpatients with severe type 2 diabetes. Nutrition and Metabolism, 2009, 6, 21.	3.0	36
144	Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. Medicine and Science in Sports and Exercise, 2018, 50, 2433-2441.	0.4	36

#	Article	IF	CITATIONS
145	Clinical impact of prevalence and severity of COPD on the decision-making process for therapeutic management of lung cancer patients. BMC Pulmonary Medicine, 2014, 14, 14.	2.0	35
146	Association of Serum Phytoestrogen Concentration and Dietary Habits in a Sample Set of the JACC Study. Journal of Epidemiology, 2005, 15, S196-S202.	2.4	34
147	Serum Folate and Methylenetetrahydrofolate Reductase (MTHFR) C677T Polymorphism Adjusted for Folate Intake. Journal of Epidemiology, 2008, 18, 125-131.	2.4	34
148	The Determination of Bile Leakage in Complex Hepatectomy Based on the Guidelines of the International Study Group of Liver Surgery. World Journal of Surgery, 2014, 38, 168-176.	1.6	34
149	Risk factors for IgA nephropathy: A case-control study in Japan. American Journal of Kidney Diseases, 1999, 33, 738-745.	1.9	33
150	Dietary Habits and Risk of Urothelial Cancer Incidence in the JACC Study. Journal of Epidemiology, 2005, 15, S190-S195.	2.4	33
151	A common variant of leucine-rich repeat-containing 16A (LRRC16A) gene is associated with gout susceptibility. Human Cell, 2014, 27, 1-4.	2.7	33
152	Coffee Consumption and Risk of Colorectal Cancer: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2014, 24, 370-378.	2.4	33
153	DUPLEX POLYMERASE CHAIN REACTION WITH CONFRONTING TWO-PAIR PRIMERS (PCR-CTPP) FOR GENOTYPING ALCOHOL DEHYDROGENASE Â SUBUNIT (ADH2) AND ALDEHYDE DEHYDROGENASE 2 (ALDH2). Alcohol and Alcoholism, 2003, 38, 407-410.	1.6	32
154	Pro-/anti-inflammatory cytokine gene polymorphisms and chronic kidney disease: a cross-sectional study. BMC Nephrology, 2012, 13, 2.	1.8	32
155	Paravertebral block via the surgical field versus epidural block for patients undergoing thoracotomy: a randomized clinical trial. Surgery Today, 2013, 43, 963-969.	1.5	32
156	A genome-wide association study in the Japanese population identifies the 12q24 locus for habitual coffee consumption: The J-MICC Study. Scientific Reports, 2018, 8, 1493.	3.3	32
157	Dietary Inflammatory Index Is Associated with Risk of All-Cause and Cardiovascular Disease Mortality but Not with Cancer Mortality in Middle-Aged and Older Japanese Adults. Journal of Nutrition, 2019, 1451-1459.	2.9	32
158	Sleep disturbance and its correlates among elderly Japanese. Archives of Gerontology and Geriatrics, 2000, 30, 85-100.	3.0	31
159	Perceived Psychologic Stress and Colorectal Cancer Mortality: Findings From the Japan Collaborative Cohort Study. Psychosomatic Medicine, 2005, 67, 72-77.	2.0	31
160	Impact of the Alcohol-Dehydrogenase (ADH) 1 Cand ADH1 Bpolymorphisms on drinking behavior in nonalcoholic Japanese. Human Mutation, 2007, 28, 506-510.	2.5	31
161	Percutaneous biliary drainage is oncologically inferior to endoscopic drainage: a propensity score matching analysis in resectable distal cholangiocarcinoma. Journal of Gastroenterology, 2016, 51, 608-619.	5.1	31
162	Cigarette smoking and bladder cancer risk: an evaluation based on a systematic review of epidemiologic evidence in the Japanese population. Japanese Journal of Clinical Oncology, 2016, 46, 273-283.	1.3	31

#	Article	IF	Citations
163	Dietary patterns and breast cancer risk in a prospective Japanese study. Breast Cancer, 2017, 24, 152-160.	2.9	31
164	Risk Factors for Breast Cancer among Japanese Women in Tokyo : A Case-Control Study. Journal of Epidemiology, 1994, 4, 65-71.	2.4	30
165	The association between tumour necrosis factor- $\hat{l}\pm$ gene polymorphism and the susceptibility to rugal hyperplastic gastritis and gastric carcinoma. European Journal of Gastroenterology and Hepatology, 2004, 16, 693-700.	1.6	30
166	Alcohol Drinking and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2007, 37, 568-574.	1.3	30
167	Body-Mass Index and Pancreatic Cancer Incidence: A Pooled Analysis of Nine Population-Based Cohort Studies With More Than 340,000 Japanese Subjects. Journal of Epidemiology, 2018, 28, 245-252.	2.4	30
168	Tooth loss and pneumonia mortality: A cohort study of Japanese dentists. PLoS ONE, 2018, 13, e0195813.	2.5	30
169	Serum oxidized low-density lipoprotein levels and risk of colorectal cancer: a case-control study nested in the Japan Collaborative Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2004, 13, 1781-7.	2.5	30
170	Effect of familial history and smoking on common cancer risks in Japan. Cancer, 2007, 109, 2116-2123.	4.1	29
171	No associations of Toll-like receptor 2 (TLR2) -196 to -174del polymorphism with the risk of Helicobacter pylori seropositivity, gastric atrophy, and gastric cancer in Japanese. Gastric Cancer, 2010, 13, 251-257.	5. 3	29
172	Dietary intakes of fat and total mortality among Japanese populations with a low fat intake: the Japan Collaborative Cohort (JACC) Study. Nutrition and Metabolism, 2014, 11, 12.	3.0	29
173	The number of metabolic syndrome components is a good risk indicator for both early- and late-stage kidney damage. Nutrition, Metabolism and Cardiovascular Diseases, 2014, 24, 277-285.	2.6	29
174	Reproducibility and validity of food group intake in a short food frequency questionnaire for the middle-aged Japanese population. Environmental Health and Preventive Medicine, 2021, 26, 28.	3.4	29
175	Prognostic Significance of Selected Lifestyle Factors in Urinary Bladder Cancer. Japanese Journal of Cancer Research, 1993, 84, 1223-1229.	1.7	28
176	Smoking and Colorectal Cancer in a Non-Western Population: a Prospective Cohort Study in Japan. Journal of Epidemiology, 2003, 13, 323-332.	2.4	28
177	Soy consumption reduces the risk of nonâ€smallâ€cell lung cancers with <i>epidermal growth factor receptor</i> mutations among Japanese. Cancer Science, 2008, 99, 1202-1208.	3.9	28
178	An Overview of Genetic Polymorphisms and Pancreatic Cancer Risk in Molecular Epidemiologic Studies. Journal of Epidemiology, 2011, 21, 2-12.	2.4	28
179	Diabetes Mellitus and Liver Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2014, 44, 986-999.	1.3	28
180	Ethnic Differences in ATP-binding Cassette Transporter, Sub-family G, Member 2 (ABCG2/BCRP): Genotype Combinations and Estimated Functions. Drug Metabolism and Pharmacokinetics, 2014, 29, 490-492.	2.2	28

#	Article	IF	Citations
181	Cigarette smoking and the risk of head and neck cancer in the Japanese population: a systematic review and meta-analysis. Japanese Journal of Clinical Oncology, 2016, 46, 580-595.	1.3	28
182	Epidemiological analysis of the association between hearing and barium in humans. Journal of Exposure Science and Environmental Epidemiology, 2016, 26, 488-493.	3.9	28
183	Decrease in Risk of Lung Cancer Death in Males after Smoking Cessation by Age at Quitting: Findings from the JACC Study. Japanese Journal of Cancer Research, 2001, 92, 821-828.	1.7	27
184	Serum Carotenoids, Retinol, and Tocopherols, and Colorectal Cancer Risk in a Japanese Cohort: Effect Modification by Sex for Carotenoids. Nutrition and Cancer, 2005, 51, 13-24.	2.0	27
185	Meat, milk, saturated fatty acids, the Pro12Ala and C161T polymorphisms of the PPAR? gene and colorectal cancer risk in Japanese. Cancer Science, 2006, 97, 1226-1235.	3.9	27
186	Associations of Periodontal Damage and Tooth Loss with Atherogenic Factors among Patients with Type 2 Diabetes Mellitus. Internal Medicine, 2007, 46, 1359-1364.	0.7	27
187	Lung Cancer Risk and Consumption of Vegetables and Fruit: An Evaluation Based on a Systematic Review of Epidemiological Evidence from Japan. Japanese Journal of Clinical Oncology, 2011, 41, 693-708.	1.3	27
188	A past history of gastric ulcers and Helicobacter pylori infection increase the risk of gastric malignant lymphoma. Carcinogenesis, 2006, 27, 1391-1397.	2.8	26
189	Rationale, Design, and Profiles of the New Integrated Suburban Seniority Investigation (NISSIN) Project: A Study of an Age-Specific, Community-Based Cohort of Japanese Elderly. Journal of Epidemiology, 2009, 19, 237-243.	2.4	26
190	Dose-Dependent Neurologic Abnormalities in Workers Exposed to 1-Bromopropane. Journal of Occupational and Environmental Medicine, 2010, 52, 769-777.	1.7	26
191	Significant Association between Serum Interleukin-6 and <i>Helicobacter pylori</i> Antibody Levels among <i>H. pylori</i> Positive Japanese Adults. Mediators of Inflammation, 2013, 2013, 1-5.	3.0	26
192	Within-visit blood pressure variability is associated with prediabetes and diabetes. Scientific Reports, 2015, 5, 7964.	3.3	26
193	No association between Helicobacter pylori infection and diabetes mellitus among a general Japanese population: a cross-sectional study. SpringerPlus, 2015, 4, 602.	1.2	26
194	Seasonal Allergic Rhinoconjunctivitis and Fatty Acid Intake. Annals of Epidemiology, 2001, 11, 59-64.	1.9	25
195	Effectiveness of Cervical Cancer Screening Over Cervical Cancer Mortality Among Japanese Women. Japanese Journal of Clinical Oncology, 2006, 36, 511-518.	1.3	25
196	Psychological attitudes and risk of breast cancer in Japan: a prospective study. Cancer Causes and Control, 2007, 18, 259-267.	1.8	25
197	No association between the frequency of forest walking and blood pressure levels or the prevalence of hypertension in a cross-sectional study of a Japanese population. Environmental Health and Preventive Medicine, 2011, 16, 299-306.	3.4	25
198	Associations of <i>apolipoprotein A5 (APOA5), glucokinase (GCK)</i> and <i>glucokinase regulatory protein (GCKR)</i> polymorphisms and lifestyle factors with the risk of dyslipidemia and dysglycemia in Japanese $\hat{a} \in \mathbb{Z}$ a cross-sectional data from the J-MICC Study. Endocrine Journal, 2012, 59, 589-599.	1.6	25

#	Article	IF	CITATIONS
199	Visit-to-visit blood pressure variability is a marker of cardiac diastolic function and carotid atherosclerosis. BMC Cardiovascular Disorders, 2014, 14, 188.	1.7	25
200	Gastric Cancer Risk and Erythrocyte Composition of Docosahexaenoic Acid with Anti-inflammatory Effects. Cancer Epidemiology Biomarkers and Prevention, 2007, 16, 2406-2415.	2.5	24
201	Subtype-specific gout susceptibility loci and enrichment of selection pressure on ABCG2 and ALDH2 identified by subtype genome-wide meta-analyses of clinically defined gout patients. Annals of the Rheumatic Diseases, 2020, 79, 657-665.	0.9	24
202	Helicobacter Pylori Seropositivity and the Myeloperoxidase G-463A Polymorphism in Combination with Interleukin-1B C-31T in Japanese Health Checkup Examinees. Japanese Journal of Clinical Oncology, 2003, 33, 192-197.	1.3	23
203	Significant interaction between LRP2 rs2544390 in intron 1 and alcohol drinking for serum uric acid levels among a Japanese population. Gene, 2012, 503, 131-136.	2.2	23
204	Vegetable consumption and colorectal cancer risk: an evaluation based on a systematic review and meta-analysis among the Japanese population. Japanese Journal of Clinical Oncology, 2015, 45, 973-979.	1.3	23
205	International pooled study on diet and bladder cancer: the bladder cancer, epidemiology and nutritional determinants (BLEND) study: design and baseline characteristics. Archives of Public Health, 2016, 74, 30.	2.4	23
206	Metallothionein <i> MT2A </i> > A-5G Polymorphism as a Risk Factor for Chronic Kidney Disease and Diabetes: Cross-Sectional and Cohort Studies. Toxicological Sciences, 2016, 152, 181-193.	3.1	23
207	Insulin-like growth factor-related components and the risk of liver cancer in a nested case-control study. Tumor Biology, 2016, 37, 15125-15132.	1.8	23
208	Risk Factors for Renal Cell Carcinoma in a Japanese Population. Asian Pacific Journal of Cancer Prevention, 2014, 15, 9065-9070.	1.2	23
209	Relationship between body mass index and the risk of ovarian cancer in the Japanese population: Findings from the Japanese Collaborate Cohort (JACC) study. Journal of Obstetrics and Gynaecology Research, 2005, 31, 452-458.	1.3	22
210	Prospective study of alcohol consumption and breast cancer risk in Japanese women. International Journal of Cancer, 2005, 116, 779-783.	5.1	22
211	Significant association of serum uric acid levels with SLC2A9 rs11722228 among a Japanese population. Molecular Genetics and Metabolism, 2011, 103, 378-382.	1.1	22
212	Effects of a moderate low-carbohydrate diet on preferential abdominal fat loss and cardiovascular risk factors in patients with type 2 diabetes. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2011, 4, 167.	2.4	22
213	Health Benefits of Daily Walking on Mortality Among Younger-Elderly Men With or Without Major Critical Diseases in the New Integrated Suburban Seniority Investigation Project: A Prospective Cohort Study. Journal of Epidemiology, 2015, 25, 609-616.	2.4	22
214	Coffee drinking and colorectal cancer risk: an evaluation based on a systematic review and meta-analysis among the Japanese population. Japanese Journal of Clinical Oncology, 2016, 46, 781-787.	1.3	22
215	Cigarette Smoking and other Risk Factors for Kidney Cancer Death in a Japanese Population: Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC study). Asian Pacific Journal of Cancer Prevention, 2013, 14, 6523-6528.	1.2	22
216	Lifestyle determinants for social activity levels among the Japanese elderly. Archives of Gerontology and Geriatrics, 1996, 22, 271-286.	3.0	21

#	Article	IF	Citations
217	Reproducibility and Validity of a Simple Checklist-type Questionnaire for Food Intake and Dietary Behavior. Journal of Epidemiology, 2003, 13, 235-245.	2.4	21
218	Inverse correlation between serum interleukin-6 and iron levels among Japanese adults: a cross-sectional study. BMC Hematology, 2014, 14, 6.	2.6	21
219	Association of gait speed with mortality among the Japanese elderly in the New Integrated Suburban Seniority Investigation Project: a prospective cohort study. Age and Ageing, 2015, 44, 153-157.	1.6	21
220	Adjustment of Cell-Type Composition Minimizes Systematic Bias in Blood DNA Methylation Profiles Derived by DNA Collection Protocols. PLoS ONE, 2016, 11, e0147519.	2.5	21
221	Smoking Habits, Local Brand Cigarettes and Lung Cancer Risk in Okinawa, Japan Journal of Epidemiology, 1997, 7, 99-105.	2.4	20
222	Updated Information on Risk Factors for Lung Cancer: Findings from the JACC Study. Journal of Epidemiology, 2005, 15, S134-S139.	2.4	20
223	Alcohol Consumption and Colorectal Cancer Risk: Findings from the JACC Study. Journal of Epidemiology, 2005, 15, S173-S179.	2.4	20
224	Alcohol Consumption and Lung Cancer Mortality in Japanese Men: Results from Japan Collaborative Cohort (JACC) Study. Journal of Epidemiology, 2006, 16, 49-56.	2.4	20
225	Alcohol Drinking and Lung Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2007, 37, 168-174.	1.3	20
226	Associations of a PTPN11 G/A polymorphism at intron 3 with Helicobactor pyloriseropositivity, gastric atrophy and gastric cancer in Japanese. BMC Gastroenterology, 2009, 9, 51.	2.0	20
227	Breastfeeding and Breast Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2012, 42, 124-130.	1.3	20
228	Smoking cessation and subsequent risk of cancer: A pooled analysis of eight population-based cohort studies in Japan. Cancer Epidemiology, 2017, 51, 98-108.	1.9	20
229	Weight Gain and Alcohol Drinking Associations with Breast Cancer Risk in Japanese Postmenopausal Women - Results from the Japan Collaborative Cohort (JACC) Study. Asian Pacific Journal of Cancer Prevention, 2016, 17, 1437-1443.	1.2	20
230	Renal Hyperfiltration in Prediabetes Confirmed by Fasting Plasma Glucose and Hemoglobin A1c. Renal Failure, 2012, 34, 1084-1090.	2.1	19
231	Fish Consumption and Colorectal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. Japanese Journal of Clinical Oncology, 2013, 43, 935-941.	1.3	19
232	Low Intake of Vegetables and Fruits and Risk of Colorectal Cancer: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2014, 24, 353-360.	2.4	19
233	Blood-borne miRNA profile-based diagnostic classifier for lung adenocarcinoma. Scientific Reports, 2016, 6, 31389.	3.3	19
234	Circulating miR-21, miR-29a, and miR-126 are associated with premature death risk due to cancer and cardiovascular disease: the JACC Study. Scientific Reports, 2021, 11, 5298.	3.3	19

#	Article	IF	CITATIONS
235	Depressive Mood and Suicide Among Middle-Aged Workers: Findings from a Prospective Cohort Study in Nagoya, Japan Journal of Epidemiology, 2000, 10, 173-178.	2.4	18
236	A prospective study on the possible association between having children and colon cancer risk: Findings from the JACC Study. Cancer Science, 2004, 95, 243-247.	3.9	18
237	A prospective study of reproductive and menstrual factors and colon cancer risk in Japanese women: Findings from the JACC study. Cancer Science, 2004, 95, 602-607.	3.9	18
238	One-Year Period Prevalence of Oral Aphthous Ulcers and Oral Health-Related Quality of Life in Patients with Behçet's Disease. Genetics Research International, 2014, 2014, 1-8.	2.0	18
239	Genome-Wide Association Study of Renal Function Traits: Results from the Japan Multi-Institutional Collaborative Cohort Study. American Journal of Nephrology, 2018, 47, 304-316.	3.1	18
240	Comparison of the analgesic effects of modified continuous intercostal block and paravertebral block under surgeon's direct vision after video-assisted thoracic surgery: a randomized clinical trial. General Thoracic and Cardiovascular Surgery, 2018, 66, 425-431.	0.9	18
241	Risk Factors for IgA Nephropathy: A Case-Control Study with Incident Cases in Japan. Nephron, 2002, 90, 16-23.	1.8	17
242	Trends in Asthma Mortality in Japan. Journal of Asthma, 2002, 39, 633-639.	1.7	17
243	<i>ATP-binding cassette transporter A1</i> (<i>ABCA1</i>) R219K (G1051A, rs2230806) polymorphism and serum high-density lipoprotein cholesterol levels in a large Japanese population: cross-sectional data from the Daiko Study. Endocrine Journal, 2015, 62, 543-549.	1.6	17
244	Leisure-time physical activity and risk of disability incidence: A 12-year prospective cohort study among young elderly of the same age at baseline. Journal of Epidemiology, 2017, 27, 538-545.	2.4	17
245	Case-control study of diabetes-related genetic variants and pancreatic cancer risk in Japan. World Journal of Gastroenterology, 2014, 20, 17456.	3.3	17
246	Favorable Life-Style Modification and Attenuation of Cardiovascular Risk Factors. Japanese Circulation Journal, 1999, 63, 184-188.	1.0	16
247	eNOS genotype modifies the effect of leisure-time physical activity on serum triglyceride levels in a Japanese population. Lipids in Health and Disease, $2012, 11, 150$.	3.0	16
248	A polymorphism near MC4R gene (rs17782313) is associated with serum triglyceride levels in the general Japanese population: the J-MICC Study. Endocrine, 2014, 47, 81-89.	2.3	16
249	Genome-wide association study-identified SNPs (rs3790844, rs3790843) in the NR5A2 gene and risk of pancreatic cancer in Japanese. Scientific Reports, 2015, 5, 17018.	3.3	16
250	Arsenic levels in cutaneous appendicular organs are correlated with digitally evaluated hyperpigmented skin of the forehead but not the sole in Bangladesh residents. Journal of Exposure Science and Environmental Epidemiology, 2018, 28, 64-68.	3.9	16
251	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep durationin the Japanese population. Sleep, 2019, 42, .	1.1	16
252	Dietary fish and $i\%$ -3 polyunsaturated fatty acids are associated with leukocyte ABCA1 DNA methylation levels. Nutrition, 2021, 81, 110951.	2.4	16

#	Article	IF	CITATIONS
253	Significance of Urinalysis for Subsequent Kidney and Urinary Tract Disorders in Mass Screening of Adults Internal Medicine, 1995, 34, 475-480.	0.7	15
254	Dietary Habits and Risk of Urothelial Cancer Death in a Large-Scale Cohort Study (JACC Study) in Japan. Nutrition and Cancer, 2004, 50, 33-39.	2.0	15
255	Inspiratory capacity as a preoperative assessment of patients undergoing thoracic surgery. Interactive Cardiovascular and Thoracic Surgery, 2012, 14, 560-564.	1.1	15
256	Associations of daily walking and television viewing time with liver cancer mortality: findings from the Japan Collaborative Cohort Study. Cancer Causes and Control, 2014, 25, 787-793.	1.8	15
257	Matrix metalloproteinaseÂ9 gene polymorphisms are associated with a multiple family history of gastric cancer. Gastric Cancer, 2017, 20, 246-253.	5.3	15
258	Lower risk of progression from prediabetes to diabetes with health checkup with lifestyle education: Japan Ningen Dock study. Nutrition, Metabolism and Cardiovascular Diseases, 2017, 27, 679-687.	2.6	15
259	Modification of the Associations of Alcohol Intake With Serum Low-Density Lipoprotein Cholesterol and Triglycerides by <i>ALDH2</i> and <i>ADH1B</i> Polymorphisms in Japanese Men. Journal of Epidemiology, 2018, 28, 185-193.	2.4	15
260	Gene-Gene Combination Effect and Interactions among ABCA1, APOA1, SR-B1, and CETP Polymorphisms for Serum High-Density Lipoprotein-Cholesterol in the Japanese Population. PLoS ONE, 2013, 8, e82046.	2.5	15
261	Estimated Prevalence of Sjogren's Syndrome in Japan : Findings from a Nationwide Epidemiological Survey. Journal of Epidemiology, 1995, 5, 125-129.	2.4	14
262	Out-of-Hospital Cardiac Arrest and Survival-An Epidemiological Analysis of Emergency Service Reports in a Large City in Japan Circulation Journal, 2004, 68, 603-609.	1.6	14
263	PRKCH gene polymorphism is associated with the risk of severe gastric atrophy. Gastric Cancer, 2010, 13, 90-94.	5.3	14
264	Serum uric acid distribution according to SLC22A12 W258X genotype in a cross-sectional study of a general Japanese population. BMC Medical Genetics, 2011, 12, 33.	2.1	14
265	Prospective cohort study on television viewing time and incidence of lung cancer: findings from the Japan Collaborative Cohort Study. Cancer Causes and Control, 2013, 24, 1547-1553.	1.8	14
266	Association of exposure level to passive smoking with hypertension among lifetime nonsmokers in Japan: a cross-sectional study. Medicine (United States), 2018, 97, e13241.	1.0	14
267	Associations between diet and mental health using the 12-item General Health Questionnaire: cross-sectional and prospective analyses from the Japan Multi-Institutional Collaborative Cohort Study. Nutrition Journal, 2020, 19, 2.	3.4	14
268	Association between plasma levels of homocysteine, folate, and vitamin B12, and dietary folate intake and hypertension in a cross-sectional study. Scientific Reports, 2020, 10, 18499.	3.3	14
269	Prediction of dialysis patients in Japan : Based on Japanese Society for Dialysis Therapy Registry. Nihon Toseki Igakkai Zasshi, 2012, 45, 599-613.	0.1	14
270	SERUM CAROTENOIDS AND OTHER ANTIOXIDATIVE SUBSTANCES ASSOCIATED WITH UROTHELIAL CANCER RISK IN A NESTED CASE-CONTROL STUDY IN JAPANESE MEN. Journal of Urology, 2005, 173, 1502-1506.	0.4	13

#	Article	IF	Citations
271	Alcohol Drinking and Total Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence among the Japanese Population. Japanese Journal of Clinical Oncology, 2007, 37, 692-700.	1.3	13
272	Lung cancer mortality and body mass index in a Japanese cohort: findings from the Japan Collaborative Cohort Study (JACC Study). Cancer Causes and Control, 2007, 18, 229-234.	1.8	13
273	Tooth loss and risk of hip fracture: a prospective study of male <scp>J</scp> apanese dentists. Community Dentistry and Oral Epidemiology, 2013, 41, 48-54.	1.9	13
274	Body Mass Index and Weight Change During Adulthood Are Associated With Increased Mortality From Liver Cancer: The JACC Study. Journal of Epidemiology, 2013, 23, 219-226.	2.4	13
275	Risk of lung cancer and consumption of vegetables and fruit in Japanese: A pooled analysis of cohort studies in Japan. Cancer Science, 2015, 106, 1057-1065.	3.9	13
276	Risk stratification by the lower limit of normal of FEV1/FVC for postoperative outcomes in patients with COPD undergoing thoracic surgery. Respiratory Investigation, 2015, 53, 117-123.	1.8	13
277	Insulin-like growth factor-1, IGF binding protein-3, and the risk of esophageal cancer in a nested case-control study. World Journal of Gastroenterology, 2017, 23, 3488.	3.3	13
278	Cigarette smoking and the risk of ovarian cancer in the Japanese population: Findings from the Japanese Collaborate Cohort study. Journal of Obstetrics and Gynaecology Research, 2005, 31, 144-151.	1.3	12
279	A prospective study of educational background and breast cancer among Japanese women. Cancer Causes and Control, 2008, 19, 931-937.	1.8	12
280	Associations of plasma IL-8 levels with <i>Helicobacter pylori</i> seropositivity, gastric atrophy, and <i>IL-8</i> T-251A genotypes. Epidemiology and Infection, 2010, 138, 512-518.	2.1	12
281	Matrix Metalloproteinase-9 Gene Polymorphisms and Chronic Kidney Disease. American Journal of Nephrology, 2012, 36, 444-450.	3.1	12
282	Transforming Growth Factor-β1 as a Predictor for the Development of Hepatocellular Carcinoma: A Nested Case–Controlled Study. EBioMedicine, 2016, 12, 68-71.	6.1	12
283	Peritoneal Dialysis Registry With 2013 Survey Report. Therapeutic Apheresis and Dialysis, 2016, 20, 557-568.	0.9	12
284	Clinical impact of the lower limit of normal of FEV1/FVC on survival in lung cancer patients undergoing thoracic surgery. Respiratory Investigation, 2016, 54, 184-192.	1.8	12
285	The impact of musculoskeletal diseases on mortality–comparison with internal diseases: A 15-year longitudinal study. Journal of Orthopaedic Science, 2017, 22, 1126-1131.	1.1	12
286	Dietary Antioxidant Micronutrients and All-Cause Mortality: The Japan Collaborative Cohort Study for Evaluation of Cancer Risk. Journal of Epidemiology, 2018, 28, 388-396.	2.4	12
287	Lifestyle and psychosocial factors and a decline in competence in daily living among Japanese early elderly people: from an age-specified community-based cohort study (NISSIN project). Environmental Health and Preventive Medicine, 2019, 24, 28.	3.4	12
288	Study profile on baseline survey of Daiko Study in the Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study). Nagoya Journal of Medical Science, 2011, 73, 187-95.	0.3	12

#	Article	IF	Citations
289	Prevalence of Helicobacter pylori infection measured with urinary antibody in an urban area of Japan, 2008-2010. Nagoya Journal of Medical Science, 2012, 74, 63-70.	0.3	12
290	Controls for monitoring the deterioration of stored blood samples in the Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study). Nagoya Journal of Medical Science, 2008, 70, 107-15.	0.3	12
291	Prostate cancer risk in relation to insulin-like growth factor (IGF)-I and IGF-binding protein-3: A nested case-control study in large scale cohort study in Japan. Asian Pacific Journal of Cancer Prevention, 2009, 10 Suppl, 57-61.	1.2	12
292	Acute metabolic responses to a high-carbohydrate meal in outpatients with type 2 diabetes treated with a low-carbohydrate diet: a crossover meal tolerance study. Nutrition and Metabolism, 2009, 6, 52.	3.0	11
293	Longitudinal Evaluation of Multi-phasic, Odontological and Nutritional Associations in Dentists (LEMONADE Study): Study Design and Profile of Nationwide Cohort Participants at Baseline. Journal of Epidemiology, 2009, 19, 72-80.	2.4	11
294	Detailed Analysis of Japanese Population Substructure with a Focus on the Southwest Islands of Japan. PLoS ONE, 2012, 7, e35000.	2.5	11
295	Genome-wide association study of genetic factors related to confectionery intake: Potential roles of the <i>ADIPOQ </i> gene. Obesity, 2013, 21, 2413-2419.	3.0	11
296	Polymorphisms in genes encoding antioxidant enzymes (SOD2, CAT, GPx, TXNRD, SEPP1, SEP15 and SELS) and risk of chronic kidney disease in Japanese - cross-sectional data from the J-MICC study. Journal of Clinical Biochemistry and Nutrition, 2013, 53, 15-20.	1.4	11
297	GCK, GCKR polymorphisms and risk of chronic kidney disease in Japanese individuals: data from the J-MICC Study. Journal of Nephrology, 2014, 27, 143-149.	2.0	11
298	Association of decrease in carbohydrate intake with reduction in abdominal fat during 3-month moderate low-carbohydrate diet among non-obese Japanese patients with type 2 diabetes. Metabolism: Clinical and Experimental, 2015, 64, 618-625.	3.4	11
299	Upper-normal waist circumference is a risk marker for metabolic syndrome in normal-weight subjects. Nutrition, Metabolism and Cardiovascular Diseases, 2016, 26, 67-76.	2.6	11
300	Genetic Variants of <i>RAMP2</i> and <i>CLR</i> are Associated with Stroke. Journal of Atherosclerosis and Thrombosis, 2017, 24, 1267-1281.	2.0	11
301	Is hemodialysis itself a risk factor for dementia? An analysis of nationwide registry data of patients on maintenance hemodialysis in Japan. Renal Replacement Therapy, 2018, 4, .	0.7	11
302	The associations of dietary patterns with all-cause mortality and other lifestyle factors in the elderly: An age-specific prospective cohort study. Clinical Nutrition, 2019, 38, 288-296.	5.0	11
303	The interaction between ABCA1 polymorphism and physical activity on the HDL-cholesterol levels in a Japanese population. Journal of Lipid Research, 2020, 61, 86-94.	4.2	11
304	Polymorphisms in <i>PPAR</i> Genes (<i>PPARD</i> , <i>PPARG</i> , and <i>PPARGC1A</i>) and the Risk of Chronic Kidney Disease in Japanese: Cross-Sectional Data from the J-MICC Study. PPAR Research, 2013, 2013, 1-8.	2.4	10
305	Impact of Thin-Section Computed Tomography-Determined Combined Pulmonary Fibrosis and Emphysema on Outcomes Among Patients With Resected Lung Cancer. Annals of Thoracic Surgery, 2016, 102, 440-447.	1.3	10
306	Smoking and subsequent risk of acute myeloid leukaemia: A pooled analysis of 9 cohort studies in Japan. Hematological Oncology, 2018, 36, 262-268.	1.7	10

#	Article	IF	CITATIONS
307	The association between self-rated health and high-sensitivity C-reactive protein level: a cross-sectional and 5-year longitudinal study. BMC Public Health, 2018, 18, 1380.	2.9	10
308	Epidemiology of Diseases of Unknown Etiology, Specified as "Intractable Diseases― Journal of Epidemiology, 1996, 6, 87-94.	2.4	9
309	Lifestyle risk factors for intrahepatic stone: Findings from a case–control study in an endemic area, Taiwan. Journal of Gastroenterology and Hepatology (Australia), 2008, 23, 1075-1081.	2.8	9
310	Significant association of RUNX3 T/A polymorphism at intron 3 (rs760805) with the risk of gastric atrophy in Helicobacter pylori seropositive Japanese. Journal of Gastroenterology, 2009, 44, 1165-1171.	5.1	9
311	Association between <i>KLK3</i> rs2735839 G/A Polymorphism and Serum PSA Levels in Japanese Men. Urologia Internationalis, 2012, 89, 39-44.	1.3	9
312	MTHFR, MTR and MTRR polymorphisms and risk of chronic kidney disease in Japanese: cross-sectional data from the J-MICC Study. International Urology and Nephrology, 2013, 45, 1613-1620.	1.4	9
313	Significant interaction between <i>RETN</i> -420 <i>G/G</i> genotype and lower BMI on decreased risk of type 2 diabetes mellitus (T2DM) in Japanese â€" the J-MICC Study [Rapid Communication]. Endocrine Journal, 2013, 60, 237-243.	1.6	9
314	Diabetes Mellitus is Associated With Low Secretion Rates of Immunoglobulin A in Saliva. Journal of Epidemiology, 2015, 25, 470-474.	2.4	9
315	Associations between polymorphisms of interleukin-6 and related cytokine genes and serum liver damage markers: a cross-sectional study in the Japan Multi-Institutional Collaborative Cohort (J-MICC) Study. Gene, 2015, 557, 158-162.	2.2	9
316	Common variant of ALPK1 is not associated with gout: a replication study. Human Cell, 2015, 28, 1-4.	2.7	9
317	Associations between polymorphisms in folate-metabolizing genes and pancreatic cancer risk in Japanese subjects. BMC Gastroenterology, 2016, 16, 83.	2.0	9
318	Meat subtypes and colorectal cancer risk: A pooled analysis of 6 cohort studies in Japan. Cancer Science, 2019, 110, 3603-3614.	3.9	9
319	A risk score predicting new incidence of hypertension in Japan. Journal of Human Hypertension, 2019, 33, 748-755.	2,2	9
320	Identification of two novel breast cancer loci through large-scale genome-wide association study in the Japanese population. Scientific Reports, 2019, 9, 17332.	3.3	9
321	Association between dietary inflammatory index and serum C-reactive protein concentrations in the Japan Collaborative Cohort Study. Nagoya Journal of Medical Science, 2020, 82, 237-249.	0.3	9
322	Diabetes Mellitus and Risk of Colorectal Cancer Mortality in Japan: the Japan Collaborative Cohort Study. Asian Pacific Journal of Cancer Prevention, 2016, 17, 4681-4688.	1.2	9
323	Dairy products and the risk of developing prostate cancer: A largeâ€scale cohort study (JACC Study) in Japan. Cancer Medicine, 2021, 10, 7298-7307.	2,8	9
324	ALDH2 polymorphism is associated with fasting blood glucose through alcohol consumption in Japanese men. Nagoya Journal of Medical Science, 2016, 78, 183-93.	0.3	9

#	Article	lF	Citations
325	Epidemiology of Intractable Diseases in Japan. Journal of Epidemiology, 1996, 6, 99-109.	2.4	8
326	Glucose Intolerance and Colorectal Cancer Risk in a Nested Case-Control Study among Japanese People. Journal of Epidemiology, 2005, 15, S180-S184.	2.4	8
327	Written Informed Consent for Participation in a Study and Reduction in Consent Rate. Journal of Epidemiology, 2008, 18, 291-294.	2.4	8
328	Reduction in urinary albumin excretion with a moderate low-carbohydrate diet in patients with type 2 diabetes: a 12-month intervention. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2012, 5, 283.	2.4	8
329	Association of genetic polymorphisms with erythrocyte traits: Verification of SNPs reported in a previous GWAS in a Japanese population. Gene, 2018, 642, 172-177.	2.2	8
330	Genome-wide association study (GWAS) of ovarian cancer in Japanese predicted regulatory variants in 22q13.1. PLoS ONE, 2018, 13, e0209096.	2.5	8
331	Circulating insulinâ€like growth factor binding proteinâ€3 and risk of gastrointestinal malignant tumors. Journal of Gastroenterology and Hepatology (Australia), 2019, 34, 2104-2111.	2.8	8
332	A genome-wide association study in Japanese identified one variant associated with a preference for a Japanese dietary pattern. European Journal of Clinical Nutrition, 2021, 75, 937-945.	2.9	8
333	A Proposal for Practical Diagnosis of Renal Hypouricemia: Evidenced from Genetic Studies of Nonfunctional Variants of URAT1/SLC22A12 among 30,685 Japanese Individuals. Biomedicines, 2021, 9, 1012.	3.2	8
334	Lack of Associations between Genetic Polymorphisms in GSTM1, GSTT1 and GSTP1 and Pancreatic Cancer Risk: A Multi-Institutional Case-Control Study in Japan. Asian Pacific Journal of Cancer Prevention, 2014, 15, 391-395.	1.2	8
335	Time spent walking or exercising and blood levels of insulin-like growth factor-I (IGF-I) and IGF-binding protein-3 (IGFBP-3): A large-scale cross-sectional study in the Japan Collaborative Cohort study. Asian Pacific Journal of Cancer Prevention, 2009, 10 Suppl, 23-7.	1.2	8
336	Epidemiological analysis of risk factors for breast cancer in Indonesian females. Medical Journal of Indonesia, 1995, 4, 163.	0.5	7
337	Newly developed ST-T abnormalities on the electrocardiogram and chronologic changes in cardiovascular risk factors. American Journal of Cardiology, 1996, 77, 823-827.	1.6	7
338	Risk factors for sudden unexpected death among workers: A nested case-control study in central Japan. Preventive Medicine, 2001, 33, 99-107.	3.4	7
339	Medical History of Circulatory Diseases and Colorectal Cancer Death in the JACC Study. Journal of Epidemiology, 2005, 15, S168-S172.	2.4	7
340	Association between Decreased Kidney Function and Endotoxin Receptor <i>CD14</i> C-159T Polymorphism among Japanese Health Check-up Examinees. Renal Failure, 2007, 29, 967-972.	2.1	7
341	Polymorphisms of genes involved in lipid metabolism and risk of chronic kidney disease in Japanese - cross-sectional data from the J-MICC study. Lipids in Health and Disease, 2014, 13, 162.	3.0	7
342	Three-graded stratification of carbohydrate restriction by level of baseline hemoglobin A1c for type 2 diabetes patients with a moderate low-carbohydrate diet. Nutrition and Metabolism, 2014, 11, 33.	3.0	7

#	Article	IF	CITATIONS
343	Genetic variants of SLC17A1 are associated with cholesterol homeostasis and hyperhomocysteinaemia in Japanese men. Scientific Reports, 2015, 5, 15888.	3.3	7
344	Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. Journal of Epidemiology, 2017, 27, 331-337.	2.4	7
345	The impact of carbohydrate intake and its sources on hemoglobin A1c levels in Japanese patients with type 2 diabetes not taking anti-diabetic medication. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2018, Volume 11, 53-64.	2.4	7
346	No association between angiotensin I converting enzyme (ACE) I/D polymorphism and gastric cancer risk among Japanese. Nagoya Journal of Medical Science, 2011, 73, 169-75.	0.3	7
347	Associations of Lifestyle Factors with Bone Mineral Density among Male University Students in Japan Journal of Epidemiology, 2003, 13, 48-55.	2.4	6
348	Significant association of urokinase plasminogen activator Pro141Leu with serum lipid profiles in a Japanese population. Gene, 2013, 524, 363-367.	2.2	6
349	A Cross-Sectional Study to Find Out the Relationship of Methylenetetrahydrofolate Reductase (MTHFR) C677T Genotype with Plasma Levels of Folate and Total Homocysteine by Daily Folate Intake in Japanese. Journal of Nutritional Science and Vitaminology, 2014, 60, 231-238.	0.6	6
350	Association between falls and depressive symptoms or visual impairment among Japanese youngâ€old adults. Geriatrics and Gerontology International, 2016, 16, 384-391.	1.5	6
351	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. British Journal of Nutrition, 2021, 126, 1843-1851.	2.3	6
352	Association of skipping breakfast and short sleep duration with the prevalence of metabolic syndrome in the general Japanese population: Baseline data from the Japan Multi-Institutional Collaborative cohort study. Preventive Medicine Reports, 2021, 24, 101613.	1.8	6
353	ASSOCIATIONS BETWEEN BODY MASS INDEX AND SERUM URIC ACID LEVELS IN A JAPANESE POPULATION WERE SIGNIFICANTLY MODIFIED BY LRP2 rs2544390. Nagoya Journal of Medical Science, 2014, 76, 333-9.	0.3	6
354	Association of Serum Carotenoid Concentration and Dietary Habits among the JACC Study Subjects. Journal of Epidemiology, 2005, 15, S220-S227.	2.4	5
355	Association of calcium urolithiasis with urokinase P141L and 3′-UTR C>T polymorphisms in a Japanese population. Urolithiasis, 2013, 41, 47-52.	2.0	5
356	Body Mass Index Is Associated with Hypertension in Japanese Young Elderly Individuals: Findings of the New Integrated Suburban Seniority Investigation. Internal Medicine, 2015, 54, 3121-3125.	0.7	5
357	Effects of <i><scp>IL</scp>6</i> Câ€634G polymorphism on tooth loss and their interaction with smoking habits. Oral Diseases, 2015, 21, 807-813.	3.0	5
358	Bilobar versus unilobar multiple colorectal liver metastases: a propensity score analysis of surgical outcomes and recurrence patterns. Journal of Hepato-Biliary-Pancreatic Sciences, 2017, 24, 153-160.	2.6	5
359	Fruit and vegetable intake and the risk of overall cancer in Japanese: AÂpooled analysis of population-based cohort studies. Journal of Epidemiology, 2017, 27, 152-162.	2.4	5
360	A genome-wide association study on fish consumption in a Japanese populationâ€"the Japan Multi-Institutional Collaborative Cohort study. European Journal of Clinical Nutrition, 2021, 75, 480-488.	2.9	5

#	Article	IF	CITATIONS
361	A study on how a 6-month aerobic exercise program can modify coronary risk factors depending on their severity in middle-aged sedentary women. Environmental Health and Preventive Medicine, 1999, 4, 117-121.	3.4	4
362	Factors associated with the incidence of dialysis. Clinical and Experimental Nephrology, 2013, 17, 890-898.	1.6	4
363	Effects of self-reported calorie restriction on correlations between SIRT1 polymorphisms and body mass index and long-term weight change. Gene, 2016, 594, 16-22.	2.2	4
364	Association of Adiponectin With Cancer and All-Cause Mortality in a Japanese Community-Dwelling Elderly Cohort: A Case-Cohort Study. Journal of Epidemiology, 2018, 28, 367-372.	2.4	4
365	Common variant of BCAS3 is associated with gout risk in Japanese population: the first replication study after gout GWAS in Han Chinese. BMC Medical Genetics, 2018, 19, 96.	2.1	4
366	Thin-section computed tomography-determined usual interstitial pneumonia pattern affects the decision-making process for resection in newly diagnosed lung cancer patients: a retrospective study. BMC Pulmonary Medicine, 2018, 18, 2.	2.0	4
367	Human serum albumin redox state is associated with decreased renal function in a community-dwelling population. American Journal of Physiology - Renal Physiology, 2019, 316, F214-F218.	2.7	4
368	Insulin-like Growth Factor-1, Insulin-like Growth Factor Binding Protein-3 and the Incidence of Malignant Neoplasms in a Nested Case–Control Study. Cancer Prevention Research, 2020, 13, 385-394.	1.5	4
369	DPP4 genetic variants influence baseline prostate-specific antigen levels: the J-MICC study. Nagoya Journal of Medical Science, 2013, 75, 73-80.	0.3	4
370	Serum insulin-like growth factors I and II, insulin-like growth factor binding protein-3 and risk of breast cancer in the Japan Collaborative Cohort study. Asian Pacific Journal of Cancer Prevention, 2009, 10 Suppl, 51-5.	1.2	4
371	Case-control study of oral cancer in Shenyang, Northeastern China. International Journal of Clinical Oncology, 1998, 3, 13-18.	2.2	3
372	Successful Aging and Social Activity in Older Japanese Adults. Journal of Aging and Physical Activity, 2000, 8, 129-139.	1.0	3
373	No association between AICDA 7888 C/T polymorphism, Helicobacter pylori seropositivity, and the risk of atrophic gastritis and gastric cancer in Japanese. Gastric Cancer, 2010, 13, 43-49.	5.3	3
374	Associations of smoking status with other lifestyle behaviors are modified by sex and occupational category among urban civil servants in Japan. Environmental Health and Preventive Medicine, 2016, 21, 539-546.	3.4	3
375	Daily sleep duration and the risk of incident disability among younger elderly Japanese adults in the New Integrated Suburban Seniority Investigation Project: A prospective study using competing event analysis. Geriatrics and Gerontology International, 2019, 19, 945-949.	1.5	3
376	Reducing Carbohydrate from Individual Sources Has Differential Effects on Glycosylated Hemoglobin in Type 2 Diabetes Mellitus Patients on Moderate Low-Carbohydrate Diets. Diabetes and Metabolism Journal, 2021, 45, 390-403.	4.7	3
377	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. Journal of Nutritional Science, 2021, 10, e61.	1.9	3
378	Alcohol intake and stomach cancer risk in Japan: A pooled analysis of six cohort studies. Cancer Science, 2022, 113, 261-276.	3.9	3

#	Article	IF	Citations
379	Macronutrient intakes and serum oestrogen, and interaction with polymorphisms in <i>CYP19A1</i> and <i>HSD17B1</i> genes: a cross-sectional study in postmenopausal Japanese women. British Journal of Nutrition, 2017, 118, 463-472.	2.3	2
380	Annual peritoneal dialysis report 2014, the peritoneal dialysis registry. Renal Replacement Therapy, $2017, 3, .$	0.7	2
381	Calculation of expected remaining lifetime of dialysis patients in Japan. Renal Replacement Therapy, 2020, 6, .	0.7	2
382	Frequency of forest walking is not associated with prevalence of hypertension based on cross-sectional studies of a general Japanese population: a reconfirmation by the J-MICC Daiko Study. Nagoya Journal of Medical Science, 2019, 81, 489-500.	0.3	2
383	Validation of the Risk Score of the Mortality and Lower Limb Loss Considering Ambulatory Status after Surgical Revascularization in Maintaining Patients with Dialysis. Annals of Vascular Diseases, 2017, 10, 192-196.	0.5	2
384	A Large-scale Survey of Health Check-up Visitors in the West-Central Area of Shizuoka Prefecture Regarding the Frequency of Walking in Forested Areas Journal of the Japanese Forest Society, 2010, 92, 110-114.	0.2	2
385	p53 labeling index in cholangioscopic biopsies is useful for determining spread of bile duct carcinomas. Gastrointestinal Endoscopy, 2002, 56, 688-695.	1.0	2
386	<editors' choice=""> Renewed Japanese spirometric reference variables and risk stratification for postoperative outcomes in COPD patients with resected lung cancer. Nagoya Journal of Medical Science, 2019, 81, 427-438.</editors'>	0.3	2
387	Tooth brushing, tooth loss, and risk of upper aerodigestive tract cancer: a cohort study of Japanese dentisits. Nagoya Journal of Medical Science, 2021, 83, 331-341.	0.3	2
388	Plasma Angiopoietin-Like Protein 2 Levels and Mortality Risk Among Younger-Old Japanese People: A Population-Based Case–Cohort Study. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2022, 77, 1150-1158.	3.6	2
389	Dose-Dependent Neurologic Abnormalities in Workers Exposed to 1-Bromopropane. Journal of Occupational and Environmental Medicine, 2011, 53, 1095-1098.	1.7	1
390	Large-scale survey of frequency of forest walking and related factors in a Japanese population inhabiting a large city, and comparison of an urban area and a rural area. Journal of Forest Research, 2013, 18, 454-461.	1.4	1
391	Associations Among Epoetin Therapy, Inflammation, Nutritional Status, and Mortality in Patients on Hemodialysis., 2014, 24, 322-329.		1
392	Institutional factors influencing regional differences in the 1-year survival of dialysis patients. Hemodialysis International, 2015, 19, S5-S10.	0.9	1
393	Association between alcohol intake pattern and metabolic syndrome components and simulated change by alcohol intake reduction: A cross-sectional study from the Japan Multi-Institutional Collaborative Cohort Study. Alcohol, 2020, 89, 129-138.	1.7	1
394	Prediction of 11 -year incidence of psychophysically dependent status or death among community-dwelling younger elderlies: from an age-specified community-based cohort study (the) Tj ETQq0 0	0 rgB1140ver	loak 10 Tf 50
395	Preliminary analysis of nutritional factors in breast cancer. Medical Journal of Indonesia, 0, , 187.	0.5	1
396	Modification of the effect of smoking on cholesterol in Japanese carriers of a PTPN11 polymorphism. Molecular Medicine Reports, 0 , , .	2.4	1

#	Article	IF	Citations
397	No association between MTHFR C677T and serum uric acid levels among Japanese with ABCG2 126QQ and SLC22A12 258WW. Nagoya Journal of Medical Science, 2013, 75, 93-100.	0.3	1
398	Food group intakes and all-cause mortality among a young older Japanese population of the same age: the New Integrated Suburban Seniority Investigation Project. Nagoya Journal of Medical Science, 2021, 83, 169-182.	0.3	1
399	Genome-wide association study of serum prostate-specific antigen levels based on 1000 Genomes imputed data in Japanese: the Japan Multi-Institutional Collaborative Cohort Study. Nagoya Journal of Medical Science, 2021, 83, 183-194.	0.3	1
400	A genome-wide association study on adherence to low-carbohydrate diets in Japanese. European Journal of Clinical Nutrition, 2022, , .	2.9	1
401	Changes in behavioral activities and transition of depressive symptoms among youngerâ€old communityâ€dwelling adults during 6Âyears: An ageâ€specific prospective cohort study. International Journal of Geriatric Psychiatry, 2022, 37, .	2.7	1
402	Response to the Letter to the Editor: Metallothionein MT2A A-5G Polymorphism and the Risk for Chronic Kidney Disease and Diabetes. Toxicological Sciences, 2016, 154, kfw205.	3.1	0
403	OP0048 \hat{a} \in GENOME-WIDE META-ANALYSIS REVEALED MULTIPLE NOVEL LOCI ASSOCIATED WITH SERUM URIC ACIDLEVELS IN JAPANESE. , 2019, , .		0
404	OP0047â€A GENOME-WIDE ASSOCIATION STUDY IDENTIFIED NOVEL LOCI ASSOCIATED WITH THE PROGRESSION FROM ASYMPTOMATIC HYPERURICEMIA TO GOUT. , 2019, , .		0
405	Population-Based Impact of Smoking, Drinking, and Genetic Factors on HDL-Cholesterol Levels in J-MICC Study Participants. Journal of Epidemiology, 2021, , .	2.4	0
406	Association between frequency of snacking and allâ€cause mortality among communityâ€dwelling youngâ€old adults: An ageâ€specific prospective cohort study. Geriatrics and Gerontology International, 2021, 21, 697-704.	1.5	0
407	Insulin-like growth factor 2 and incidence of liver cancer in a nested case-control study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, cebp.EPI-21-0481-E.2021.	2.5	0
408	516Association of dietary diversity with all-cause mortality by body mass index in Japanese older adults. International Journal of Epidemiology, 2021, 50, .	1.9	0
409	Response to Dr Shikata's letter: â€~Secondhand smoke exposure and risk of lung cancer in Japan: a systematic review and meta-analysis of epidemiologic studies'. Japanese Journal of Clinical Oncology, 2021, 51, 661-661.	1.3	O
410	Identification of risk factors for ipsilateral lower limb loss after minor amputation. The Journal of Japanese Society of Limb Salvage and Podiatric Medicine, 2015, 7, 168-172.	0.0	0
411	Associations of Dietary Salt and Its Sources with Hemoglobin A1c in Patients with Type 2 Diabetes Not Taking Anti-Diabetic Medications: Analysis Based on 6-Month Intervention with a Moderate Low-Carbohydrate Diet. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2021, Volume 14. 4569-4578.	2.4	O