Leonardo Furi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11536771/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Methylation Warfare: Interaction of Pneumococcal Bacteriophages with Their Host. Journal of Bacteriology, 2019, 201, .	2.2	22
2	Dissemination of Novel Antimicrobial Resistance Mechanisms through the Insertion Sequence Mediated Spread of Metabolic Genes. Frontiers in Microbiology, 2016, 7, 1008.	3.5	40
3	Significant Differences Characterise the Correlation Coefficients between Biocide and Antibiotic Susceptibility Profiles in Staphylococcus aureus. Current Pharmaceutical Design, 2015, 21, 2054-2057.	1.9	35
4	Mutations upstream of fabl in triclosan resistant Staphylococcus aureus strains are associated with elevated fabl gene expression. BMC Genomics, 2015, 16, 345.	2.8	17
5	Novel insight into antimicrobial resistance and sensitivity phenotypes associated to qac and norA genotypes in Staphylococcus aureus. Microbiological Research, 2015, 170, 184-194.	5.3	27
6	The Role of Host and Microbial Factors in the Pathogenesis of Pneumococcal Bacteraemia Arising from a Single Bacterial Cell Bottleneck. PLoS Pathogens, 2014, 10, e1004026.	4.7	66
7	ls adsorption an artifact in experimentation with Triclosan?. Desalination and Water Treatment, 2014, 52, 7101-7107.	1.0	7
8	A random six-phase switch regulates pneumococcal virulence via global epigenetic changes. Nature Communications, 2014, 5, 5055.	12.8	264
9	Evaluation of Reduced Susceptibility to Quaternary Ammonium Compounds and Bisbiguanides in Clinical Isolates and Laboratory-Generated Mutants of Staphylococcus aureus. Antimicrobial Agents and Chemotherapy, 2013, 57, 3488-3497.	3.2	102
10	Lack of Evidence for Reduced Fitness of Clinical Staphylococcus aureus Isolates with Reduced Susceptibility to Triclosan. Antimicrobial Agents and Chemotherapy, 2012, 56, 6068-6069.	3.2	8
11	A novel resistance mechanism to triclosan that suggests horizontal gene transfer and demonstrates a potential selective pressure for reduced biocide susceptibility in clinical strains of Staphylococcus aureus. International Journal of Antimicrobial Agents, 2012, 40, 210-220.	2.5	92