

Susann Steudte-Schmiedgen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11574705/publications.pdf>

Version: 2024-02-01

27
papers

1,373
citations

623734

14
h-index

580821

25
g-index

28
all docs

28
docs citations

28
times ranked

1934
citing authors

#	ARTICLE	IF	CITATIONS
1	Stress-related and basic determinants of hair cortisol in humans: A meta-analysis. <i>Psychoneuroendocrinology</i> , 2017, 77, 261-274.	2.7	556
2	An integrative model linking traumatization, cortisol dysregulation and posttraumatic stress disorder: Insight from recent hair cortisol findings. <i>Neuroscience and Biobehavioral Reviews</i> , 2016, 69, 124-135.	6.1	127
3	Hair cortisol concentrations and cortisol stress reactivity predict PTSD symptom increase after trauma exposure during military deployment. <i>Psychoneuroendocrinology</i> , 2015, 59, 123-133.	2.7	119
4	The Cortisol Paradox of Trauma-Related Disorders: Lower Phasic Responses but Higher Tonic Levels of Cortisol Are Associated with Sexual Abuse in Childhood. <i>PLoS ONE</i> , 2015, 10, e0136921.	2.5	74
5	Hair cortisol concentrations and cortisol stress reactivity in generalized anxiety disorder, major depression and their comorbidity. <i>Journal of Psychiatric Research</i> , 2017, 84, 184-190.	3.1	71
6	Glucocorticoid receptor gene methylation moderates the association of childhood trauma and cortisol stress reactivity. <i>Psychoneuroendocrinology</i> , 2018, 90, 68-75.	2.7	66
7	Sweat-inducing physiological challenges do not result in acute changes in hair cortisol concentrations. <i>Psychoneuroendocrinology</i> , 2015, 53, 108-116.	2.7	53
8	Social Support and Optimism as Protective Factors for Mental Health among 7765 Healthcare Workers in Germany during the COVID-19 Pandemic: Results of the VOICE Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 3827.	2.6	52
9	The Impact of Parental Role Distributions, Work Participation, and Stress Factors on Family Health-Related Outcomes: Study Protocol of the Prospective Multi-Method Cohort "Dresden Study on Parenting, Work, and Mental Health" (DREAM). <i>Frontiers in Psychology</i> , 2019, 10, 1273.	2.1	32
10	Sense of coherence, social support and religiosity as resources for medical personnel during the COVID-19 pandemic: A web-based survey among 4324 health care workers within the German Network University Medicine. <i>PLoS ONE</i> , 2021, 16, e0255211.	2.5	32
11	The Effects of Modifying Dysfunctional Appraisals in Posttraumatic Stress Disorder Using a Form of Cognitive Bias Modification: Results of a Randomized Controlled Trial in an Inpatient Setting. <i>Psychotherapy and Psychosomatics</i> , 2021, 90, 386-402.	8.8	28
12	The impact of the COVID-19 pandemic on the mental health of medical staff considering the interplay of pandemic burden and psychosocial resources – A rapid systematic review. <i>PLoS ONE</i> , 2022, 17, e0264290.	2.5	28
13	Trauma exposure is associated with increased context-dependent adjustments of cognitive control in patients with posttraumatic stress disorder and healthy controls. <i>Cognitive, Affective and Behavioral Neuroscience</i> , 2014, 14, 1310-1319.	2.0	26
14	The utility of hair cortisol concentrations in the prediction of PTSD symptoms following traumatic physical injury. <i>Social Science and Medicine</i> , 2017, 175, 228-234.	3.8	20
15	Correlates and Predictors of PTSD Symptoms Among Healthcare Workers During the COVID-19 Pandemic: Results of the egePan-VOICE Study. <i>Frontiers in Psychiatry</i> , 2021, 12, 686667.	2.6	16
16	Biological stress indicators as risk markers for increased alcohol use following traumatic experiences. <i>Addiction Biology</i> , 2018, 23, 281-290.	2.6	12
17	Investigating d-cycloserine as a potential pharmacological enhancer of an emotional bias learning procedure. <i>Journal of Psychopharmacology</i> , 2018, 32, 569-577.	4.0	11
18	Ecological momentary assessment in posttraumatic stress disorder and coping. An eHealth study protocol. <i>HÅrre Utbildning</i> , 2019, 10, 1654064.	3.0	9

#	ARTICLE	IF	CITATIONS
19	Hair cortisol predicts avoidance behavior and depressiveness after first-time and single-event trauma exposure in motor vehicle crash victims. <i>Stress</i> , 2020, 23, 567-576.	1.8	8
20	Augmenting inpatient treatment for post-traumatic stress disorder with a computerised cognitive bias modification procedure targeting appraisals (CBM-App): protocol for a randomised controlled trial. <i>BMJ Open</i> , 2018, 8, e019964.	1.9	7
21	Hydrocortisone as an adjunct to brief cognitive-behavioural therapy for specific fear: Endocrine and cognitive biomarkers as predictors of symptom improvement. <i>Journal of Psychopharmacology</i> , 2021, 35, 641-651.	4.0	5
22	Reply to: Linking Hair Cortisol Levels to Phenotypic Heterogeneity of Posttraumatic Stress Symptoms in Highly Traumatized Chinese Women. <i>Biological Psychiatry</i> , 2015, 77, e23-e24.	1.3	3
23	Cognitive functioning in posttraumatic stress disorder before and after cognitive-behavioral therapy. <i>Journal of Anxiety Disorders</i> , 2020, 74, 102265.	3.2	3
24	Human perceptual learning is delayed by the N-methyl-D-aspartate receptor partial agonist D-cycloserine. <i>Journal of Psychopharmacology</i> , 2021, 35, 253-264.	4.0	2
25	Lifetime trauma history and cognitive functioning in major depression and their role for cognitive-behavioral therapy outcome. <i>Clinical Psychology in Europe</i> , 2021, 3, .	1.1	1
26	The predictive role of hair cortisol concentrations for treatment outcome in PTSD inpatients. <i>Psychoneuroendocrinology</i> , 2021, 131, 105326.	2.7	1
27	Reply to the commentary by Parrot and Downey (2017). <i>Psychoneuroendocrinology</i> , 2017, 81, 160.	2.7	0