

Yuqi He

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11590506/publications.pdf>

Version: 2024-02-01

13
papers

409
citations

933447

10
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

748
citing authors

#	ARTICLE	IF	CITATIONS
1	Exploring <sc>microRNAs</sc> in diabetic chronic cutaneous ulcers: Regulatory mechanisms and therapeutic potential. <i>British Journal of Pharmacology</i> , 2020, 177, 4077-4095.	5.4	15
2	Gypenosides Altered Hepatic Bile Acids Homeostasis in Mice Treated with High Fat Diet. <i>Evidence-based Complementary and Alternative Medicine</i> , 2018, 2018, 1-10.	1.2	15
3	All-trans retinoic acid regulates hepatic bile acid homeostasis. <i>Biochemical Pharmacology</i> , 2014, 91, 483-489.	4.4	39
4	Metabolomic and Genomic Evidence for Compromised Bile Acid Homeostasis by Senecionine, a Hepatotoxic Pyrrolizidine Alkaloid. <i>Chemical Research in Toxicology</i> , 2014, 27, 775-786.	3.3	39
5	Biological functional annotation of retinoic acid alpha and beta in mouse liver based on genome-wide binding. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 307, G205-G218.	3.4	5
6	Genome-Wide Binding and Transcriptome Analysis of Human Farnesoid X Receptor in Primary Human Hepatocytes. <i>PLoS ONE</i> , 2014, 9, e105930.	2.5	50
7	The role of retinoic acid in hepatic lipid homeostasis defined by genomic binding and transcriptome profiling. <i>BMC Genomics</i> , 2013, 14, 575.	2.8	57
8	Genome-wide binding and transcriptome analysis of human farnesoid X receptor in the liver. <i>FASEB Journal</i> , 2013, 27, 663.2.	0.5	0
9	Function Annotation of Hepatic Retinoid x Receptor ± Based on Genome-Wide DNA Binding and Transcriptome Profiling. <i>PLoS ONE</i> , 2012, 7, e50013.	2.5	10
10	The comparative pharmacokinetics of two pyrrolizidine alkaloids, senecionine and adonifoline, and their main metabolites in rats after intravenous and oral administration by UPLC/ESIMS. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 401, 275-287.	3.7	34
11	Simultaneous determination of senecionine, adonifoline and their metabolites in rat serum by UPLC-ESIMS and its application in pharmacokinetic studies. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009, 50, 1070-1074.	2.8	16
12	Identification of metabolites of adonifoline, a hepatotoxic pyrrolizidine alkaloid, by liquid chromatography/tandem and high-resolution mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2009, 23, 3907-3916.	1.5	25
13	Bile Acids Metabonomic Study on the CCl ₄ - and ±-Naphthylisothiocyanate-Induced Animal Models: Quantitative Analysis of 22 Bile Acids by Ultrapformance Liquid Chromatography~Mass Spectrometry. <i>Chemical Research in Toxicology</i> , 2008, 21, 2280-2288.	3.3	104