

Xiaolong Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11596424/publications.pdf>

Version: 2024-02-01

16
papers

354
citations

1040056

9
h-index

1058476

14
g-index

17
all docs

17
docs citations

17
times ranked

522
citing authors

#	ARTICLE	IF	CITATIONS
1	EGFR suppresses p53 function by promoting p53 binding to DNA-PKcs: a noncanonical regulatory axis between EGFR and wild-type p53 in glioblastoma. <i>Neuro-Oncology</i> , 2022, 24, 1712-1725.	1.2	8
2	PARP-mediated PARylation of MGMT is critical to promote repair of temozolomide-induced O6-methylguanine DNA damage in glioblastoma. <i>Neuro-Oncology</i> , 2021, 23, 920-931.	1.2	58
3	The promise of DNA damage response inhibitors for the treatment of glioblastoma. <i>Neuro-Oncology Advances</i> , 2021, 3, vdab015.	0.7	16
4	EGFR Amplification Induces Increased DNA Damage Response and Renders Selective Sensitivity to Talazoparib (PARP Inhibitor) in Glioblastoma. <i>Clinical Cancer Research</i> , 2020, 26, 1395-1407.	7.0	26
5	Tie2-FGFR1 Interaction Induces Adaptive PI3K Inhibitor Resistance by Upregulating Aurora A/PLK1/CDK1 Signaling in Glioblastoma. <i>Cancer Research</i> , 2019, 79, 5088-5101.	0.9	17
6	Wild-type defined gamma-secretase inhibitor sensitivity and synergistic activity with doxorubicin in GSCs. <i>American Journal of Cancer Research</i> , 2019, 9, 1734-1745.	1.4	3
7	BRCA1 identified as a modulator of temozolomide resistance in P53 wild-type GBM using a high-throughput shRNA-based synthetic lethality screening. <i>American Journal of Cancer Research</i> , 2019, 9, 2428-2441.	1.4	1
8	DDIS-03. EGFR AMPLIFICATION INDUCED INCREASED DNA DAMAGE RESPONSE AND PREDICTED SELECTIVE SENSITIVITY TO TALAZOPARIB (PARP INHIBITOR) IN GLIOBLASTOMA STEM-LIKE CELLS. <i>Neuro-Oncology</i> , 2018, 20, vi69-vi69.	1.2	0
9	EXTH-11. GLIOBLASTOMA STEM CELL GROWTH DEPENDENCE ON NUTRIENTS: MORE THAN BASAL METABOLIC ACTIVITIES. <i>Neuro-Oncology</i> , 2018, 20, vi87-vi87.	1.2	0
10	Molecular Aberrance in Papillary Thyroid Microcarcinoma Bearing High Aggressiveness: Identifying a Tibetan Mastiff Dog-From Puppies. <i>Journal of Cellular Biochemistry</i> , 2016, 117, 1491-1496.	2.6	10
11	Low expression of sodium iodide symporter expression in aggressive variants of papillary thyroid carcinoma. <i>International Journal of Clinical Oncology</i> , 2014, 19, 800-804.	2.2	9
12	Hemithyroidectomy increases the risk of disease recurrence in patients with ipsilateral multifocal papillary thyroid carcinoma. <i>Oncology Letters</i> , 2013, 5, 1412-1416.	1.8	14
13	Achieving cell penetration with distance-matching cysteine cross-linkers: a facile route to cell-permeable peptide dual inhibitors of Mdm2/Mdmx. <i>Chemical Communications</i> , 2011, 47, 9396.	4.1	79
14	Conjugation of spermine enhances cellular uptake of the stapled peptide-based inhibitors of p53-Mdm2 interaction. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011, 21, 7412-7415.	2.2	14
15	Synthesis of cell-permeable stapled peptide dual inhibitors of the p53-Mdm2/Mdmx interactions via photoinduced cycloaddition. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011, 21, 1472-1475.	2.2	97
16	Multi-analysis with mathematic model of 3125 non-thyrogeous masses of the neck. <i>Chinese-German Journal of Clinical Oncology</i> , 2008, 7, 319-325.	0.1	0