

Andreas G Wibmer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11625837/publications.pdf>

Version: 2024-02-01

13
papers

751
citations

1163117

8
h-index

1199594

12
g-index

14
all docs

14
docs citations

14
times ranked

1330
citing authors

#	ARTICLE	IF	CITATIONS
1	Brown adipose tissue is associated with cardiometabolic health. <i>Nature Medicine</i> , 2021, 27, 58-65.	30.7	332
2	Background, current role, and potential applications of radiogenomics. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 604-620.	3.4	137
3	Molecular Imaging of Prostate Cancer. <i>Radiographics</i> , 2016, 36, 142-159.	3.3	83
4	Brown adipose tissue is associated with healthier body fat distribution and metabolic benefits independent of regional adiposity. <i>Cell Reports Medicine</i> , 2021, 2, 100332.	6.5	51
5	Comparison of Magnetic Resonance Imaging-stratified Clinical Pathways and Systematic Transrectal Ultrasound-guided Biopsy Pathway for the Detection of Clinically Significant Prostate Cancer: A Systematic Review and Meta-analysis of Randomized Controlled Trials. <i>European Urology Oncology</i> , 2019, 2, 605-616.	5.4	30
6	Trends in oncologic hybrid imaging. <i>European Journal of Hybrid Imaging</i> , 2018, 2, 1.	1.5	25
7	Local Extent of Prostate Cancer at MRI versus Prostatectomy Histopathology: Associations with Long-term Oncologic Outcomes. <i>Radiology</i> , 2022, 302, 595-602.	7.3	12
8	International Multi-Site Initiative to Develop an MRI-Inclusive Nomogram for Side-Specific Prediction of Extraprostatic Extension of Prostate Cancer. <i>Cancers</i> , 2021, 13, 2627.	3.7	11
9	MRI-detectability of clinically significant prostate cancer relates to oncologic outcomes after prostatectomy. <i>Clinical Genitourinary Cancer</i> , 2022, , .	1.9	6
10	Endovascular Treatment of Delayed Rupture Following Prior Abdominal Aortic Aneurysm Repair Achieves Better Survival Rates. <i>Journal of Endovascular Therapy</i> , 2013, 20, 609-618.	1.5	4
11	Endovascular conversion into aorto-uniiliac configuration of acute failed endovascular aneurysm repair is associated with better one-year survival rates compared to open conversion. <i>Journal of the Royal College of Surgeons of Edinburgh</i> , 2017, 15, 131-138.	1.8	2
12	Secondary modification into aortouniiliac configuration to salvage failed endovascular aneurysm repair is safe and effective but not associated with higher intervention rates during long-term follow-up. <i>American Journal of Surgery</i> , 2014, 208, 435-443.	1.8	0
13	Rates of adverse events and correction procedures after elective versus emergent aortouniiliac endografting during mid-term follow-up: A prospective cohort study. <i>International Journal of Surgery</i> , 2015, 18, 104-109.	2.7	0