

# Maurizio Brigotti

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1163140/publications.pdf>

Version: 2024-02-01

63  
papers

1,545  
citations

257450

24  
h-index

330143

37  
g-index

64  
all docs

64  
docs citations

64  
times ranked

1368  
citing authors

#	ARTICLE	IF	CITATIONS
1	Damage to nuclear DNA induced by Shiga toxin 1 and ricin in human endothelial cells1. <i>FASEB Journal</i> , 2002, 16, 365-372.	0.5	133
2	Novel Dyskerin-Mediated Mechanism of p53 Inactivation through Defective mRNA Translation. <i>Cancer Research</i> , 2010, 70, 4767-4777.	0.9	95
3	Identification of TLR4 as the Receptor That Recognizes Shiga Toxins in Human Neutrophils. <i>Journal of Immunology</i> , 2013, 191, 4748-4758.	0.8	76
4	Early Volume Expansion and Outcomes of Hemolytic Uremic Syndrome. <i>Pediatrics</i> , 2016, 137, .	2.1	74
5	Clinical Relevance of Shiga Toxin Concentrations in the Blood of Patients With Hemolytic Uremic Syndrome. <i>Pediatric Infectious Disease Journal</i> , 2011, 30, 486-490.	2.0	67
6	Effects of osmolarity, ions and compatible osmolytes on cell-free protein synthesis. <i>Biochemical Journal</i> , 2003, 369, 369-374.	3.7	59
7	Creatine as a compatible osmolyte in muscle cells exposed to hypertonic stress. <i>Journal of Physiology</i> , 2006, 576, 391-401.	2.9	57
8	Human ribosomes from cells with reduced dyskerin levels are intrinsically altered in translation. <i>FASEB Journal</i> , 2015, 29, 3472-3482.	0.5	57
9	Shiga Toxins Present in the Gut and in the Polymorphonuclear Leukocytes Circulating in the Blood of Children with Hemolytic-Uremic Syndrome. <i>Journal of Clinical Microbiology</i> , 2006, 44, 313-317.	3.9	52
10	Flow cytometry detection of Shiga toxins in the blood from children with hemolytic uremic syndrome. <i>Cytometry</i> , 2004, 61B, 40-44.	1.8	50
11	Interactions between Shiga toxins and human polymorphonuclear leukocytes. <i>Journal of Leukocyte Biology</i> , 2008, 84, 1019-1027.	3.3	50
12	Dyskerin depletion increases VEGF mRNA internal ribosome entry site-mediated translation. <i>Nucleic Acids Research</i> , 2013, 41, 8308-8318.	14.5	50
13	Ribosome-inactivating proteins depurinate poly(ADP-ribosyl)ated poly(ADP-ribose) polymerase and have transforming activity for 3T3 fibroblasts. <i>FEBS Letters</i> , 2003, 538, 178-182.	2.8	44
14	Endothelial damage induced by Shiga toxins delivered by neutrophils during transmigration. <i>Journal of Leukocyte Biology</i> , 2010, 88, 201-210.	3.3	40
15	Shiga-like toxin I is a polynucleotide:adenosine glycosidase. <i>Molecular Microbiology</i> , 1998, 29, 661-662.	2.5	37
16	Attenuated Expression of 70-kDa Heat Shock Protein in WI-38 Human Fibroblasts during Aging in Vitro. <i>Experimental Cell Research</i> , 1999, 252, 20-32.	2.6	35
17	Shiga toxin 1 and ricin inhibit the repair of H2O2-induced DNA single strand breaks in cultured mammalian cells. <i>DNA Repair</i> , 2005, 4, 271-277.	2.8	34
18	Molecular Damage and Induction of Proinflammatory Cytokines in Human Endothelial Cells Exposed to Shiga Toxin 1, Shiga Toxin 2, and Î±-Sarcin. <i>Infection and Immunity</i> , 2007, 75, 2201-2207.	2.2	34

#	ARTICLE	IF	CITATIONS
19	Shiga toxin 1 and ricin A chain bind to human polymorphonuclear leucocytes through a common receptor. <i>Biochemical Journal</i> , 2010, 432, 173-180.	3.7	32
20	Differential requirement of ATP and extra-ribosomal proteins for ribosome inactivation by eight RNA N-glycosidases. <i>Biochemical and Biophysical Research Communications</i> , 1992, 182, 579-582.	2.1	31
21	The RNA-N-glycosidase activity of Shiga-like toxin I: Kinetic parameters of the native and activated toxin. <i>Toxicon</i> , 1997, 35, 1431-1437.	1.6	30
22	A reconstituted cell-free assay for the evaluation of the intrinsic activity of purified human ribosomes. <i>Nature Protocols</i> , 2016, 11, 1309-1325.	12.0	29
23	The Interactions of Human Neutrophils with Shiga Toxins and Related Plant Toxins: Danger or Safety?. <i>Toxins</i> , 2012, 4, 157-190.	3.4	28
24	Shiga toxin 1: damage to DNA in vitro. <i>Toxicon</i> , 2001, 39, 341-348.	1.6	26
25	Galloflavin prevents the binding of lactate dehydrogenase A to single stranded DNA and inhibits RNA synthesis in cultured cells. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 466-469.	2.1	25
26	Serum Shiga toxin 2 values in patients during acute phase of diarrhoea-associated haemolytic uraemic syndrome. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2015, 104, e564-8.	1.5	19
27	Particulate Shiga Toxin 2 in Blood is Associated to the Development of Hemolytic Uremic Syndrome in Children. <i>Thrombosis and Haemostasis</i> , 2020, 120, 107-120.	3.4	16
28	Bloody Diarrhea and Shiga Toxin-producing Escherichia coli Hemolytic Uremic Syndrome in Children: Data from the ItalKid-HUS Network. <i>Journal of Pediatrics</i> , 2021, 237, 34-40.e1.	1.8	16
29	Differential up-regulation by tRNAs of ribosome-inactivating proteins. <i>FEBS Letters</i> , 1995, 373, 115-118.	2.8	15
30	Change in Conformation with Reduction of $\alpha$ -Helix Content Causes Loss of Neutrophil Binding Activity in Fully Cytotoxic Shiga Toxin 1. <i>Journal of Biological Chemistry</i> , 2011, 286, 34514-34521.	3.4	14
31	An Improved Method for the Sensitive Detection of Shiga Toxin 2 in Human Serum. <i>Toxins</i> , 2018, 10, 59.	3.4	13
32	The structure of the Shiga toxin 2a subunit dictates the interactions of the toxin with blood components. <i>Cellular Microbiology</i> , 2019, 21, e13000.	2.1	13
33	Shiga toxin 1 acting on DNA in vitro is a heat-stable enzyme not requiring proteolytic activation. <i>Biochimie</i> , 2004, 86, 305-309.	2.6	12
34	Human monocytes stimulated by Shiga toxin 1a via globotriaosylceramide release proinflammatory molecules associated with hemolytic uremic syndrome. <i>International Journal of Medical Microbiology</i> , 2018, 308, 940-946.	3.6	12
35	Is Shiga toxin 1 protective for the development of Shiga toxin 2-related hemolytic uremic syndrome in children? Data from the ItalKid-HUS Network. <i>Pediatric Nephrology</i> , 2020, 35, 1997-2001.	1.7	12
36	Hemidesmus indicus induces apoptosis via proteasome inhibition and generation of reactive oxygen species. <i>Scientific Reports</i> , 2019, 9, 7199.	3.3	11

#	ARTICLE	IF	CITATIONS
37	Uncompetitive inhibition by adenine of the RNA-N-glycosidase activity of ribosome-inactivating proteins. <i>BBA - Proteins and Proteomics</i> , 1998, 1384, 277-284.	2.1	10
38	A Rapid and Sensitive Method to Measure the Functional Activity of Shiga Toxins in Human Serum. <i>Toxins</i> , 2015, 7, 4564-4576.	3.4	10
39	Identification of the tRNAs which up-regulate agrostin, barley RIP and PAP-S, three ribosome-inactivating proteins of plant origin. <i>FEBS Letters</i> , 1998, 431, 259-262.	2.8	9
40	Soluble Toll-Like Receptor 4 Impairs the Interaction of Shiga Toxin 2a with Human Serum Amyloid P Component. <i>Toxins</i> , 2018, 10, 379.	3.4	9
41	Cofactor requirement of ribosome-inactivating proteins from plants. <i>Journal of Experimental Botany</i> , 1997, 48, 1519-1523.	4.8	8
42	5â€™-Untranslated region of heat shock protein 70 mRNA drives translation under hypertonic conditions. <i>Biochemical and Biophysical Research Communications</i> , 2013, 431, 321-325.	2.1	8
43	Shiga Toxin 1, as DNA Repair Inhibitor, Synergistically Potentiates the Activity of the Anticancer Drug, Mafofamide, on Raji Cells. <i>Toxins</i> , 2013, 5, 431-444.	3.4	8
44	RiboAbacus: a model trained on polyribosome images predicts ribosome density and translational efficiency from mammalian transcriptomes. <i>Nucleic Acids Research</i> , 2015, 43, e153-e153.	14.5	8
45	The Antibiotic Polymyxin B Impairs the Interactions between Shiga Toxins and Human Neutrophils. <i>Journal of Immunology</i> , 2016, 196, 1177-1185.	0.8	8
46	The inhibition of lactate dehydrogenase A hinders the transcription of histone 2B gene independently from the block of aerobic glycolysis. <i>Biochemical and Biophysical Research Communications</i> , 2017, 485, 742-745.	2.1	8
47	Inhibition by suramin of protein synthesis in vitro. Ribosomes as the target of the drug. <i>Biochimie</i> , 2006, 88, 497-503.	2.6	7
48	Extracellular Vesicles and Renal Endothelial Cells. <i>American Journal of Pathology</i> , 2021, 191, 795-804.	3.8	7
49	Interaction of diphtheria toxin fragment A and of elongation factor 2 with Cibacron blue. <i>Bioscience Reports</i> , 1987, 7, 737-743.	2.4	6
50	Alpha-sarcin impairs the N-glycosidase activity of ricin on ribosomes. <i>Biochemical and Biophysical Research Communications</i> , 1989, 160, 857-861.	2.1	6
51	Nucleotides U28-A42 and A37 in unmodified yeast tRNA <sup>Trp</sup> as negative identity elements for bovine tryptophanyl-tRNA synthetase. <i>FEBS Letters</i> , 2001, 492, 238-241.	2.8	6
52	Identity elements in bovine tRNA <sup>Trp</sup> required for the specific stimulation of gelonin, a plant ribosome-inactivating protein. <i>Rna</i> , 1999, 5, 1357-1363.	3.5	5
53	A survey of adenine and 4-aminopyrazolo[3,4-d]pyrimidine (4-APP) as inhibitors of ribosome-inactivating proteins (RIPs). <i>Life Sciences</i> , 2000, 68, 331-336.	4.3	5
54	Partial purification of two proteins which sensitize ribosomes to gelonin: Sensitization is not linked to phosphorylation of ribosomal proteins. <i>Toxicon</i> , 1993, 31, 989-996.	1.6	4

#	ARTICLE	IF	CITATIONS
55	Elongation factor 2 from <i>Artemia salina</i> embryos and its affinity for ribosomes. <i>FEBS Journal</i> , 1991, 200, 13-18.	0.2	3
56	Cap-independent protein synthesis is enhanced by betaine under hypertonic conditions. <i>Biochemical and Biophysical Research Communications</i> , 2017, 483, 936-940.	2.1	3
57	Shiga Toxin 2 Triggers C3a-Dependent Glomerular and Tubular Injury through Mitochondrial Dysfunction in Hemolytic Uremic Syndrome. <i>Cells</i> , 2022, 11, 1755.	4.1	3
58	Deuterium Incorporation Protects Cells from Oxidative Damage. <i>Oxidative Medicine and Cellular Longevity</i> , 2019, 2019, 1-13.	4.0	2
59	Method for the Detection of the Cleaved Form of Shiga Toxin 2a Added to Normal Human Serum. <i>Toxins</i> , 2021, 13, 94.	3.4	2
60	tRNATrp as cofactor of gelonin, a ribosome-inactivating protein with RNA-N-glycosidase activity features required for the cofactor activity. <i>IUBMB Life</i> , 1996, 40, 181-188.	3.4	1
61	Primer tRNATrp of RSV-transformed or RAV-1-infected cells up-regulates the antiribosomal activity of gelonin. <i>Biochimie</i> , 1998, 80, 575-578.	2.6	1
62	No convergence for references. <i>Nature</i> , 1998, 393, 301-301.	27.8	0
63	Variable biological properties of two different preparations of Shiga toxins yielding new insights into eHUS pathogenesis. <i>Molecular Immunology</i> , 2017, 89, 159.	2.2	0