

Lone Petersen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11635260/publications.pdf>

Version: 2024-02-01

28
papers

2,688
citations

331670

21
h-index

501196

28
g-index

29
all docs

29
docs citations

29
times ranked

2387
citing authors

#	ARTICLE	IF	CITATIONS
1	The Momentum trial: the efficacy of using a smartphone application to promote patient activation and support shared decision making in people with a diagnosis of schizophrenia in outpatient treatment settings: a randomized controlled single-blind trial. <i>BMC Psychiatry</i> , 2019, 19, 185.	2.6	13
2	Recovery-oriented practice: Participant observations of the interactions between patients and health professionals in mental health inpatient settings. <i>International Journal of Mental Health Nursing</i> , 2019, 28, 318-329.	3.8	29
3	Recovery orientation in mental health inpatient settings: Inpatient experiences?. <i>International Journal of Mental Health Nursing</i> , 2018, 27, 1177-1187.	3.8	25
4	Recovery-Oriented Practice in Mental Health Inpatient Settings: A Literature Review. <i>Psychiatric Services</i> , 2016, 67, 596-602.	2.0	61
5	Long-term trajectories of positive and negative symptoms in first episode psychosis: A 10year follow-up study in the OPUS cohort. <i>Schizophrenia Research</i> , 2015, 168, 84-91.	2.0	141
6	From research to practice: how OPUS treatment was accepted and implemented throughout Denmark. <i>Microbial Biotechnology</i> , 2015, 9, 156-162.	1.7	74
7	Predictors of recovery in first episode psychosis: The OPUS cohort at 10year follow-up. <i>Schizophrenia Research</i> , 2013, 150, 163-168.	2.0	144
8	Cost-effectiveness of early intervention in first-episode psychosis: economic evaluation of a randomised controlled trial (the OPUS study). <i>British Journal of Psychiatry</i> , 2013, 202, 35-41.	2.8	106
9	Predictors of recovery from psychosis. <i>Schizophrenia Research</i> , 2011, 125, 257-266.	2.0	143
10	Treatment of first-episode psychosis patient, psychosocial aspects. <i>Drug Discovery Today: Therapeutic Strategies</i> , 2011, 8, 37-41.	0.5	1
11	The effect of five years versus two years of specialised assertive intervention for first episode psychosis - OPUS II: study protocol for a randomized controlled trial. <i>Trials</i> , 2011, 12, 72.	1.6	35
12	The quality of life among first-episode psychotic patients in the opus trial. <i>Schizophrenia Research</i> , 2010, 116, 27-34.	2.0	33
13	Predictors of poor adherence to medication among patients with first-episode schizophrenia-spectrum disorder. <i>Microbial Biotechnology</i> , 2009, 3, 66-74.	1.7	38
14	Course of illness in a sample of 265 patients with first-episode psychosis—Five-year follow-up of the Danish OPUS trial. <i>Schizophrenia Research</i> , 2009, 107, 173-178.	2.0	85
15	Does a detection team shorten duration of untreated psychosis?. <i>Microbial Biotechnology</i> , 2008, 2, 22-26.	1.7	7
16	Effect of integrated treatment on the use of coercive measures in first-episode schizophrenia-spectrum disorder. A randomized clinical trial. <i>International Journal of Law and Psychiatry</i> , 2008, 31, 72-76.	0.9	17
17	Five-Year Follow-up of a Randomized Multicenter Trial of Intensive Early Intervention vs Standard Treatment for Patients With a First Episode of Psychotic Illness. <i>Archives of General Psychiatry</i> , 2008, 65, 762.	12.3	399
18	Predictors of Remission and Recovery in a First-Episode Schizophrenia Spectrum Disorder Sample: 2-Year Follow-up of the OPUS Trial. <i>Canadian Journal of Psychiatry</i> , 2008, 53, 660-670.	1.9	87

#	ARTICLE	IF	CITATIONS
19	Frequency and predictive values of first rank symptoms at baseline among 362 young adult patients with first-episode schizophrenia. <i>Schizophrenia Research</i> , 2007, 97, 60-67.	2.0	19
20	Gender Differences in Young Adults With First-Episode Schizophrenia Spectrum Disorders at Baseline in the Danish OPUS Study. <i>Journal of Nervous and Mental Disease</i> , 2007, 195, 396-405.	1.0	158
21	Substance abuse and first-episode schizophrenia spectrum disorders. The Danish OPUS trial. <i>Microbial Biotechnology</i> , 2007, 1, 88-96.	1.7	43
22	Transition rates from schizotypal disorder to psychotic disorder for first-contact patients included in the OPUS trial. A randomized clinical trial of integrated treatment and standard treatment. <i>Schizophrenia Research</i> , 2006, 83, 29-40.	2.0	154
23	Social network among young adults with first-episode schizophrenia spectrum disorders. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2006, 41, 761-770.	3.1	55
24	Treatment of patients with first-episode psychosis: two-year outcome data from the Danish National Schizophrenia Project. <i>World Psychiatry</i> , 2006, 5, 100-3.	10.4	16
25	A randomised multicentre trial of integrated versus standard treatment for patients with a first episode of psychotic illness. <i>BMJ: British Medical Journal</i> , 2005, 331, 602.	2.3	480
26	Integrated treatment of first-episode psychosis: effect of treatment on family burden. <i>British Journal of Psychiatry</i> , 2005, 187, s85-s90.	2.8	125
27	Improving 1-year outcome in first-episode psychosis. <i>British Journal of Psychiatry</i> , 2005, 187, s98-s103.	2.8	131
28	The Danish National Schizophrenia Project: prospective, comparative longitudinal treatment study of first-episode psychosis. <i>British Journal of Psychiatry</i> , 2005, 186, 394-399.	2.8	69