Monsalud del Olmo-Iruela

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1167632/publications.pdf

Version: 2024-02-01

186265 58 1,978 28 citations h-index papers

g-index 61 61 61 2144 docs citations times ranked citing authors all docs

254184

43

#	Article	IF	Citations
1	Multiclass cyanotoxin analysis in reservoir waters: Tandem solid-phase extraction followed by zwitterionic hydrophilic interaction liquid chromatography-mass spectrometry. Talanta, 2022, 237, 122929.	5.5	15
2	A novel approach based on capillary liquid chromatography for the simultaneous determination of neonicotinoid residues in cereal samples. Microchemical Journal, 2021, 161, 105756.	4.5	9
3	Micellar electrokinetic chromatography as efficient alternative for the multiresidue determination of seven neonicotinoids and 6-chloronicotinic acid in environmental samples. Analytical and Bioanalytical Chemistry, 2020, 412, 6231-6240.	3.7	11
4	Capillary liquid chromatography as an effective method for the determination of seven neonicotinoid residues in honey samples. Journal of Separation Science, 2020, 43, 3847-3855.	2.5	9
5	Aminoâ€functionalized material from a bioâ€template for silver adsorption: process evaluation in batch and fixed bed. Journal of Chemical Technology and Biotechnology, 2019, 94, 590-599.	3.2	10
6	Monitoring of cyanotoxins in water from hypersaline microalgae colonies by ultra high performance liquid chromatography with diode array and tandem mass spectrometry detection following salting-out liquid-liquid extraction. Journal of Chromatography A, 2019, 1608, 460409.	3.7	13
7	Ultra-high performance liquid chromatography with fluorescence detection following salting-out assisted liquid–liquid extraction for the analysis of benzimidazole residues in farm fish samples. Journal of Chromatography A, 2018, 1543, 58-66.	3.7	10
8	Collision cross section (CCS) as a complementary parameter to characterize human and veterinary drugs. Analytica Chimica Acta, 2018, 1043, 52-63.	5.4	43
9	Use of carbon paste electrodes as a novel strategy to study adsorption mechanism of silver ions onto functionalized grapefruit peel. Journal of Electroanalytical Chemistry, 2018, 830-831, 20-26.	3.8	7
10	Green and simple analytical method to determine benzimidazoles in milk samples by using salting-out assisted liquid-liquid extraction and capillary liquid chromatography. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2018, 1091, 46-52.	2.3	24
11	Determination of benzimidazoles in meat samples by capillary zone electrophoresis tandem mass spectrometry following dispersive liquid–liquid microextraction. Journal of Chromatography A, 2017, 1490, 212-219.	3.7	26
12	Validation of a new method based on salting-out assisted liquid-liquid extraction and UHPLC-MS/MS for the determination of betalactam antibiotics in infant dairy products. Talanta, 2017, 167, 493-498.	5.5	33
13	Coupling sweeping-micellar electrokinetic chromatography with tandem mass spectrometry for the therapeutic monitoring of benzimidazoles in animal urine by dilute and shoot. Talanta, 2017, 175, 542-549.	5.5	15
14	Capillary electrochromatography coupled with dispersive liquid-liquid microextraction for the analysis of benzimidazole residues in water samples. Talanta, 2016, 161, 8-14.	5.5	20
15	Use of an ionic liquidâ€based surfactant as pseudostationary phase in the analysis of carbamates by micellar electrokinetic chromatography. Electrophoresis, 2015, 36, 955-961.	2.4	22
16	Vortex-assisted ionic liquid dispersive liquid–liquid microextraction for the determination of sulfonylurea herbicides in wine samples by capillary high-performance liquid chromatography. Food Chemistry, 2015, 170, 348-353.	8.2	70
17	Salting-out assisted liquid–liquid extraction combined with capillary HPLC for the determination of sulfonylurea herbicides in environmental water and banana juice samples. Talanta, 2014, 127, 51-58.	5.5	70
18	Hollowâ€fiber liquidâ€phase microextraction combined with capillary <scp>HPLC</scp> for the selective determination of six sulfonylurea herbicides in environmental waters. Journal of Separation Science, 2013, 36, 3395-3401.	2.5	28

#	Article	IF	CITATIONS
19	Dispersive Liquid–Liquid Microextraction Followed by Capillary High-Performance Liquid Chromatography for the Determination of Six Sulfonylurea Herbicides in Fruit Juices. Food Analytical Methods, 2013, 7, 1465.	2.6	6
20	On-line anion exchange solid-phase extraction coupled to liquid chromatography with fluorescence detection to determine quinolones in water and human urine. Journal of Chromatography A, 2013, 1310, 91-97.	3.7	34
21	Ion-paired extraction of cephalosporins in acetone prior to their analysis by capillary liquid chromatography in environmental water and meat samples. Talanta, 2013, 115, 943-949.	5 . 5	24
22	Evaluation of dispersive liquid–liquid microextraction for the determination of patulin in apple juices using micellar electrokinetic capillary chromatography. Food Control, 2013, 31, 353-358.	5.5	62
23	Advances in the determination of \hat{l}^2 -lactam antibiotics by liquid chromatography. TrAC - Trends in Analytical Chemistry, 2012, 38, 52-66.	11.4	74
24	Convenient solid phase extraction of cephalosporins in milk using a molecularly imprinted polymer. Food Chemistry, 2012, 135, 775-779.	8.2	49
25	Analysis of cephalosporin residues in environmental waters by capillary zone electrophoresis with off-line and on-line preconcentration. Analytical Methods, 2012, 4, 2341.	2.7	20
26	Trace determination of sulfonylurea herbicides in water and grape samples by capillary zone electrophoresis using large volume sample stacking. Analytical and Bioanalytical Chemistry, 2010, 397, 2593-2601.	3.7	44
27	Multiresidue determination of penicillins in environmental waters and chicken muscle samples by means of capillary electrophoresisâ€ŧandem mass spectrometry. Electrophoresis, 2009, 30, 1708-1717.	2.4	33
28	Applications of capillary electrophoresis to the determination of antibiotics in food and environmental samples. Analytical and Bioanalytical Chemistry, 2009, 395, 967-986.	3.7	81
29	Trace determination of $10\hat{l}^2$ -lactam antibiotics in environmental and food samples by capillary liquid chromatography. Journal of Chromatography A, 2009, 1216, 8355-8361.	3.7	58
30	Trace determination of \hat{l}^2 -lactam antibiotics in environmental aqueous samples using off-line and on-line preconcentration in capillary electrophoresis. Journal of Chromatography A, 2008, 1185, 273-280.	3.7	71
31	Largeâ€volume sample stacking for the analysis of seven βâ€lactam antibiotics in milk samples of different origins by CZE. Electrophoresis, 2007, 28, 4082-4090.	2.4	39
32	Large volume sample stacking in capillary zone electrophoresis for the monitoring of the degradation products of metribuzin in environmental samples. Journal of Chromatography A, 2007, 1164, 320-328.	3.7	32
33	Determination of the herbicide metribuzin and its major conversion products in soil by micellar electrokinetic chromatography. Journal of Chromatography A, 2006, 1102, 280-286.	3.7	35
34	Use of solid-phase microextraction followed by on-column silylation for determining chlorinated bisphenol A in human plasma by gas chromatography–mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2005, 817, 167-172.	2.3	47
35	Development of Methodologies for Different Degrees of Resolution of Linear Alkylbenzene Sulfonates in Groundwater and Wastewater by Liquid Chromatography Using Sodium Dodecyl Sulphate. Chromatographia, 2004, 60, .	1.3	2
36	Sensitive determination of carbaryl in vegetal food and natural waters by flow-injection analysis based on the luminol chemiluminescence reaction. Analytica Chimica Acta, 2004, 524, 161-166.	5.4	35

#	Article	IF	CITATIONS
37	Gas chromatographic–mass spectrometric method for the determination of bisphenol A and its chlorinated derivatives in urban wastewater. Water Research, 2003, 37, 735-742.	11.3	84
38	Determination of bisphenol-a and related compounds in human saliva by gas chromatography—mass spectrometry. Chromatographia, 2002, 56, 213-218.	1.3	24
39	Determination of trace amounts of bisphenol F, bisphenol A and their diglycidyl ethers in wastewater by gas chromatography–mass spectrometry. Analytica Chimica Acta, 2001, 431, 31-40.	5 . 4	90
40	Mortars, pigments and binding media of wall paintings in the †Carrera del Darro' in Granada, Spain. Journal of Cultural Heritage, 2000, 1, 19-28.	3.3	21
41	Determination of carbetamide in groundwater by micro liquid-liquid extraction and gas chromatography-mass spectrometry. Chromatographia, 2000, 52, 233-236.	1.3	6
42	Resolution of mixtures of three nonsteroidal anti-inflammatory drugs by fluorescence using partial least squares multivariate calibration with previous wavelength selection by Kohonen artificial neural networks. Talanta, 2000, 52, 1069-1079.	5 . 5	33
43	Determination of bisphenol A (BPA) in the presence of phenol by first-derivative fluorescence following micro liquid–liquid extraction (MLLE). Talanta, 2000, 50, 1141-1148.	5.5	38
44	Kohonen artificial neural networks as a tool for wavelength selection in multicomponent spectrofluorimetric PLS modelling: application to phenol, o-cresol, m-cresol and p-cresol mixtures. TrAC - Trends in Analytical Chemistry, 1999, 18, 93-98.	11.4	45
45	Trace determination of phenol, bisphenol A and bisphenol A diglycidyl ether in mixtures by excitation fluorescence following micro liquid–liquid extraction using partial least squares regression. Analyst, The, 1999, 124, 385-390.	3. 5	18
46	Simultaneous determination of naproxen, salicylic acid and acetylsalicylic acid by spectrofluorimetry using partial least-squares (PLS) multivariate calibration. Talanta, 1999, 48, 469-475.	5 . 5	68
47	Use of Transmitted Room-Temperature Phosphorescence to Improve Nalidixic Acid Determination. Applied Spectroscopy, 1998, 52, 101-105.	2.2	6
48	Determination of Bisphenol A in Water by Micro Liquid-Liquid Extraction Followed by Silylation and Gas Chromatography-Mass Spectrometry Analysis. Journal of Chromatographic Science, 1998, 36, 565-570.	1.4	87
49	Determination of trace amounts of carbaryl in water by solid-phase laser-induced fluorescence. Talanta, 1997, 44, 443-449.	5. 5	18
50	Determination of bisphenol A (BPA) in water by gas chromatography-mass spectrometry. Analytica Chimica Acta, 1997, 346, 87-92.	5 . 4	107
51	Determination of Synthetic Food Antioxidants in Multicomponent Mixtures Using UV-Visible Spectrophotometry and Partial Least-Squares Calibration. Applied Spectroscopy, 1996, 50, 449-453.	2.2	27
52	Resolution of phenol, o-cresol, m-cresol and p-cresol mixtures by excitation fluorescence using partial least-squares (PLS) multivariate calibration. Analytica Chimica Acta, 1996, 335, 23-33.	5.4	46
53	Laser-Excited Synchronous Fluorescence System for the Analysis of Polycyclic Aromatic Compounds. Polycyclic Aromatic Compounds, 1996, 9, 265-272.	2.6	3
54	Close overlapping discrimination of polycyclic aromatic hydrocarbons by synchronous scanning at variable-angle solid-phase spectrofluorimetry. Analytica Chimica Acta, 1995, 302, 193-200.	5.4	17

#	Article	IF	CITATIONS
55	Determination of polycyclic aromatic hydrocarbon residues in water by synchronous solid-phase spectrofluorimetry. Analyst, The, 1994, 119, 1211-1214.	3.5	28
56	Determination of pyrene and benzo(a)pyrene residues in water by derivative synchronous solid-phase spectrofluorimetry. Mikrochimica Acta, 1993, 112, 55-62.	5.0	7
57	Determination of Ultra-Traces of Anthracene in Water Samples by Solid-Phase Spectrofluorometry. Analytical Sciences, 1993, 9, 117-120.	1.6	10
58	Determination of Benzo(a)pyrene in Water by Synchronous Fluorimetry Following Preconcentration on Sephadex Gels. Analytical Letters, 1993, 26, 2443-2454.	1.8	11