
## Bl Roth

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11696158/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Serotonin-2C and -2a receptor co-expression on cells in the rat medial prefrontal cortex. Neuroscience, 2015, 297, 22-37.	2.3	61
2	The PDZ-binding domain is essential for the dendritic targeting of 5-HT2A serotonin receptors in cortical pyramidal neurons in vitro. Neuroscience, 2003, 122, 907-920.	2.3	71
3	Localization of 5-HT2A receptors on dopamine cells in subnuclei of the midbrain A10 cell group. Neuroscience, 2002, 111, 163-176.	2.3	154
4	Similar ultrastructural distribution of the 5-HT2A serotonin receptor and microtubule-associated protein MAP1A in cortical dendrites of adult rat. Neuroscience, 2002, 113, 23-35.	2.3	89
5	Clozapine and other 5-hydroxytryptamine-2A receptor antagonists alter the subcellular distribution of 5-hydroxytryptamine-2A receptors in vitro and in vivo. Neuroscience, 1999, 91, 599-606.	2.3	153
6	Cross-linking of [1251]β-endorphin to μ-opioid receptors during development. Developmental Brain Research, 1989, 45, 283-289.	1.7	9
7	Phorbol esters inhibit alpha1-adrenergic receptor-stimulated phosphoinositide hydrolysis and contraction in rat aorta: Evidence for a link between vascular contraction and phosphoinositide turnover. Biochemical and Biophysical Research Communications, 1986, 134, 970-974.	2.1	53
8	Detection and characterization of beta-adrenergic receptors and adenylate cyclase in coated vesicles isolated from bovine brain. Journal of Neuroscience, 1986, 6, 2578-2584.	3.6	44
9	Stereospecific opiate-binding sites occur in coated vesicles. Journal of Neuroscience, 1985, 5, 3010-3015.	3.6	36
10	Differential ontogeny of multiple opioid receptors (mu, delta, and kappa). Journal of Neuroscience, 1985, 5, 584-588.	3.6	224
11	Activation of phosphatidylinositol turnover in rat aorta by α1-adrenergic receptor stimulation. European Journal of Pharmacology, 1985, 110, 389-390.	3.5	59
12	Evidence for distinct subcellular sites of opiate receptors. Demonstration of opiate receptors in smooth microsomal fractions isolated from rat brain Journal of Biological Chemistry, 1981, 256, 10117-10123.	3.4	74