

# Hongtao Wang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11697058/publications.pdf>

Version: 2024-02-01

8  
papers

370  
citations

1163117

8  
h-index

1588992

8  
g-index

9  
all docs

9  
docs citations

9  
times ranked

596  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Timing and Location of Glial Cell Line-Derived Neurotrophic Factor Expression Determine Enteric Nervous System Structure and Function. <i>Journal of Neuroscience</i> , 2010, 30, 1523-1538.	3.6	79
2	PAI-1 deficiency reduces liver fibrosis after bile duct ligation in mice through activation of tPA. <i>FEBS Letters</i> , 2007, 581, 3098-3104.	2.8	76
3	LRH-1 mitigates intestinal inflammatory disease by maintaining epithelial homeostasis and cell survival. <i>Nature Communications</i> , 2018, 9, 4055.	12.8	58
4	Transcriptional profiling after bile duct ligation identifies PAI-1 as a contributor to cholestatic injury in mice. <i>Hepatology</i> , 2005, 42, 1099-1108.	7.3	56
5	Hepatocyte Growth Factor and MET Support Mouse Enteric Nervous System Development, the Peristaltic Response, and Intestinal Epithelial Proliferation in Response to Injury. <i>Journal of Neuroscience</i> , 2015, 35, 11543-11558.	3.6	34
6	Tissue-type plasminogen activator deficiency exacerbates cholestatic liver injury in mice. <i>Hepatology</i> , 2007, 45, 1527-1537.	7.3	30
7	Development of the First Low Nanomolar Liver Receptor Homolog-1 Agonist through Structure-guided Design. <i>Journal of Medicinal Chemistry</i> , 2019, 62, 11022-11034.	6.4	21
8	Serum Markers May Distinguish Biliary Atresia From Other Forms of Neonatal Cholestasis. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2010, 50, 411-416.	1.8	15