

# Peter HÃ,gh

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11743042/publications.pdf>

Version: 2024-02-01

47  
papers

1,337  
citations

331670

21  
h-index

377865

34  
g-index

48  
all docs

48  
docs citations

48  
times ranked

2002  
citing authors

#	ARTICLE	IF	CITATIONS
1	Sixteen Weeks of Aerobic Exercise does not Alter Resting-state Connectivity of the Precuneus in Patients with Alzheimer's Disease. <i>Current Alzheimer Research</i> , 2022, 19, 171-177.	1.4	3
2	The Added Value of Cerebrospinal Fluid Neurofilament Light Chain to Existing Diagnostic Methods and Biomarkers in a Mixed Memory Clinic Cohort of Consecutive Patients. <i>Journal of Alzheimer's Disease Reports</i> , 2022, 6, 121-127.	2.2	2
3	Saliva Neurofilament Light Chain Is Not a Diagnostic Biomarker for Neurodegeneration in a Mixed Memory Clinic Population. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 659898.	3.4	9
4	Lactoferrin in cerebrospinal fluid and saliva is not a diagnostic biomarker for Alzheimer's disease in a mixed memory clinic population. <i>EBioMedicine</i> , 2021, 67, 103361.	6.1	23
5	Cerebrospinal fluid and saliva lactoferrin as a diagnostic biomarker for Alzheimer's disease in a mixed memory clinic population. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.8	1
6	A visual rating scale for cingulate island sign on 18F-FDG-PET to differentiate dementia with Lewy bodies and Alzheimer's disease. <i>Journal of the Neurological Sciences</i> , 2020, 410, 116645.	0.6	15
7	The role of physical and cognitive function in performance of activities of daily living in patients with mild-to-moderate Alzheimer's disease – a cross-sectional study. <i>BMC Geriatrics</i> , 2020, 20, 513.	2.7	24
8	Electroencephalographic Cross-Frequency Coupling as a Sign of Disease Progression in Patients With Mild Cognitive Impairment: A Pilot Study. <i>Frontiers in Neuroscience</i> , 2020, 14, 790.	2.8	27
9	Detecting seizure patterns in patients with Alzheimer's disease using long-term EEG monitoring: A feasibility study. <i>Alzheimer's and Dementia</i> , 2020, 16, e042025.	0.8	0
10	Cerebrospinal Fluid Biomarkers to Differentiate Idiopathic Normal Pressure Hydrocephalus from Subcortical Ischemic Vascular Disease. <i>Journal of Alzheimer's Disease</i> , 2020, 75, 937-947.	2.6	13
11	Physical Exercise May Increase Plasma Concentration of High-Density Lipoprotein-Cholesterol in Patients With Alzheimer's Disease. <i>Frontiers in Neuroscience</i> , 2020, 14, 532.	2.8	3
12	Changes in the left temporal microstate are a sign of cognitive decline in patients with Alzheimer's disease. <i>Brain and Behavior</i> , 2020, 10, e01630.	2.2	22
13	Cerebrospinal Fluid/Plasma Albumin Ratio as a Biomarker for Blood-Brain Barrier Impairment Across Neurodegenerative Dementias. <i>Journal of Alzheimer's Disease</i> , 2020, 75, 429-436.	2.6	35
14	Cholinergic dysfunction, neurodegeneration, and amyloid-beta pathology in neurodegenerative diseases. <i>Psychiatry Research - Neuroimaging</i> , 2020, 302, 111099.	1.8	9
15	Oscillatory connectivity as a diagnostic marker of dementia due to Alzheimer's disease. <i>Clinical Neurophysiology</i> , 2019, 130, 1889-1899.	1.5	30
16	Microstates as Disease and Progression Markers in Patients With Mild Cognitive Impairment. <i>Frontiers in Neuroscience</i> , 2019, 13, 563.	2.8	53
17	Brief Assessment of Impaired Cognition (BASIC) – Validation of a new dementia case-finding instrument integrating cognitive assessment with patient and informant report. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 1724-1733.	2.7	14
18	Moderate-to high-intensity exercise does not modify cortical $\beta$ 2-microglobulin in Alzheimer's disease. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2019, 5, 208-215.	3.7	20

#	ARTICLE	IF	CITATIONS
19	Patients with Alzheimer's disease who carry the <i>APOE</i> $\epsilon$ 4 allele benefit more from physical exercise. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2019, 5, 99-106.	3.7	40
20	Exercise as a potential modulator of inflammation in patients with Alzheimer's disease measured in cerebrospinal fluid and plasma. <i>Experimental Gerontology</i> , 2019, 121, 91-98.	2.8	72
21	Altered Low-Frequency EEG Connectivity in Mild Cognitive Impairment as a Sign of Clinical Progression. <i>Journal of Alzheimer's Disease</i> , 2019, 68, 947-960.	2.6	19
22	Quantitative Electroencephalography Analyzed by Statistical Pattern Recognition as a Diagnostic and Prognostic Tool in Mild Cognitive Impairment: Results from a Nordic Multicenter Cohort Study. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2019, 8, 426-438.	1.3	14
23	Progressive DNA and RNA damage from oxidation after aneurysmal subarachnoid haemorrhage in humans. <i>Free Radical Research</i> , 2018, 52, 51-56.	3.3	9
24	A 16-Week Aerobic Exercise Intervention Does Not Affect Hippocampal Volume and Cortical Thickness in Mild to Moderate Alzheimer's Disease. <i>Frontiers in Aging Neuroscience</i> , 2018, 10, 293.	3.4	27
25	Use of Flutemetamol F 18 $\epsilon$ -Labeled Positron Emission Tomography and Other Biomarkers to Assess Risk of Clinical Progression in Patients With Amnesic Mild Cognitive Impairment. <i>JAMA Neurology</i> , 2018, 75, 1114.	9.0	75
26	Change in Fitness and the Relation to Change in Cognition and Neuropsychiatric Symptoms After Aerobic Exercise in Patients with Mild Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 137-145.	2.6	45
27	EEG Theta Power Is an Early Marker of Cognitive Decline in Dementia due to Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2018, 64, 1359-1371.	2.6	100
28	Decreased Parietal Beta Power as a Sign of Disease Progression in Patients with Mild Cognitive Impairment. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 475-487.	2.6	25
29	Effect of physical exercise on markers of neuronal dysfunction in cerebrospinal fluid in patients with Alzheimer's disease. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2017, 3, 284-290.	3.7	23
30	[P3 $\epsilon$ -174]: EFFECT OF PHYSICAL EXERCISE ON MARKERS OF NEURONAL DYSFUNCTION IN CEREBROSPINAL FLUID IN PATIENTS WITH ALZHEIMER'S DISEASE. <i>Alzheimer's and Dementia</i> , 2017, 13, P1000.	0.8	0
31	Moderate-to-High Intensity Physical Exercise in Patients with Alzheimer's Disease: A Randomized Controlled Trial. <i>Journal of Alzheimer's Disease</i> , 2016, 50, 443-453.	2.6	210
32	Electroencephalography Is a Good Complement to Currently Established Dementia Biomarkers. <i>Dementia and Geriatric Cognitive Disorders</i> , 2016, 42, 80-92.	1.5	30
33	Cerebrospinal Fluid Amyloid Beta and Tau Concentrations Are Not Modulated by 16 Weeks of Moderate- to High-Intensity Physical Exercise in Patients with Alzheimer Disease. <i>Dementia and Geriatric Cognitive Disorders</i> , 2016, 42, 146-158.	1.5	40
34	Effect of aerobic exercise on physical performance in patients with Alzheimer's disease. <i>Alzheimer's and Dementia</i> , 2016, 12, 1207-1215.	0.8	76
35	Associations between physical function, dual-task performance and cognition in patients with mild Alzheimer's disease. <i>Aging and Mental Health</i> , 2016, 20, 1139-1146.	2.8	28
36	P1-069: Moderate-to-high intensity physical training does not alter cerebrospinal amyloid- $\beta$ 1-42 levels in patients with Alzheimer's disease. , 2015, 11, P364-P365.		0

#	ARTICLE	IF	CITATIONS
37	Autonomic Dysfunction in Patients with Mild to Moderate Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2015, 47, 681-689.	2.6	31
38	O5-04-06: Moderate to high-intensity physical exercise in patients with Alzheimer's disease. , 2015, 11, P324-P325.		2
39	Discrepancy between stimulus response and tolerance of pain in Alzheimer disease. <i>Neurology</i> , 2015, 84, 1575-1581.	1.1	25
40	P4-059: CEREBROSPINAL FLUID ROUTINE PARAMETERS AND BIOMARKERS AS POTENTIAL INDICATORS OF CLINICAL PROGRESSION IN MILD COGNITIVE IMPAIRMENT. , 2014, 10, P803-P804.		0
41	Visual Rating and ROI-Based Parametric Analysis of rCBF SPECT in Patients with Mild or Questionable Dementia: A Comparative Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2007, 24, 429-433.	1.5	10
42	Temporal Lobe Hypoperfusion in Isolated Amnesia with Slow Onset: A Single Photon Emission Computer Tomography Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2004, 18, 15-23.	1.5	15
43	Diagnostic profile of young and middle-aged memory clinic patients. <i>Neurology</i> , 2002, 59, 1259-1262.	1.1	35
44	Single Photon Emission Computed Tomography and Apolipoprotein E in Alzheimer's Disease: Impact of the $\epsilon 4$ Allele on Regional Cerebral Blood Flow. <i>Journal of Geriatric Psychiatry and Neurology</i> , 2001, 14, 42-51.	2.3	27
45	A multidisciplinary memory clinic in a neurological setting: diagnostic evaluation of 400 consecutive patients. <i>European Journal of Neurology</i> , 1999, 6, 279-288.	3.3	37
46	Functional Brain Imaging With Single-Photon Emission Computed Tomography in the Diagnosis of Alzheimer's Disease. <i>International Psychogeriatrics</i> , 1997, 9, 223-227.	1.0	19
47	Potentially Reversible Conditions in Memory Clinic Patients. , 0, , 123-128.		0