

# Abdalla Rifai

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11758250/publications.pdf>

Version: 2024-02-01

31  
papers

1,110  
citations

430874

18  
h-index

477307

29  
g-index

31  
all docs

31  
docs citations

31  
times ranked

1073  
citing authors

#	ARTICLE	IF	CITATIONS
1	Candesartan suppresses chronic renal inflammation by a novel antioxidant action independent of AT1R blockade. <i>Kidney International</i> , 2008, 74, 1128-1138.	5.2	74
2	Hepatocyte Growth Factor Suppresses Proinflammatory NF $\kappa$ B Activation through GSK3 $\beta$ Inactivation in Renal Tubular Epithelial Cells. <i>Journal of Biological Chemistry</i> , 2008, 283, 7401-7410.	3.4	89
3	Anti-Inflammatory Effect of Hepatocyte Growth Factor in Chronic Kidney Disease: Targeting the Inflamed Vascular Endothelium. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 2464-2473.	6.1	83
4	Platelet derived growth factor-D may be a possible therapeutic target for advanced IgA nephropathy. <i>Nephrology</i> , 2005, 10, A439-A439.	1.6	0
5	Activation of PI3K $\rightarrow$ Akt $\rightarrow$ GSK3 $\beta$ pathway mediates hepatocyte growth factor inhibition of RANTES expression in renal tubular epithelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2005, 330, 27-33.	2.1	35
6	Hepatocyte Growth Factor Ameliorates Renal Interstitial Inflammation in Rat Remnant Kidney by Modulating Tubular Expression of Macrophage Chemoattractant Protein-1 and RANTES. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 2868-2881.	6.1	99
7	Hepatocyte growth factor ameliorates progression of interstitial fibrosis in rats with established renal injury. <i>Kidney International</i> , 2004, 65, 409-419.	5.2	66
8	Hepatocyte Growth Factor Modulates Matrix Metalloproteinases and Plasminogen Activator/Plasmin Proteolytic Pathways in Progressive Renal Interstitial Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2003, 14, 3047-3060.	6.1	88
9	Profiling the IgA nephropathy renal transcriptome: analysis by complementary DNA array hybridization. <i>Nephrology</i> , 2002, 7, S140-S144.	1.6	2
10	IgA nephropathy and mesangial cell proliferation: shared global gene expression profiles. <i>Nephrology</i> , 2002, 7, S106-S113.	1.6	2
11	IgA nephropathy and mesangial cell proliferation: shared global gene expression profiles. <i>Nephrology</i> , 2002, 7, S106.	1.6	1
12	Profiling the IgA nephropathy renal transcriptome: analysis by complementary DNA array hybridization. <i>Nephrology</i> , 2002, 7, S140.	1.6	2
13	Profiling the adult human liver transcriptome: analysis by cDNA array hybridization. <i>Journal of Hepatology</i> , 2001, 35, 178-186.	3.7	35
14	Comprehensive gene expression profile of the adult human renal cortex: Analysis by cDNA array hybridization. <i>Kidney International</i> , 2000, 57, 1452-1459.	5.2	32
15	Genomic repertoire of human mesangial cells: comprehensive analysis of gene expression by cDNA array hybridization. <i>Nephrology</i> , 2000, 5, 215-223.	1.6	8
16	The N-Glycans Determine the Differential Blood Clearance and Hepatic Uptake of Human Immunoglobulin (Ig) $\alpha$ 1 and Ig $\alpha$ 2 Isotypes. <i>Journal of Experimental Medicine</i> , 2000, 191, 2171-2182.	8.5	144
17	Characterization and Evaluation of Detoxification Functions of a Nontumorigenic Immortalized Porcine Hepatocyte Cell Line (HepLiu). <i>Cell Transplantation</i> , 1999, 8, 219-232.	2.5	37
18	Phenotypic characterization of cytokine expression in patients with IgA nephropathy. <i>Journal of Clinical Immunology</i> , 1997, 17, 396-402.	3.8	59

#	ARTICLE	IF	CITATIONS
19	Increase of HLA-DR-positive natural killer cells in peripheral blood from patients with IgA nephropathy. <i>Human Immunology</i> , 1996, 49, 64-70.	2.4	19
20	Immunopathogenesis of experimental IgA nephropathy. <i>Seminars in Immunopathology</i> , 1994, 16, 81-95.	4.0	1
21	Extrarenal cytokines modulate the glomerular response to IgA immune complexes. <i>Kidney International</i> , 1992, 42, 341-353.	5.2	65
22	Pathogenesis of IgA Immune Complex-Mediated Glomerulonephritis. , 1991, , 996-1003.		1
23	Characteristics of Nephritogenic IgA Immune Complexes. <i>American Journal of Kidney Diseases</i> , 1988, 12, 402-405.	1.9	11
24	Composition of IgA immune complexes precipitated with polyethylene glycol. <i>Journal of Immunological Methods</i> , 1987, 103, 239-245.	1.4	21
25	Experimental models for IgA-associated nephritis. <i>Kidney International</i> , 1987, 31, 1-7.	5.2	44
26	Complement activation in experimental IgA nephropathy: An antigen-mediated process. <i>Kidney International</i> , 1987, 32, 838-844.	5.2	41
27	Preparation of phosphorylcholine-conjugated antigens. <i>Journal of Immunological Methods</i> , 1986, 94, 25-30.	1.4	18
28	IgA Immune Complexes and Disease: An Experimental Perspective. <i>Pathology and Immunopathology Research</i> , 1986, 5, 278-285.	0.8	1
29	Experimental IgA Nephropathy <sup>1</sup> . <i>Contributions To Nephrology</i> , 1984, 40, 37-44.	1.1	11
30	THE CLEARANCE KINETICS AND HEPATIC LOCALIZATION OF IgA-IMMUNE COMPLEXES IN MICE. <i>Annals of the New York Academy of Sciences</i> , 1983, 409, 861-863.	3.8	2
31	Stable, soluble, model immune complexes made with a versatile multivalent affinity-labeling antigen. <i>Biochemistry</i> , 1982, 21, 301-308.	2.5	19