Ulrike Philippar

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11772334/publications.pdf Version: 2024-02-01



Πισικε Ομιιισσλα

#	Article	IF	CITATIONS
1	Discovery of Highly Potent and Selective IRAK1 Degraders to Probe Scaffolding Functions of IRAK1 in ABC DLBCL. Journal of Medicinal Chemistry, 2021, 64, 10878-10889.	6.4	12
2	Assessing IRAK4 Functions in ABC DLBCL by IRAK4 Kinase Inhibition and Protein Degradation. Cell Chemical Biology, 2020, 27, 1500-1509.e13.	5.2	31
3	Influence of the bispecific antibody IgG subclass on T cell redirection. MAbs, 2019, 11, 1012-1024.	5.2	11
4	Specific covalent inhibition of MALT1 paracaspase suppresses B cell lymphoma growth. Journal of Clinical Investigation, 2018, 128, 4397-4412.	8.2	51
5	Discovery of 1-(1 <i>H</i> -Pyrazolo[4,3- <i>c</i>]pyridin-6-yl)urea Inhibitors of Extracellular Signal-Regulated Kinase (ERK) for the Treatment of Cancers. Journal of Medicinal Chemistry, 2016, 59, 6501-6511.	6.4	26
6	PTP1B-dependent regulation of receptor tyrosine kinase signaling by the actin-binding protein Mena. Molecular Biology of the Cell, 2015, 26, 3867-3878.	2.1	31
7	Discovery of a Novel ERK Inhibitor with Activity in Models of Acquired Resistance to BRAF and MEK Inhibitors. Cancer Discovery, 2013, 3, 742-750.	9.4	559
8	Predictive Genes in Adjacent Normal Tissue Are Preferentially Altered by sCNV during Tumorigenesis in Liver Cancer and May Rate Limiting. PLoS ONE, 2011, 6, e20090.	2.5	68
9	DLK1-DIO3 Genomic Imprinted MicroRNA Cluster at 14q32.2 Defines a Stemlike Subtype of Hepatocellular Carcinoma Associated with Poor Survival. Journal of Biological Chemistry, 2011, 286, 30706-30713.	3.4	147
10	Gene Signatures Derived from a c-MET-Driven Liver Cancer Mouse Model Predict Survival of Patients with Hepatocellular Carcinoma. PLoS ONE, 2011, 6, e24582.	2.5	26
11	Degeneration of the mouse retina upon dysregulated activity of serum response factor. Molecular Vision, 2011, 17, 1110-27.	1.1	7
12	Identification of invasion specific splice variants of the cytoskeletal protein Mena present in mammary tumor cells during invasion inÂvivo. Clinical and Experimental Metastasis, 2009, 26, 153-159.	3.3	107
13	A Mena Invasion Isoform Potentiates EGF-Induced Carcinoma Cell Invasion and Metastasis. Developmental Cell, 2008, 15, 813-828.	7.0	242
14	Human Mena+11a Isoform Serves as a Marker of Epithelial Phenotype and Sensitivity to Epidermal Growth Factor Receptor Inhibition in Human Pancreatic Cancer Cell Lines. Clinical Cancer Research, 2008, 14, 4943-4950.	7.0	63
15	Ena/VASP Is Required for Neuritogenesis in the Developing Cortex. Neuron, 2007, 56, 441-455.	8.1	204
16	Filopodia are required for cortical neurite initiation. Nature Cell Biology, 2007, 9, 1347-1359.	10.3	276
17	Simultaneous imaging of GFP, CFP and collagen in tumors in vivo using multiphoton microscopy. BMC Biotechnology, 2005, 5, 14.	3.3	119
18	Neuronal migration in the murine rostral migratory stream requires serum response factor. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 6148-6153.	7.1	131

ULRIKE PHILIPPAR

#	Article	IF	CITATIONS
19	SRF regulates Bcl-2 expression and promotes cell survival during murine embryonic development. EMBO Journal, 2004, 23, 1834-1844.	7.8	67
20	The SRF Target Gene Fhl2 Antagonizes RhoA/MAL-Dependent Activation of SRF. Molecular Cell, 2004, 16, 867-880.	9.7	137
21	Serum response factor is crucial for actin cytoskeletal organization and focal adhesion assembly in embryonic stem cells. Journal of Cell Biology, 2002, 156, 737-750.	5.2	176