

# Jeremiah Morrissey

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11783082/publications.pdf>

Version: 2024-02-01

22  
papers

2,936  
citations

361413

20  
h-index

677142

22  
g-index

22  
all docs

22  
docs citations

22  
times ranked

2515  
citing authors

#	ARTICLE	IF	CITATIONS
1	Myocardial effects of VDR activators in renal failure. Journal of Steroid Biochemistry and Molecular Biology, 2010, 121, 188-192.	2.5	52
2	Combination Therapy with an Angiotensin-Converting Enzyme Inhibitor and a Vitamin D Analog Suppresses the Progression of Renal Insufficiency in Uremic Rats. Journal of the American Society of Nephrology: JASN, 2007, 18, 1796-1806.	6.1	186
3	Bone morphogenic protein-7 (BMP-7), a novel therapy for diabetic nephropathy <sup>11</sup> Professor Robert Chevalier served as a guest editor for this paper.. Kidney International, 2003, 63, 2037-2049.	5.2	209
4	Obstructive nephropathy and renal fibrosis: The role of bone morphogenic protein-7 and hepatocyte growth factor. Kidney International, 2003, 64, S105-S112.	5.2	75
5	ACE inhibition increases expression of the ETB receptor in kidneys of mice with unilateral obstruction. American Journal of Physiology - Renal Physiology, 2003, 284, F209-F217.	2.7	31
6	Transforming Growth Factor- $\beta^2$ Induces Renal Epithelial Jagged-1 Expression in Fibrotic Disease. Journal of the American Society of Nephrology: JASN, 2002, 13, 1499-1508.	6.1	130
7	Obstructive nephropathy and renal fibrosis. American Journal of Physiology - Renal Physiology, 2002, 283, F861-F875.	2.7	500
8	Comparative effects of ACE inhibition and angiotensin II receptor blockade in the prevention of renal damage. Kidney International, 2002, 62, S23-S26.	5.2	20
9	Bone Morphogenetic Protein-7 Improves Renal Fibrosis and Accelerates the Return of Renal Function. Journal of the American Society of Nephrology: JASN, 2002, 13, S14-S21.	6.1	168
10	Contributions of angiotensin II and tumor necrosis factor- $\alpha$ to the development of renal fibrosis. American Journal of Physiology - Renal Physiology, 2001, 280, F777-F785.	2.7	122
11	Osteogenic protein-1 prevents renal fibrogenesis associated with ureteral obstruction. American Journal of Physiology - Renal Physiology, 2000, 279, F130-F143.	2.7	215
12	Induction of CD14 in Tubular Epithelial Cells During Kidney Disease. Journal of the American Society of Nephrology: JASN, 2000, 11, 1681-1690.	6.1	30
13	Role of TNFR1 and TNFR2 receptors in tubulointerstitial fibrosis of obstructive nephropathy. American Journal of Physiology - Renal Physiology, 1999, 277, F766-F772.	2.7	89
14	Delayed treatment with enalapril halts tubulointerstitial fibrosis in rats with obstructive nephropathy. Kidney International, 1996, 49, 1110-1119.	5.2	139
15	Angiotensin II receptor antagonist ameliorates renal tubulointerstitial fibrosis caused by unilateral ureteral obstruction. Kidney International, 1995, 47, 1285-1294.	5.2	316
16	Role of angiotensin II in the tubulointerstitial fibrosis of obstructive nephropathy. American Journal of Kidney Diseases, 1995, 26, 141-146.	1.9	93
17	Enalapril reduces collagen type IV synthesis and expansion of the interstitium in the obstructed rat kidney. Kidney International, 1994, 45, 1637-1647.	5.2	201
18	Effects of high concentrations of glucose on PTH secretion in parathyroid cells. Kidney International, 1990, 37, 1522-1527.	5.2	104

#	ARTICLE	IF	CITATIONS
19	Eicosanoid production by isolated glomeruli of rats with unilateral ureteral obstruction. <i>Kidney International</i> , 1990, 37, 1528-1535.	5.2	41
20	Effect of 1,25-Dihydroxyvitamin D <sub>3</sub> on Phospholipid Metabolism in Cultured Bovine Parathyroid Cells*. <i>Endocrinology</i> , 1988, 122, 2387-2392.	2.8	16
21	Suppression of parathyroid hormone secretion by aluminum. <i>Kidney International</i> , 1983, 23, 699-704.	5.2	178
22	The Role of Na <sup>+</sup> -Ca <sup>++</sup> Exchange in Parathyroid Hormone Secretion*. <i>Endocrinology</i> , 1982, 111, 225-230.	2.8	21