
Ricardo Alberto Moreno

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11790692/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Physical activity as an adjuvant therapy for depression and influence on peripheral inflammatory markers: A randomized clinical trial. Mental Health and Physical Activity, 2022, 22, 100442.	1.8	2
2	Treatment of mixed depression with theta-burst stimulation (TBS): results from a double-blind, randomized, sham-controlled clinical trial. Neuropsychopharmacology, 2021, 46, 2257-2265.	5.4	10
3	Altered brain creatine cycle metabolites in bipolar I disorder with childhood abuse: A 1H magnetic resonance spectroscopy study. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2021, 109, 110233.	4.8	2
4	Efficacy, Safety, and Tolerability of Theta-Burst Stimulation in Mixed Depression: Design, Rationale, and Objectives of a Randomized, Double-Blinded, Sham-Controlled Trial. Frontiers in Psychiatry, 2020, 11, 435.	2.6	2
5	A randomized controlled trial comparing lithium plus valproic acid versus lithium plus carbamazepine in young patients with type 1 bipolar disorder: the LICAVAL study. Trials, 2019, 20, 608.	1.6	9
6	Anterior Cingulate Cortex Glutamatergic Metabolites and Mood Stabilizers in Euthymic Bipolar I Disorder Patients: A Proton Magnetic Resonance Spectroscopy Study. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, 2018, 3, 985-991.	1.5	19
7	What is the nonverbal communication of depression? Assessing expressive differences between depressive patients and healthy volunteers during clinical interviews. Journal of Affective Disorders, 2018, 238, 636-644.	4.1	14
8	Lithium-associated anterior cingulate neurometabolic profile in euthymic Bipolar I disorder: A 1H-MRS study. Journal of Affective Disorders, 2018, 241, 192-199.	4.1	18
9	Is nonverbal behavior in patients and interviewers relevant to the assessment of depression and its recovery? A study with Dutch and Brazilian patients. Psychiatry Research, 2017, 250, 59-64.	3.3	8
10	Investigation of associations between recurrence of major depressive disorder and spinal posture alignment: A quantitative cross-sectional study. Gait and Posture, 2017, 52, 258-264.	1.4	20
11	Predicting response to treatment and discriminating bipolar and depression symptoms using Hamilton Depression Rating Scale. Jornal Brasileiro De Psiquiatria, 2017, 66, 125-130.	0.7	2
12	Antidepressant Efficacy of Adjunctive Aerobic Activity and Associated Biomarkers in Major Depression: A 4-Week, Randomized, Single-Blind, Controlled Clinical Trial. PLoS ONE, 2016, 11, e0154195.	2.5	40
13	Hamilton depression rating scale and montgomery–asberg depression rating scale in depressed and bipolar I patients: psychometric properties in a Brazilian sample. Health and Quality of Life Outcomes, 2015, 13, 42.	2.4	58
14	Bcl-2 rs956572 Polymorphism is Associated with Increased Anterior Cingulate Cortical Glutamate in Euthymic Bipolar I Disorder. Neuropsychopharmacology, 2013, 38, 468-475.	5.4	65
15	Gender effects of the COMT Val158Met genotype on verbal fluency in healthy adults. Molecular Medicine Reports, 2013, 8, 837-844.	2.4	18
16	Burden of maternal bipolar disorder on at-risk offspring: A controlled study on family planning and maternal care. Journal of Affective Disorders, 2012, 143, 172-178.	4.1	13
17	COMT polymorphisms as predictors of cognitive dysfunction during manic and mixed episodes in bipolar I disorder. Bipolar Disorders, 2012, 14, 554-564.	1.9	46
18	Posture and body image in individuals with major depressive disorder: a controlled study. Revista Brasileira De Psiquiatria, 2010, 32, 375-380.	1.7	73

#	Article	IF	CITATIONS
19	The prevalence of psychopathology in offspring of bipolar women from a Brazilian tertiary center. Revista Brasileira De Psiquiatria, 2009, 31, 240-246.	1.7	26